

# Accepted Manuscript

Synthesis and antiproliferative activity of 6,7-disubstituted-4-phenoxyquinoline derivatives bearing the 1,8-naphthyridin-2-one moiety

Qidong Tang, Yongli Duan, Hehua Xiong, Ting Chen, Zhen Xiao, Linxiao Wang, Yueyue Xiao, Shunmin Huang, Yinhua Xiong, Wufu Zhu, Ping Gong, Pengwu Zheng



PII: S0223-5234(18)30738-4

DOI: [10.1016/j.ejmech.2018.08.066](https://doi.org/10.1016/j.ejmech.2018.08.066)

Reference: EJMECH 10677

To appear in: *European Journal of Medicinal Chemistry*

Received Date: 30 April 2018


Revised Date: 23 August 2018

Accepted Date: 24 August 2018

Please cite this article as: Q. Tang, Y. Duan, H. Xiong, T. Chen, Z. Xiao, L. Wang, Y. Xiao, S. Huang, Y. Xiong, W. Zhu, P. Gong, P. Zheng, Synthesis and antiproliferative activity of 6,7-disubstituted-4-phenoxyquinoline derivatives bearing the 1,8-naphthyridin-2-one moiety, *European Journal of Medicinal Chemistry* (2018), doi: 10.1016/j.ejmech.2018.08.066.

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

## Graphical abstract



A series of 6,7-disubstituted-4-phenoxyquinoline derivatives bearing the 1,8-naphthyridin-2-one moiety (**25–56**) were designed, synthesized and evaluated for their activity against c-Met kinase and cancer cell lines. The most promising compound **33** showed excellent *in vitro* activity.

Download English Version:

<https://daneshyari.com/en/article/10147447>

Download Persian Version:

<https://daneshyari.com/article/10147447>

[Daneshyari.com](https://daneshyari.com)