



ELSEVIER

Contents lists available at ScienceDirect

Comptes Rendus Chimie

www.sciencedirect.com



Full paper/Mémoire

A practical and highly efficient transfer hydrogenation of aryl azides using a $[\text{Ru}(p\text{-cymene})\text{Cl}_2]_2$ catalyst and sodium borohydride

Benan Kilbas^{a, b, *}, Yunus Emre Yilmaz^a, Sinem Ergen^a^a Department of Chemistry, Faculty of Sciences, Duzce University, 81620 Duzce, Turkey^b Moltek A. S. Gebze Organize Sanayi, 41400 Gebze, Kocaeli, Turkey

ARTICLE INFO

Article history:

Received 22 May 2018

Accepted 17 July 2018

Available online xxxxx

Keywords:

 $[\text{Ru}(p\text{-cymene})\text{Cl}_2]_2$

Homogeneous catalyst

Transfer hydrogenation

Aryl azide

ABSTRACT

Various aniline derivatives were synthesized by selective reduction of aryl azides in the presence of a dichloro(*p*-cymene)ruthenium(II) dimer ($[\text{Ru}(p\text{-cymene})\text{Cl}_2]_2$) via hydrolysis of sodium borohydride. The hydrogenation reactions were carried out in aqueous media at room temperature. Most of the reactions were completed within 10 min with quantitative yields.

© 2018 Académie des sciences. Published by Elsevier Masson SAS. All rights reserved.

1. Introduction

Reduction of azidoarenes to the aniline derivatives is an industrially important process, since the aniline derivatives are crucial intermediates for the synthesis of dyes, polymers, pharmaceuticals, agrochemicals and biologically active molecules [1–5]. Various synthetic approaches for the reduction of azidoarenes proceed through catalytic hydrogenation [6,7], NaBH_4 /phase-transfer catalysis [8], triphenylphosphine [9], $\text{BF}_3 \cdot \text{OEt}_2/\text{EtSH}$ [10], metal mediated reductions such as $\text{NiCl}_2\text{--Zn}$ [11], $\text{FeCl}_3\text{--Zn}$ [12], Sm/I_2 [13] and tellurium metal [14]. These traditional methods have some disadvantages such as high temperature, high pressure, long reaction time and tedious work-up process. Furthermore, desired transformation is more complicated by using polyfunctional azide substrates in which poor chemoselectivity can be observed. Transfer Hydrogenation (TH), hydrogen abstraction from the reagent by catalyst contribution followed by hydrogen addition to the

unsaturated functional system, is a more convenient method for the reduction of diverse functional groups bearing mainly carbonyls and imines [15,16]. This method has several important advantages such as lack of explosive free hydrogen, chemoselectivity, inexpensive and readily available hydrogen sources and also the use of readily accessible, recyclable and reusable catalysts [17–19]. A plenty of homogeneous and heterogeneous catalysts have been already reported in the literature for the TH process. Although heterogeneous catalysts are recyclable and reusable, homogeneous catalysts are usually preferred for TH due to their more activity and selectivity. Recently, considerable attention has been focused on arene ruthenium complexes as TH catalysts due to their unique properties, inherent catalytic ability and versatile building blocks for coordination cages and macrocycles [20]. Arene rings provided sterical hindrance of the ruthenium center, which prevent rapid oxidation, and they are also resistant to the substitution reactions. Besides, there are three available coordination sites located on the opposite side of the arene ligand can be easily replaced by donor ligands to employ more catalytically active Ru(II) complexes [21,22].

* Corresponding author.

E-mail address: bkilbas@gmail.com (B. Kilbas).

Although exponential growth has been observed for TH of carbonyl and imine polar bonds by the aid of arene ruthenium (II) complexes, the usage of these catalysts for azide hydrogenation is rare.


Herein we report for the first time that various azido arenes were catalytically converted to aniline derivatives by the contribution of the $[\text{Ru}(p\text{-cymene})\text{Cl}_2]_2$ [23] dimeric structure and sodiumborohydride in the aqueous medium with shorter reaction times and high yields.

2. Results and discussion

Monomeric piano-stool type arene ruthenium (II) complexes are generally derived by the cleavage of the $[\text{Ru}(p\text{-cymene})\text{Cl}_2]_2$ dimeric structure with $-\text{N}$, $-\text{S}$, $-\text{P}$ type strong donor ligands to provide efficient performance and catalytic activity for TH [16]. Recently, Cetinkaya et al. described novel monomeric 3,4-dihydroquinazoline ruthenium(II) complexes for the reduction of acetophenone to the corresponding secondary alcohol with high yield after 1 h [24]. They claimed that only 15% conversion was observed when $[\text{Ru}(p\text{-cymene})\text{Cl}_2]_2$ was used as a catalyst.

As it is known that, azido compounds are more reactive, therefore we initially utilized the $[\text{Ru}(p\text{-cymene})\text{Cl}_2]_2$ dimer to check the completion of the reduction process. The careful GC analysis indicated that absolute conversions of azidoarenes to corresponding anilines were achieved within 10 min at room temperature. These results demonstrate that there is no need for an extra reaction step for the generation of a new catalyst to afford reduction. Furthermore, $[\text{Ru}(p\text{-cymene})\text{Cl}_2]_2$ is a promising candidate for the chemoselective TH as previously published monomeric arene ruthenium (II) complexes. On the other hand, a hydrogen source played another crucial role in TH. Apparently, a compound which has lower oxidation potential can act as a versatile hydrogen carrier, because these unique properties enable hydrogen transfer from the donor to the substrate in the presence of the catalyst. Alcohols, formic acid, hydrazine and 1,4-dihydropyridine derivatives have been extensively used as hydrogen carriers [25]. Even they are more accessible, eco-friendly and safe, TH requires harsh reaction conditions such as high temperature and long reaction time. The nature of the hydrogen carriers also influences the chemoselectivity and activity of TH [26,19c].

Table 1
 $[\text{Ru}(p\text{-cymene})\text{Cl}_2]_2$ catalyzed reduction of various $\text{R}-\text{N}_3$ compounds.^a



Entry	R-N ₃	Product	Conversion (%)	Yield ^b (%)
1	Phenyl-	Aniline	100	>95
2	o-Chlorophenyl-	o-Chloroaniline	100	>95
3	p-Chlorophenyl-	p-Chloroaniline	100	>95
4	o-Bromophenyl-	o-Bromoaniline	100	>95
5	p-Bromophenyl-	p-Bromoaniline	100	>95
6	p-Iodophenyl-	p-Iodoaniline	100	>95
7	o-Tolyl-	o-Toluidine	100	>95
8	m-Tolyl-	m-Toluidine	100	>95
9	p-Tolyl-	p-Toluidine	100	>95
10	o-Methoxyphenyl-	o-Methoxyaniline	100	>95
11	m-Methoxyphenyl-	m-Methoxyaniline	100	>95
12	p-Methoxyphenyl-	p-Methoxyaniline	100	>95
13	o-(Trifluoromethyl)phenyl-	o-(Trifluoromethyl)aniline	100	>95
14	p-(Trifluoromethyl)phenyl-	p-(Trifluoromethyl)aniline	100	>95

^a Reaction conditions, substrate (0.25 mmol), NaBH_4 (0.5 mmol) and $[\text{Ru}(p\text{-cymene})\text{Cl}_2]_2$ (10 mg) was used with 1.5 ml of water/methanol ($v/v=2/1$) at room temperature.

^b Isolated yield.

Table 2
Use of different commercially available catalysts for the reduction of 1-azido-2-chlorobenzene to 2-chloroaniline.^a

Entry	Catalyst	Amount of catalyst, mg	Time (min)	Yield ^c (%)	TON ($\frac{\text{mol}_{\text{product}}}{\text{mol}_{\text{metal(s)}}}$)	TOF ($\frac{\text{mol}_{\text{product}}}{\text{mol}_{\text{metal(s)}}/\text{h}}$)
1	$\text{Pd}/\text{C}^{\text{b}}$	3.4	50	70	109.3	131.2
2	$\text{Pd}(\text{OAc})_2$	6.7	60	72	6.0	6.0
3	PdCl_2	5.3	60	35	2.9	2.9
4	$[\text{Ru}(p\text{-cymene})\text{Cl}_2]_2$	10.0	10	>95	28.8	172.7

^a Reaction conditions, substrate (0.25 mmol), NaBH_4 (0.5 mmol) and catalyst (0.03 mmol metal content) was used with 1.5 ml of water/methanol ($v/v=2/1$) at room temperature.

^b 5 wt % Pd.

^c isolated yield.

Download English Version:

<https://daneshyari.com/en/article/10150302>

Download Persian Version:

<https://daneshyari.com/article/10150302>

[Daneshyari.com](https://daneshyari.com)