

Accepted Manuscript

Title: Detection of pacific ciguatoxins using liquid chromatography coupled to either low or high resolution mass spectrometry (LC-MS/MS)

Authors: Manoella Sibat, Christine Herrenknecht, Hélène Taiana Darius, Mélanie Roué, Mireille Chinain, Philipp Hess



PII: S0021-9673(18)30992-0

DOI: <https://doi.org/10.1016/j.chroma.2018.08.008>

Reference: CHROMA 359604

To appear in: *Journal of Chromatography A*

Received date: 6-4-2018

Revised date: 12-7-2018

Accepted date: 3-8-2018

Please cite this article as: Sibat M, Herrenknecht C, Darius HT, Roué M, Chinain M, Hess P, Detection of pacific ciguatoxins using liquid chromatography coupled to either low or high resolution mass spectrometry (LC-MS/MS), *Journal of Chromatography A* (2018), <https://doi.org/10.1016/j.chroma.2018.08.008>

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

Detection of pacific ciguatoxins using liquid chromatography coupled to either low or high resolution mass spectrometry (LC-MS/MS)

Manoella Sibat^{a*}, Christine Herrenknecht^b, Hélène Taiana Darius^c, Mélanie Roué^d, Mireille Chinain^c, Philipp Hess^a

^a Ifremer, Laboratoire Phycotoxines, Rue de l'Ile d'Yeu, 44311 Nantes, France;
manoella.sibat@ifremer.fr, philipp.hess@ifremer.fr,

^b LUNAM, Université de Nantes, MMS EA2160, Faculté de Pharmacie, 9 rue Bias, 44035 Nantes, France, christine.herrenknecht@univ-nantes.fr

^c Institut Louis Malardé (ILM), Laboratory of Toxic Microalgae - UMR 241-EIO, PO box 30, 98713 Papeete, Tahiti, French Polynesia; tdarius@ilm.pf, mchinain@ilm.pf

^d Institut de Recherche pour le Développement (IRD) - UMR 241-EIO, PO box 53267, 98716 Pirae, Tahiti, French Polynesia; melanie.roue@ird.fr

*Corresponding author at: Ifremer/Laboratoire phycotoxines, Rue de l'Ile D'Yeu, 44311 Nantes, France.
Tel.: +33 (2) 40 37 42 54. Fax: +33 (2) 02 40 37 42 41 E-mail address: manoella.sibat@ifremer.fr

Highlights

- MeOH improved selectivity and sensitivity for P-CTXs over ACN-based mobile phase
- HRMS fragmentation allowed for development of selective LRMS method
- LRMS method using $[M+NH_4]^+$ precursor gave better selectivity & less matrix effects
- Low LODs thanks to methanol-based mobile phase and $[M+NH_4]^+$ precursor ion
- HRMS 2- to 12-fold higher LODs but confirmed P-CTXs and eliminated artefacts

Download English Version:

<https://daneshyari.com/en/article/10154454>

Download Persian Version:

<https://daneshyari.com/article/10154454>

[Daneshyari.com](https://daneshyari.com)