




King Saud University  
Arabian Journal of Chemistry

www.ksu.edu.sa  
www.sciencedirect.com



# Molecules and functions of rosewood: *Diospyros celebica*

Juntao Chen<sup>b,1</sup>, Changyu Ni<sup>b,1</sup>, Junwei Lou<sup>c,1</sup>, Wanxi Peng<sup>a,b,\*</sup>

<sup>a</sup> School of Forestry, Henan Agricultural University, Zhengzhou 450002, China

<sup>b</sup> School of Materials Science and Engineering, Central South University of Forestry and Technology, Changsha 410004, China

<sup>c</sup> Department of Architectural Engineering, University of Zhejiang Business Technology Institute, Ningbo 315012, China

Received 3 September 2017; accepted 24 December 2017

## KEYWORDS

Pterocarpus;  
*Diospyros celebica*;  
PY–GC–MS;  
GC–MS;  
TDS–GC–MS;  
Health care ingredients

**Abstract** Pterocarpus and Pterocarpus products have certain human health function. In this paper, *Diospyros celebica* Bakh as an example, we study its human health components by using PY–GC–MS, TDS–GC–MS and GC–MS. The composition of known human health functions was studied by reviewing the literature. 3-O-Methyl-d-glucose has a certain conservation property, and it can protect the pancreatic B cells against the toxicity of alloxan. P-Cresol plays a role in endothelial dysfunction in uremic patients, and it can repair wounds and reduce endothelial progression. 2(3H)-Furanone, 5-methyl-has certain biological resistance, and has high antimicrobial activity against NCIM 2501 and NCIM 5021.

© 2017 Production and hosting by Elsevier B.V. on behalf of King Saud University. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

## 1. Introduction

*Diospyros celebica* Bakh is mainly grown in Sulawesi, Indonesia, belonging to Ebenaceae, Diospyros. *Diospyros celebica* Bakh is the best of the *Diospyros* ebum Koenig, and the paradise bird, the red dragon fish collectively known as the three national treasures of Indonesia, known as “the black pearl in

the wood”. *Diospyros celebica* Bakh grows mainly in the annual drought, rare rain cliff; Because of its poor growth environment, *Diospyros celebica* Bakh has a long period of material, generally take hundreds or even thousands of years. *Diospyros celebica* Bakh is evergreen broad-leaved trees, with tree height up to 40 m and DBH up to 1 m. *Diospyros celebica* Bakh wood for the hollow material, heartwood and sapwood are significantly different, heartwood was black, the surface with shiny and deep and white stripes. Wood with high strength, big hardness, corrosion resistance and strong insect resistance, air dry density of 1.09 g/cm<sup>3</sup>. *Diospyros celebica* Bakh can emit fungal and antibiotics and other natural molecules and it enter the body to inhibit and kill the body of native viruses and bacteria, with human health function. In this paper, the *Diospyros celebica* Bakh powder was analyzed by PY–GC–MS, TDS–GC–MS, TG and FT-IR; The extracts of ethanol, ethanol/benzene and ethanol/methanol in

\* Corresponding author at: School of Forestry, Henan Agricultural University, Zhengzhou 450002, China.

E-mail address: [pengwanxi@163.com](mailto:pengwanxi@163.com) (W. Peng).

<sup>1</sup> Juntao Chen and Changyu Ni are co-first authors.

Peer review under responsibility of King Saud University.



Production and hosting by Elsevier

<https://doi.org/10.1016/j.arabjc.2017.12.033>

1878-5352 © 2017 Production and hosting by Elsevier B.V. on behalf of King Saud University.

This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

Please cite this article in press as: Chen, J. et al., Molecules and functions of rosewood: *Diospyros celebica*. Arabian Journal of Chemistry (2018), <https://doi.org/10.1016/j.arabjc.2017.12.033>

the *Diospyros celebica* Bakh were analyzed by GC–MS and FT-IR. To determine the active molecules of *Diospyros celebica* Bakh, figurative effect of human care function.

## 2. Materials and methods

### 2.1. Materials

*Diospyros celebica* used in the experiment was produced in Indonesia. When we do the experiment, the *Diospyros celebica* Bakh are first pulverized and then tested with the obtained wood powder. The ethanol, benzene and methanol used in the experiments were purely chromatographed. Quantitative filter paper should be extracted with ethanol for 12 h. The three extracts used in the experiment were ethanol, ethanol/benzene (volume ratio of 1:2) and ethanol/methanol (volume ratio of 1:1).

### 2.2. Experimental methods

#### 2.2.1. Extraction method

The crushed and processed *Diospyros celebica* Bakh's powder was weighed 3 parts and the mass was 10 g (accuracy was 1.0 mg). A well-weighed powder and 250 mL of ethanol, ethanol/benzene (1:2 by volume) and ethanol/methanol (1:1 by volume) were added in the three round bottom flasks respectively. And then refluxed at 85 °C, 82 °C and 80 °C for 4.5 h. The obtained extract was subjected to suction filtration on a circulating water type vacuum pump (YUHUA SHZ-D (III)) using a quantitative filter paper subjected to ethanol extraction treatment for 12 h. Finally, the obtained extract was steamed and concentrated by a rotary evaporator (YUHUA RE-2000A).

#### 2.2.2. Ft-ir method

*Diospyros celebica*'s powder and the concentrated extract refluxed by three kinds of extractants were subjected to FT-IR detection (ThermoFisher Nicolet, 670FT-IR). The scanning of each powder was collected at a spectral resolution of 4 cm<sup>-1</sup> and the spectral range was 400–4000 cm<sup>-1</sup> (Botha and Strydom, 2003; Basheer et al., 2017; Razali and Said, 2017).

#### 2.2.3. TG method

The powder of *Diospyros celebica* was analyzed by thermogravimetric analyzer (TGA Q50 V20.8 Build 34). The carrier gas used in the experiment was high purity nitrogen and the nitrogen release rate was 60 mL/min. The temperature program of TG starts at 30 °C and rises to 250 °C at a rate of 5 °C/min. During the test, the sample's weight (%), Deriv. Weight (%/°C) were recorded (Bassilakis et al., 2001; Ismail and Hanafiah, 2017; Halim et al., 2017).

#### 2.2.4. GC–Ms method

The three extracts were analyzed using a gas chromatography-mass spectrometer (Agilent GC–MS 7890B 5977A). Column HP-5MS (30 m × 250 µm × 0.25 µm). Elastic quartz capillary column, the carrier gas used for high purity helium, flow rate of 1 mL/min. The split ratio is 20:1. The temperature program of the GC starts at 50 °C, rises to 250 °C at a rate of 8 °C/min, and then rises to 300 °C at a rate of 5 °C/min. MS program scan mass range of 30–600 amu, ionization voltage of 70 eV, ionization current of 150 µA electron ionization (EI). The

ion source and the quadrupole temperature were set at 230 °C and 150 °C, respectively.

#### 2.2.5. TDS–GC–MS method

The *Diospyros celebica*' powder was analyzed with thermal desorption–gas chromatography–mass spectrometry. TDS starting temperature of 30 °C, for 1 min, at 10 °C/min rate rose to 100 °C, keep 5 min, then 10 °C/min rate rose to 200 °C, the transmission line temperature of 230 °C. CIS starting temperature of –50 °C, hold 0.1 min, and then 10 °C/s rate rose to 230 °C, keep 1 min. Gas Chromatography–Mass Spectrometer (Agilent GC–MS 7890B 5977A). The temperature program of the GC starts at 50 °C, rises to 250 °C at a rate of 8 °C/min, and then rises to 300 °C at a rate of 5 °C/min. MS program scan mass range of 30–600 amu, ionization voltage of 70 eV, ionization current of 150 µA electron ionization (EI). The ion source and the quadrupole temperature were set at 230 °C and 150 °C, respectively. The analytical standard library was analyzed by NIST14.L.


#### 2.2.6. PY–GC–MS method

The powder of *Diospyros celebica* was analyzed by thermal cracking–gas chromatography–mass spectrometry (CDS5200-trace1310 ISQ). The carrier gas used for high purity helium, the pyrolysis temperature was 500 °C, the heating rate was 20 °C/ms, and the pyrolysis time was 15 s. The pyrolysis product transfer line and the injection valve temperature are set to 300 °C; Column TR-5MS; Capillary column (30 m × 0.25 mm × 0.25 µm); Shunt mode, split ratio of 1:60, shunt rate of 50 mL/min. The temperature of the GC program starts at 40 °C for 2 min, rises to 120 °C at a rate of 5 °C/min, and then rises to 200 °C at a rate of 10 °C/min for 15 min. Ion source (EI) temperature of 280 °C, scanning range of 28–500 amu.

## 3. Results and analysis

### 3.1. Ft-ir analysis

Fig. 1 shows the infrared contrast spectra of the *Diospyros celebica* Bakh powder and the three extracts. The infrared



**Fig. 1** FT-IR comparison spectra of *Diospyros celebica* Bakh powders and three extracts.

Download English Version:

<https://daneshyari.com/en/article/10154577>

Download Persian Version:

<https://daneshyari.com/article/10154577>

[Daneshyari.com](https://daneshyari.com)