

Available online at www.sciencedirect.com

ScienceDirect

journal homepage: www.elsevier.com/locate/pepo

Original research article/Artykuł oryginalny

Thalassemia major may decrease the frequency of febrile convulsions in children



Ali Akbar Momen¹, Reza Azizi Malamiri^{1,*}, Bijan Keykhaei Dehdezi²,
Mohsen Fathi³

¹Department of Paediatric Neurology, Golestan Medical, Educational, and Research Center, Ahvaz Jundishapur University of Medical Sciences, Ahvaz, Iran

²Department of Paediatric Hematology and Oncology, Shafa Medical, Educational, and Research Center, Ahvaz Jundishapur University of Medical Sciences, Ahvaz, Iran

³Department of Paediatrics, Golestan Medical, Educational, and Research Center, Ahvaz Jundishapur University of Medical Sciences, Ahvaz, Iran

ARTICLE INFO

Article history:

Received: 02.09.2013

Accepted: 21.10.2013

Available online: 28.10.2013

Keywords:

- Febrile convulsion
- Thalassemia
- Children

ABSTRACT

Aim: We aimed to determine the relative frequency of febrile convulsion in children with major thalassemia to theorize that higher serum iron levels could reduce the incidence of febrile convulsion. **Background:** Febrile convulsion is the most common type of seizure in childhood that its causes are not fully understood. However, some risk factors have been cited such as the serum iron level. **Materials and methods:** Three hundred and fifty-nine children aged more than 5 years with major thalassemia who were receiving blood were enrolled as the case group. The control group consisted of 357 children without thalassemia aged 4–7 years (151 boys, 206 girls) who were referred to healthcare centers for routine health monitoring. Included data were the history of febrile convulsion, age of onset and type and the frequency of convulsions. **Results:** Children in control group significantly experienced more febrile convulsions than thalassemic children [4/359 (1.1%) in the thalassemic children and 14/357 (3.9%) in the control group had experienced febrile convulsions ($P = 0.017$)]. **Conclusion:** The frequency of febrile convulsion in children with major thalassemia is less than that of normal children. Children with thalassemia major may have higher serum levels of iron and such high serum iron levels might have a protective role in the children who have a vulnerability for febrile convulsions.

© 2013 Polish Pediatric Society. Published by Elsevier Urban & Partner Sp. z o.o. All rights reserved.

Introduction

Febrile convulsion is the most common type of seizure during childhood and has a prevalence of 2–4% in different

societies. Febrile convulsion usually occurs in 9 month to 10-year-old children, reaching its peak incidence at 14–18 months of age [1–3]. Although its mortality and morbidity rates are low, but many parents are concerned about the recurrence of seizures [1–3].

* Corresponding author at: Department of Paediatric Neurology, Golestan Medical, Educational, and Research Center, Ahvaz Jundishapur University of Medical Sciences, Golestan Blvd., P.O. Box 6135733118, Ahvaz, Khuzestan, Iran. Tel.: +98 611 3743063; fax: +98 611 3743063; mobile: +98 916 111 6296.

E-mail address: azizi.amin@gmail.com (R.A. Malamiri).

0031-3939/\$ – see front matter © 2013 Polish Pediatric Society. Published by Elsevier Urban & Partner Sp. z o.o. All rights reserved.
<http://dx.doi.org/10.1016/j.pepo.2013.10.002>

The cause and pathophysiology of febrile convulsions are not fully understood. Genetic studies have shown a relation between the genes on chromosomes 8 and 19 and susceptibility to this entity [2]. Some studies have assessed the effect of microelements deficiency, and recently a few studies such as one performed in Iran have focused on iron deficiency, and have recommended the use of iron supplements [4-6]. Two recent studies in Iran and Thailand mentioned a lower frequency of febrile convulsions in patients with major thalassemia [7, 8].

In patients who have thalassemia major, iron is accumulated in the body as a result of ineffective erythropoiesis and frequent blood transfusions. A few studies had reported lower incidence of febrile seizures in children with major thalassemia; therefore, iron accumulation might have a protective or preventive role in the occurrence of febrile convulsions in patients with major thalassemia [7].

In one study in Thailand on 430 patients with thalassemia aged 6 months to 10 years, the researchers found that the frequency of febrile convulsion was 4.4 times lower in children with thalassemia compared with the general population. In the mentioned study, the annual incidence of febrile convulsion was 1.1/1000 individuals in patients with thalassemia, compared with 4.8 in the normal population [7].

In a study performed in Iran comparing patients with febrile convulsion and febrile patients without convulsion, no significant association was found between anemia and the incidence of febrile convulsions [9].

Major thalassemia is more frequent in southwest of Iran (Khouzestan) than that of the other provinces because of the more frequency of thalassemia genes in the general population and consanguineous marriages in this area of Iran. To the best of our knowledge, no study has been performed to measure the frequency of febrile convulsions in patients with major thalassemia in Khouzestan, southwest Iran. We aimed to assess the relative frequency of febrile convulsion in children with major thalassemia comparing to a group of healthy children to theorize that higher serum levels of Iron in children with major thalassemia might have a protective role against febrile convulsions in these children.

Patients and methods

This cross sectional study was performed in Jundishapur University of Medical Sciences, Ahvaz, southwest Iran from April 2010 to April 2011.

Data were collected using related questionnaires. We enrolled 363 patients with major thalassemia over the age of 5 who referred to the thalassemia clinics of Shafa, Abuzar, and Naft Hospitals in Ahvaz as the case group. The patients were confirmed as having major thalassemia based on their history, medical records, blood tests, and hemoglobin electrophoresis. In the control group, 363 healthy children with an age range of 4-7 years who had referred to healthcare centers in Ahvaz for growth monitoring and vaccination were also selected after informing their parents about the aims of the study.

A trained nurse interviewed the participants regarding their demographic information, history of febrile convulsion,

age of the initial onset of convulsion, number of recurrent febrile convulsions, history of febrile convulsions in first degree relatives, familial history of epilepsy, developmental condition, history of related hospital admission, type of febrile convulsion, history of anti-convulsant intake, and history of hypoparathyroidism. A precise history regarding the patients' febrile convulsion and their medical records was taken.

In order to prevent bias, the control group was selected from healthy children who had only referred to the healthcare centers for routine growth assessment.

If the patients were admitted to the hospitals, their medical records were used to diagnose febrile convulsion. If the medical records were unavailable, febrile convulsion was confirmed based on a previous diagnosis made by a pediatrician or family physician.

Febrile convulsion was considered if the seizures were accompanied by fever and no signs of other diseases such as meningitis, encephalitis, and colitis or history of psychomotor retardation, epilepsy, or brain tumor were present.

In the case group, we included patients with beta thalassemia major who had experienced febrile convulsion, neither have psychomotor retardation nor suffer from meningitis, encephalitis, and colitis during seizures, with the informed consent of their parents. Patients whose parents were incapable of remembering the accurate history of convulsions were excluded from our study. The study protocol conforms to the ethical guidelines of the 1975 Declaration of Helsinki as reflected in *a priori* approval by the institution's human research committee.

Statistical analysis was performed using SPSS version 16.0 statistical software (SPSS, Inc., Chicago, IL). χ^2 , *t*, and Fisher's exact tests were used when appropriate. A biostatistician who was blinded to the study groups performed the statistical analysis.

Results

We enrolled to this study, 363 children aged more than 5 years with major thalassemia (169 boys, 194 girls) who were receiving blood as the patient group and 363 children without thalassemia aged 4-7 years (154 boys, 209 girls) who had referred to healthcare centers for routine health monitoring as the control group.

Of the 363 patients with thalassemia major, 4 patients were excluded from the study because of psychomotor retardation (PMR), suspected shivering, suspected breath holding, and history of convulsion at the age of 10 months and long hospital stay. In the control group, 6 children were excluded for reasons such as having meningitis (*n* = 1), shigellosis (*n* = 2), and suspected shivering (*n* = 3).

Among the children with thalassemia major, 4/359 (1.1%) had a history of febrile convulsion as compared with 14/357 (3.9%) children in the control group (*P* = 0.017, χ^2 test). Among the four children in the case group who had a history of febrile convulsion, 3 (1.8%) were girls and 1 (0.5%) was a boy (*P* = 0.25), compared to 9 (5.8%) boys and 5 (2.4%) girls in the control group (*P* = 0.09).

In overall, 18 children had a history of febrile convulsion in both groups including 12 (66.7%) boys and 6 (33.3%) girls.

Download English Version:

<https://daneshyari.com/en/article/10163392>

Download Persian Version:

<https://daneshyari.com/article/10163392>

[Daneshyari.com](https://daneshyari.com)