

## Accepted Manuscript


Title: Eyes should be taped closed during rapid sequence induction of anaesthesia

Author: Hans-Joachim Priebe

PII: S2352-5568(17)30378-8

DOI: <https://doi.org/doi:10.1016/j.accpm.2017.12.010>

Reference: ACCPM 332



To appear in:

Received date: 19-12-2017

Accepted date: 24-12-2017

Please cite this article as: Hans-Joachim Priebe Eyes should be taped closed during rapid sequence induction of anaesthesia (2018), <https://doi.org/10.1016/j.accpm.2017.12.010>

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

**TITLE:****Eyes should be taped closed during rapid sequence induction of anaesthesia**

Hans-Joachim Priebe, MD, FRCA, FACI

Professor Emeritus of Anaesthesia

Department of Anesthesiology and Intensive Care Medicine

University Hospital Freiburg

79106 Freiburg, Germany

Correspondence to: Hans-Joachim Priebe, MD

Email: [hans-joachim.priebe@uniklinik-freiburg.de](mailto:hans-joachim.priebe@uniklinik-freiburg.de)**TITLE:****Eyes should be taped closed during rapid sequence induction of anaesthesia**

Hans-Joachim Priebe, MD, FRCA, FACI

Professor Emeritus of Anaesthesia

Department of Anaesthesiology and Intensive Care Medicine

University Hospital Freiburg

79106 Freiburg, Germany

Correspondence to: Hans-Joachim Priebe, MD

Email: [hans-joachim.priebe@uniklinik-freiburg.de](mailto:hans-joachim.priebe@uniklinik-freiburg.de)

Download English Version:

<https://daneshyari.com/en/article/10211905>

Download Persian Version:

<https://daneshyari.com/article/10211905>

[Daneshyari.com](https://daneshyari.com)