




Contents lists available at ScienceDirect

International Journal of Cardiology

journal homepage: www.elsevier.com/locate/ijcard

Low population prevalence of atrial fibrillation in rural Uganda: A community-based cross-sectional study

Rahul G. Muthalaly^{a,b}, Bruce A. Koplan^{a,b}, Alfred Albano^c, Crystal North^{b,d}, Jeffrey I. Campbell^{b,e}, Bernard Kakuhikire^f, Dagmar Vořechovská^f, John D. Kraemer^g, Alexander C. Tsai^{b,d,f}, Mark J. Siedner^{b,d,f,*}

^a Arrhythmia Service, Cardiovascular Division, Brigham and Women's Hospital, Boston, MA, USA

^b Harvard Medical School, Boston, MA, USA

^c Spectrum Health, Grand Rapids, MI, USA

^d Massachusetts General Hospital, Boston, MA, USA

^e Boston Medical Center, Boston, MA, USA

^f Mbarara University of Science & Technology, Mbarara, Uganda

^g Department of Health Systems Administration, Georgetown University, Washington D.C., USA

ARTICLE INFO

Article history:

Received 17 February 2018

Received in revised form 1 May 2018

Accepted 21 May 2018

Available online xxxxx

Keywords:

Atrial fibrillation

Africa

Uganda

Prevalence

Epidemiology

ABSTRACT

Objectives: Atrial fibrillation (AF) is a major risk factor for stroke, which is the leading cause of cardiovascular mortality in sub-Saharan Africa. However, there is limited population-based epidemiological data on AF in sub-Saharan Africa. We sought to estimate the prevalence and correlates of AF in rural Uganda.

Methods: We conducted a cross-sectional study using community health fairs in 2015 targeting eight villages in rural Uganda. Study participants completed a medical history, a clinical exam, blood collection, and 12 lead electrocardiographic (ECG) screening. Of 1814 participants enrolled in a parent cohort study that includes 98% of adults residing in the geographic area, 856 attended a health fair and were included in this study. Our primary outcome was AF or atrial flutter. We modelled population prevalence of the outcome with inverse probability of treatment weighting using data collected from the full population.

Results: 856 (47.2%) adults in the area attended a health fair and were included in the analysis. Health fair attendees were older (42 vs 34 years, $P < 0.0001$), in worse self-reported health ($P < 0.0001$) and more likely to be female (62% vs 49%, $P < 0.0001$) compared with non-attendees. After applying weights, the estimated population mean age was 37.7 ± 14.9 years. 15% of the population was overweight or obese and 1.9% had left atrial enlargement on ECG. Despite this, the weighted estimate of AF was 0% (95%CI 0–0.54%).

Conclusions: AF appears less prevalent in rural Uganda than in developed countries. The explanations for this finding may be genetic, environmental or related to survivorship bias.

© 2017 Elsevier B.V. All rights reserved.

1. Introduction

Atrial fibrillation (AF) is a major public health concern in developed nations with clinical sequelae including heart failure, cardio-embolic stroke and cardiovascular mortality. Although data from sub-Saharan Africa are sparse, model-based estimates from the Global Burden of Disease Study suggest AF is the fastest growing cardiovascular disease in the region, their data also suggests that factors other than rheumatic heart disease are responsible for the AF burden [1]. The authors

highlight the need for significant additional surveillance and detailed analysis of the mechanisms of atrial fibrillation. If corroborated by population-based data, such estimated trends in AF would be of high public health import, because its risk factors, such as alcohol, hypertension, peripheral vascular disease and tobacco use, and clinical sequelae, such as stroke, are among the highest burden non-communicable public health problems in the region [1,2].

Quality epidemiological data on AF in sub-Saharan Africa are needed to better inform burden of disease models [1]. A meta-analysis of cross-sectional studies including both clinic-based and community-based data from the U.S., Europe, and Commonwealth countries, estimated a prevalence of AF of 2.3% rising to 4.4% among those ≥ 65 years old [3]. Similar attempts to estimate the prevalence of AF in sub-Saharan Africa have been conducted, but these studies were commonly hospital-based and/or have been limited by selection and measurement bias [4,5]. The objective of this study was to use population-

Abbreviations: AF, atrial fibrillation; NCD, non-communicable disease; ECG, electrocardiogram; HbA1c, hemoglobin A1c; HIV, human immunodeficiency virus; LVH, left ventricular hypertrophy; BMI, body mass index; MET, metabolic equivalent of task; IPT, inverse probability of treatment.

* Corresponding author at: Massachusetts General Hospital, 55 Fruit Street, Boston, MA, USA.

E-mail address: msiedner@mgh.harvard.edu (M.J. Siedner).

<https://doi.org/10.1016/j.ijcard.2018.05.074>

0167-5273/© 2017 Elsevier B.V. All rights reserved.

Please cite this article as: R.G. Muthalaly, et al., Low population prevalence of atrial fibrillation in rural Uganda: A community-based cross-sectional study, *Int J Cardiol* (2017), <https://doi.org/10.1016/j.ijcard.2018.05.074>

representative data to estimate the prevalence of electrocardiogram (ECG)-detected AF and its correlates in a rural Ugandan community.

2. Material and methods

2.1. Study design and participants

Residents of Nyakabare Parish were invited to attend free community health fairs by means of radio announcements, printed posters, church announcements and endorsements from community leaders. The health fairs were conducted daily between June 25–29, 2015 in a geographical region approximately 22 km outside of Mbarara Town in southwestern Uganda. The area is rural, the local economy is based primarily on subsistence farming, and both food and water insecurity are common [6]. Prior to the health fairs, during June 2014–May 2015, all adult residents in the parish were enumerated in a population census and invited to participate in a longitudinal cohort study. Approximately 98% (1814 of 1851) of eligible adults consented to participate and completed interviews.

2.2. Data collection

There were two primary sources of data from this study: [1] population census interviews and [2] the community health fairs. During the 2014–2015 population census, we collected demographic, social and geographical variables from virtually all Nyakabare Parish residents. Thus, we had detailed information about the whole population (including demographic, social, psychological and geographical data) eligible to attend the health fairs, whether they attended or not. During health fairs, participants completed a survey administered by trained research assistants to ascertain medical history and sociodemographic characteristics. Physical activity was assessed using the International Physical Activity Questionnaire and converted into metabolic equivalent of task (MET) minutes and divided into three categories: active, minimally active and inactive [7]. Smoking was assessed using a version of the WHO Steps Questionnaire [8].

Participants also underwent short clinical examinations and blood sampling to assess height, weight, blood pressure, creatinine, hemoglobin A1c (HbA1c, Siemens DCA Vantage, Munich, Germany), serum lipids, and human immunodeficiency virus (HIV) rapid antibody testing based on the 2010 Ugandan National HIV Testing Guidelines algorithm. Lastly, the patients underwent a 10-second seated ECG recording using a portable ECG machine (CardioCard Digital ECG Box with CardioCard software, Nasiff Associates, New York, USA).

2.3. ECG interpretation

ECGs were interpreted by a board-certified cardiac electrophysiologist (AA) using a standardised interpretation sheet with 89 possible itemized categories. ECG left atrial enlargement was defined as either a biphasic P-wave in lead V1 with a large negative deflection (>0.1 mV) or prolongation of the P-wave duration to >0.12 s with a widely notched appearance [9]. ECG left ventricular hypertrophy (LVH) was defined using the Sokolow-Lyon criteria of the S wave in V1 or V2 and an R wave in V5 or V6 ≥ 3.5 mV [10]. Atrial fibrillation was defined as the presence of fibrillatory waves associated with an irregular ventricular response. Atrial flutter was defined as the presence of rapid flutter waves [11]. Left bundle branch block was defined as a QRS duration of >0.12 s with a wide slurred R wave in leads I, aVL, V5 and V6, and absent Q waves in leads I, V5 and V6. A Q wave myocardial infarction was defined per the ACC/AHA definition [12].

2.4. Statistical analysis

We described our sample using both standard descriptive techniques of the health fair sample and weighted population estimates. For unweighted characteristics, continuous data are presented as mean \pm SD or median with interquartile range (IQR), depending on their distribution. Normality was assessed visually with the use of histograms. Categorical data are expressed as a percentage with frequency with missing data recorded as missing.

To estimate weighted population characteristics and outcomes, we constructed sampling weights using inverse probability of treatment (IPT) weighting. We first estimated propensity to participate in the health fair using a logistic regression model that included variables which we expected to determine participation: sex, age, marital status, educational attainment, household asset wealth, village of residence, distance from the health fair, difference between the altitude of the household residence and the altitude of the health fair, heavy alcohol use, self-reported HIV status, self-reported overall health, social network size, index of social participation, food insecurity, and water insecurity. The inverse of the predicted conditional probabilities of health fair attendance were applied as IPT weights [13]. The IPT weights enabled population estimates to be generated from the health fair data. We assessed the accuracy of this method using variables that were not included in the IPT model for which we had values for the whole population (as shown in Supplementary Table 2). We also assessed the sensitivity of the model to extreme weights using trimming at the 95th/5th percentiles (as shown in Supplementary Table 4).

We generated population-representative estimates of risk factors and ECG findings applying the sampling weight technique described above. Our primary outcome of interest was the prevalence of AF. Most proportions are presented with logit-transformed confidence intervals and survey design incorporated via Taylor linearization. However, this

approach cannot estimate confidence intervals if the sample proportion is 0%, as it was for some of our outcomes. For such variables, we approximated the confidence interval by calculating Wilson confidence intervals, and then multiplying the upper bound by the design factor from a similar outcome (left bundle branch block). We were unable to conduct our pre-planned regression modelling to identify AF correlates due to zero cells in our primary outcome. All analyses were performed using Stata version 14 (StataCorp LP, College Station, Texas).

2.5. Ethical considerations

All study procedures were approved by the institutional review committees of the Mbarara University of Science and Technology and Partners Healthcare. Consistent with national guidelines, we also obtained clearance for the study from the Ugandan National Council of Science and Technology and from the Research Secretariat in the Office of the Ugandan President. All participants gave written informed consent to participate.

3. Results

Of the 1814 adults included in the Nyakabare parish census, 856 (47.3%) presented to one of the five health fairs. The sample consisted of 320 (37.5%) men, the mean age was 42.3 ± 17.5 years, and the mean BMI was 24.7 ± 5.1 kg/m² as shown in Table 1. There were 105 (12.3%) participants aged <25 years old and 127 (14.8%) aged >65 years old. Forty participants (4.7%) had a positive HIV test result. Four-hundred-and-six (47.4%) had completed at least primary education. Comparing health fair attendees to those in the census who did not attend, women were more likely to attend than men ($P < 0.0001$, Supplementary Table 1). Health fair attendees were older ($P < 0.0001$) and more likely to report bad or very bad overall health ($P < 0.0001$), but were similar in household asset wealth ($P = 0.3$) and village location ($P = 0.1$). Twenty-eight (3.3%) participants had missing ECG data, but were similar in other characteristics as the rest of the sample (Supplementary Table 3).

3.1. Weighted population estimates of atrial fibrillation risk factors

Using IPT weights, the entire population was defined by a mean age of 37.7 ± 14.9 years, BMI 24.4 ± 4.5 kg/m², 2.1% with diabetes, 8.7% with hypertension, 3.9% with self-report of a prior myocardial infarct or heart failure and 3.1% with a prior stroke (Table 1). Almost 30% of the population had consumed alcohol within the past year, of which 17% met criteria for binge drinking (≥ 6 drinks in one occasion). Approximately one quarter (26.4%) were former or current smokers. The majority (75.3%) had high levels of physical activity.

3.2. ECG findings

In weighted estimates, approximately two-thirds of the population had a normal ECG (67.7%, 95% CI 62.4 to 72.6, Table 2). ECG-defined left atrial enlargement was found in 1.9% (95% CI 1.2 to 3.2) and left ventricular hypertrophy in 1.1% (95% CI 0.6 to 1.9). Despite this, there were no cases of atrial fibrillation or atrial flutter detected (0%, 95% CI 0% to 0.54%). Trimmed weight estimates at the 95th/5th percentiles were similar to the untrimmed estimates (Supplementary Table 4).

4. Discussion

In this cross-sectional, population-based study in rural Uganda, we found a 0% prevalence of AF or atrial flutter using single-point 12 lead ECG screening. By applying weights derived from whole-population data to account for self-selection into health fair participation, we obtained a population estimate of AF prevalence with an upper bound confidence interval of approximately 0.5%. These findings are unexpected given the prevalence of multiple established AF risk factors in this cohort, including obesity, hypertension, left atrial enlargement and ischemic heart disease/heart failure [14–17]. Our results contrast with a large meta-analysis of 30 studies—all from developed countries—that used similar single-point screening to estimate the

Download English Version:

<https://daneshyari.com/en/article/10213155>

Download Persian Version:

<https://daneshyari.com/article/10213155>

[Daneshyari.com](https://daneshyari.com)