


Accepted Manuscript

Relation of coronary culprit lesion morphology determined by optical coherence tomography and cardiac outcomes to preinfarction angina in patients with acute myocardial infarction

Nobuaki Kobayashi, Noritake Hata, Masafumi Tsurumi, Yusaku Shibata, Hirotake Okazaki, Akihiro Shirakabe, Masamichi Takano, Kuniya Asai, Yoshihiko Seino, Wataru Shimizu



PII: S0167-5273(18)32180-6
DOI: doi:[10.1016/j.ijcard.2018.07.074](https://doi.org/10.1016/j.ijcard.2018.07.074)
Reference: IJCA 26741

To appear in: *International Journal of Cardiology*

Received date: 3 April 2018
Revised date: 1 June 2018
Accepted date: 16 July 2018

Please cite this article as: Nobuaki Kobayashi, Noritake Hata, Masafumi Tsurumi, Yusaku Shibata, Hirotake Okazaki, Akihiro Shirakabe, Masamichi Takano, Kuniya Asai, Yoshihiko Seino, Wataru Shimizu, Relation of coronary culprit lesion morphology determined by optical coherence tomography and cardiac outcomes to preinfarction angina in patients with acute myocardial infarction. *Ijca* (2018), doi:[10.1016/j.ijcard.2018.07.074](https://doi.org/10.1016/j.ijcard.2018.07.074)

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

Relation of Coronary Culprit Lesion Morphology Determined by Optical Coherence Tomography and Cardiac Outcomes to Preinfarction Angina in Patients with Acute Myocardial Infarction

Brief title: Lesion Morphology, Outcome and Preinfarction Angina

Nobuaki Kobayashi^{a*1}, Noritake, Hata^a, Masafumi Tsurumi^b, Yusaku Shibata^a, Hirotake Okazaki^a, Akihiro Shirakabe^a, Masamichi Takano^c, Kuniya Asai^a, Yoshihiko Seino^c, Wataru Shimizu^d

^aDivision of Intensive Care Unit, Nippon Medical School Chiba Hokusoh Hospital, Chiba, Japan;

^bDepartment of Cardiovascular Medicine, Ishikawajima Memorial Hospital, Tokyo, Japan;

^cDepartment of Cardiology, Nippon Medical School Chiba Hokusoh Hospital, Chiba, Japan;

and ^dDepartment of Cardiovascular Medicine, Nippon Medical School, Tokyo, Japan.

Disclosures: The current study was funded by MSD K. K. and Otsuka Pharmaceutical Co., Ltd.. The funding agencies had no role in designing or conducting the trial.

Total word count: (body text) 2,938 words

* **Corresponding author:** Nobuaki Kobayashi, MD, PhD.

Division of Intensive Care Unit, Nippon Medical School Chiba Hokusoh Hospital

1715, Kamagari, Inzai, Chiba, 270-1694, Japan

E-mail: s5047@nms.ac.jp

1 This author takes responsibility for all aspects of the reliability and freedom from bias of the data presented and their discussed interpretation.

Download English Version:

<https://daneshyari.com/en/article/10213577>

Download Persian Version:

<https://daneshyari.com/article/10213577>

[Daneshyari.com](https://daneshyari.com)