Accepted Manuscript

Increased urinary liver-type fatty acid-binding protein level predicts worsening renal function in patients with acute heart failure

Yousaku Okubo MD , Akinori Sairaku MD , Nobuyuki Morishima MD , Hiroshi Ogi MD , Takeshi Matsumoto MD , Hiroki Kinoshita MD , Yasuki Kihara MD

PII: \$1071-9164(18)30276-8 DOI: 10.1016/j.cardfail.2018.07.003


Reference: YJCAF 4152



Received date: 4 January 2018 Revised date: 1 May 2018 Accepted date: 8 July 2018

Please cite this article as: Yousaku Okubo MD, Akinori Sairaku MD, Nobuyuki Morishima MD, Hiroshi Ogi MD, Takeshi Matsumoto MD, Hiroki Kinoshita MD, Yasuki Kihara MD, Increased urinary liver-type fatty acid-binding protein level predicts worsening renal function in patients with acute heart failure, *The End-to-end Journal* (2018), doi: 10.1016/j.cardfail.2018.07.003

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.



ACCEPTED MANUSCRIPT

Highlights

- An increased urinary L-FABP level was an independent predictor of WRF.
- A higher urinary L-FABP level tended to predict recurrent acute heart failure.
- A high urinary L-FABP level was not associated with the all-cause mortality.

Download English Version:

https://daneshyari.com/en/article/10214059

Download Persian Version:

https://daneshyari.com/article/10214059

<u>Daneshyari.com</u>