

Accepted Manuscript

Nerve ultrasound findings differentiate Charcot-Marie-Tooth disease (CMT) 1A from other demyelinating CMTs

Giampietro Zanette, Gian Maria Fabrizi, Federica Taioli, Matteo Francesco Lauriola, Andrea Badari, Moreno Ferrarini, Tiziana Cavallaro, Stefano Tamburin


PII: S1388-2457(18)31202-1
DOI: <https://doi.org/10.1016/j.clinph.2018.08.016>
Reference: CLINPH 2008636

To appear in: *Clinical Neurophysiology*

Received Date: 19 May 2018
Revised Date: 4 August 2018
Accepted Date: 21 August 2018

Please cite this article as: Zanette, G., Maria Fabrizi, G., Taioli, F., Francesco Lauriola, M., Badari, A., Ferrarini, M., Cavallaro, T., Tamburin, S., Nerve ultrasound findings differentiate Charcot-Marie-Tooth disease (CMT) 1A from other demyelinating CMTs, *Clinical Neurophysiology* (2018), doi: <https://doi.org/10.1016/j.clinph.2018.08.016>

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.



Nerve ultrasound findings differentiate Charcot-Marie-Tooth disease (CMT) 1A from other demyelinating CMTs

Giampietro Zanette,^a Gian Maria Fabrizi,^{b,c} Federica Taioli,^{b,c} Matteo Francesco Lauriola,^a Andrea Badari,^a Moreno Ferrarini,^{b,c} Tiziana Cavallaro,^c Stefano Tamburin,^{b,c}

^aNeurology Division, Pederzoli Hospital, Peschiera del Garda, Verona, Italy;

^bDepartment of Neurosciences, Biomedicine and Movement Sciences, University of Verona, Verona, Italy;

^cNeurology Division, Department of Neuroscience AOUI Verona, Verona, Italy.

Corresponding author:

Stefano Tamburin, MD, PhD

Department of Neurosciences, Biomedicine and Movement Sciences, University of Verona, Piazzale Scuro 10, I-37134 Verona, Italy

Tel.: +39-045-812-4285; +39-347-523-5580

Fax: +39-045-802-7276

Email: stefano.tamburin@univr.it

ORCID: 0000-0002-1561-2187

Highlights

1. Nerve cross sectional area (CSA) is heterogenous across demyelinating CMTs, despite the overlap in nerve conduction study data.
2. Average nerve CSA is significantly enlarged in CMT1A vs. other demyelinating CMTs.
3. The large CSA spectrum in CMT1A may be related to its clinical variability.

Download English Version:

<https://daneshyari.com/en/article/10215353>

Download Persian Version:

<https://daneshyari.com/article/10215353>

[Daneshyari.com](https://daneshyari.com)