Accepted Manuscript



Distinct modes of functional connectivity induced by movie-watching

Murat Demirtaş, Adrian Ponce-Alvarez, Matthieu Gilson, Patric Hagmann, Dante Mantini, Viviana Betti, Gian Luca Romani, Karl Friston, Maurizio Corbetta, Gustavo Deco

PII: S1053-8119(18)30827-9

DOI: 10.1016/j.neuroimage.2018.09.042

Reference: YNIMG 15278

To appear in: *NeuroImage*

Received Date: 14 June 2018

Revised Date: 1 September 2018

Accepted Date: 16 September 2018

Please cite this article as: Demirtaş, M., Ponce-Alvarez, A., Gilson, M., Hagmann, P., Mantini, D., Betti, V., Romani, G.L., Friston, K., Corbetta, M., Deco, G., Distinct modes of functional connectivity induced by movie-watching, *NeuroImage* (2018), doi: https://doi.org/10.1016/j.neuroimage.2018.09.042.

Distinct Modes of Functional Connectivity induced by Movie-Watching

Murat Demirtaş[1,2], Adrian Ponce-Alvarez[2], Matthieu Gilson[2], Patric Hagmann[4],Dante Mantini[5,6], Viviana Betti[7], Gian Luca Romani[8], Karl Friston[9], Maurizio Corbetta[10,11], Gustavo Deco[2,3,12,13]

[1]N3 Division, Department of Psychiatry, Yale University, 40 Temple Street, New Haven, 06511, Connecticut, United States

[2]Center for Brain and Cognition, Computational Neuroscience Group, Department of Information and Communication Technologies, Universitat Pompeu Fabra, Roc Boronat 138, Barcelona, 08018, Spain

[3]Institució Catalana de la Recerca i Estudis Avançats (ICREA), Universitat Pompeu Fabra, Passeig Lluís Companys 23, Barcelona, 08010, Spain

[4]Department of Radiology, Lausanne University Hospital and University of Lausanne (CHUV-UNIL), Rue du Bugnon 46, 1011 Lausanne, Switzerland.

[5]Research Center for Motor Control and Neuroplasticity, KU Leuven, Tervuursevest 101, 3001 Leuven, Belgium.

[6]Functional Neuroimaging Laboratory, IRCCS San Camillo Hospital Foundation, via Alberoni 70, 30126 Venice Lido, Italy.

[7]Department of Psychology, Sapienza University of Rome, via dei Marsi 78, 00185 Rome; Fondazione Santa Lucia and Istituto Di Ricovero e Cura a Carattere Scientifico, 00142, Rome, Italy.

[8]Institute for Advanced Biomedical Technologies, "G. d'Annunzio" University of Chieti-Pescara, 66100, Chieti, Italy.

[9]Wellcome Trust Centre for Neuroimaging, Institute of Neurology, University College London, 12 Queen Square, London WC1N 3BG, United Kingdom.

[10]Department of Neuroscience and Padova Neuroscience Center (PNC), University of Padova, Italy

[11]Departments of Neurology, Radiology, Anatomy of Neurobiology, School of Medicine, Washington University, St. Louis, St Louis, USA

[12]Department of Neuropsychology, Max Planck Institute for Human Cognitive and Brain Sciences, 04103 Leipzig, Germany

[13]School of Psychological Sciences, Monash University, Melbourne, Clayton VIC 3800, Australia

Corresponding author:

Murat Demirtaş, Yale University, 40 Temple Street, New Haven, 06511, Connecticut, United States. murat.demirtas@yale.edu

Funding

GD is supported by the Spanish Research Project PSI2016-75688-P (AEI/FEDER) and by the European Union's Horizon 2020 Framework Programme for Research and Innovation under the Specific Grant Agreement. No. 785907 (Human Brain Project SGA2). MC is supported by NIH grant 5R01NS095741

Acknowledgements

We thank Alan Anticevic, John Murray, Markus Helmer and Joshua Burt for the insightful discussions and their comments during the preparation of this paper.