Accepted Manuscript

Improvement in insulin resistance after gastric bypass surgery is correlated with a decline in plasma 2-hydroxybutyric acid

Prapimporn Chattranukulchai Shantavasinkul , Michael J Muehlbauer , James R Bain , Olga R Ilkayeva , Damian M Craig , Christopher B Newgard , Laura P Svetkey , Svati H Shah , Alfonso Torquati

PII: S1550-7289(18)30173-4 DOI: 10.1016/j.soard.2018.03.033

Reference: SOARD 3345

To appear in: Surgery for Obesity and Related Diseases

Received date: 6 February 2018
Revised date: 13 March 2018
Accepted date: 30 March 2018



Please cite this article as: Prapimporn Chattranukulchai Shantavasinkul, Michael J Muehlbauer, James R Bain, Olga R Ilkayeva, Damian M Craig, Christopher B Newgard, Laura P Svetkey, Svati H Shah, Alfonso Torquati, Improvement in insulin resistance after gastric bypass surgery is correlated with a decline in plasma 2-hydroxybutyric acid, *Surgery for Obesity and Related Diseases* (2018), doi: 10.1016/j.soard.2018.03.033

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

ACCEPTED MANUSCRIPT

Highlight

This is the first report of a decline in 2-hydroxybutyric acid in response to bariatric surgery. Correlation between insulin resistance and 2-hydroxybutyric acid suggests the utility of the latter as an excellent biomarker for tracking glycemic improvement, and offers further insight into the pathways that control diabetes.

Download English Version:

https://daneshyari.com/en/article/10218204

Download Persian Version:

https://daneshyari.com/article/10218204

<u>Daneshyari.com</u>