

Association Between Video Laryngoscopy and Adverse Tracheal Intubation-Associated Events in the Neonatal Intensive Care Unit

Nicole R. Pouppirt, MD¹, Rula Nassar, MD², Natalie Napolitano, MPH, RRT-NPS³, Ursula Nawab, MD¹, Akira Nishisaki, MD, MSCE⁴, Vinay Nadkarni, MD⁴, Anne Ades, MD, MSEd¹, and Elizabeth E. Foglia, MD, MSCE¹

The effect of video laryngoscopy on adverse events during neonatal tracheal intubation is unknown. In this single site retrospective cohort study, video laryngoscopy was independently associated with decreased risk for adverse events during neonatal intubation. (*J Pediatr* 2018;■■■:■■■-■■■).

Neonatal tracheal intubation is a challenging yet life-saving procedure that is associated with adverse events.¹⁻⁵ Video laryngoscopy, which improves the view of the glottis during intubation, may lower the risk of such events.^{5,6} Studies that investigated the impact of video laryngoscopy on adverse events in the pediatric and adult populations report conflicting results.^{6,7} Evidence regarding the effect of video laryngoscopy on adverse events during neonatal intubation is insufficient.⁸⁻¹¹ Randomized controlled trials demonstrated improved neonatal intubation success rates for trainees using video laryngoscopy compared with conventional laryngoscopy, but these trials only examined a limited number of adverse events.^{12,13} We hypothesized that video laryngoscopy is associated with a decrease in adverse events during neonatal tracheal intubation.

Methods

We conducted a retrospective cohort study of neonatal intubations performed at our institution between July 1, 2013 and June 30, 2016. We retrospectively queried the National Emergency Airway Registry for Neonates (NEAR4NEOS), a prospectively developed database, for all intubation encounters in our neonatal intensive care unit (NICU). NEAR4NEOS is a multicenter neonatal airway registry that developed from the pediatric airway registry, National Emergency Airway Registry for Children.¹⁴ Data on patient, provider and practice characteristics, and proximal outcomes are recorded at the time of intubation by the clinical team. The NEAR4NEOS registry is deemed an ongoing Quality Improvement Initiative by the Children's Hospital of Philadelphia Institutional Review Board, and thus, this study of NEAR4NEOS data was not subject to Institutional Review Board approval.

The Children's Hospital of Philadelphia is an urban, academic, training hospital in the US. The Children's Hospital of Philadelphia NICU is a level IV, 98-bed referral center with a small percentage of inborn patients with prenatally diagnosed congenital anomalies. Only intubations occurring in the

NICU setting were included; intubations occurring in the specialized delivery center or other hospital units were excluded. Endotracheal tube exchanges (ie, upsizing or replacing an existing endotracheal tube in an intubated patient) were excluded from this analysis. Patients remain intubated throughout most of the tube exchange procedure, and, thus, the process is distinct from a traditional intubation in a nonintubated patient. Intubations performed by neonatologists, neonatology fellows, pediatric residents, and other NICU staff (nurse practitioners, physician's assistants, and hospitalists) were included. Our unit has a general guideline that limits the number of intubation attempts per provider (up to 2 attempts) and encourages the use of premedication for intubation. The types and dosages of premedication are at the discretion of the neonatologist. An attending neonatologist supervises the majority of NICU intubations.

The exposure of interest was the first laryngoscopy device used for the intubation encounter: video laryngoscopy vs conventional laryngoscopy. The most commonly used video laryngoscope in our unit was the C-MAC (Karl-Storz, Tuttlingen, Germany), introduced into our unit in July 2014. The decision to use the C-MAC for intubation was based on the provider's discretion. Intubations using devices other than the C-MAC video laryngoscope or conventional laryngoscope (ie, other types of video laryngoscope or fiberoptic bronchoscope) were excluded, as these intubations were rare and typically occurred in patients with difficult airways.

We used NEAR4NEOS operational definitions.^{1,6} Briefly, an "encounter" is a single intubation procedure beginning with delivery of premedication and ending 20 minutes after

NEAR4NEOS	National Emergency Airway Registry for Neonates
NICU	Neonatal intensive care unit
TIAE	Tracheal intubation associated event

From the ¹Department of Pediatrics, Division of Neonatology, The Children's Hospital of Philadelphia and The University of Pennsylvania Perelman School of Medicine, Philadelphia, PA; ²Division of Neonatology, Christiana Care Health System, Wilmington, DE; ³Respiratory Therapy Department, The Children's Hospital of Philadelphia; and ⁴Department of Anesthesiology and Critical Care Medicine, The Children's Hospital of Philadelphia and The University of Pennsylvania Perelman School of Medicine, Philadelphia, PA

A.N. and N.N. are supported by the National Institutes of Child Health and Human Development (NICHD) (1R21HD089151, AHRQ R18HS022464, R18HS024511). E.F. is supported by a NICHD Career Development Award (K23HD084727). The authors declare no conflicts of interest.

Portions of this study were presented at the Pediatric Academic Societies annual meeting, May 6-9, 2017, San Francisco, California, and at the Eastern Society for Pediatric Research annual meeting, March 24-26, 2017, Philadelphia, Pennsylvania.

0022-3476/\$ - see front matter. © 2018 Elsevier Inc. All rights reserved.
<https://doi.org/10.1016/j.jpeds.2018.05.046>

securement of endotracheal tube. A “course” is defined as one method of intubation, including the initial device, approach, and medications used. An “attempt” begins with the insertion of a laryngoscope into the patient’s mouth and ends when the laryngoscope is withdrawn. There can be many attempts during a course and multiple courses within an encounter. Only the first course of each intubation encounter was included in this analysis as the study question related to the initial approach to intubation. If a patient underwent multiple intubation encounters in the NICU throughout the study period, the first course of each of these encounters were included in the analysis.

Adverse tracheal intubation associated events (TIAEs) are categorized by NEAR4NEOS as nonsevere or severe. Examples of nonsevere TIAEs include mainstem bronchial intubation, esophageal intubation with immediate recognition, dysrhythmia (including heart rate <60 beats per minute), lip trauma, and pain or agitation requiring additional medication with delay in intubation. Examples of severe TIAEs include cardiac arrest with or without return of spontaneous circulation, esophageal intubation with delayed recognition, pneumothorax, pneumomediastinum, and laryngospasm. Severe oxygen desaturations are defined as $\geq 20\%$ decrease in SpO₂ (oxygen saturation) from the highest value prior to the procedure to the lowest value recorded at any point during the intubation. The highest SpO₂ value is obtained just prior to the intubation, during bag mask ventilation and premedication administration (if administered). Severe oxygen desaturation events are collected separately from TIAEs.

The primary outcome of this study was any adverse TIAE occurring during the first intubation course. Secondary outcomes included severe TIAEs, severe oxygen desaturation events, first attempt success rate, number of attempts during the intubation course, and overall success rates in the first course.

Statistical Analyses

Statistical analysis was performed using STATA 14.0 (Stata Corp, College Station, Texas). Baseline characteristics between groups who were intubated with video laryngoscopy and conventional laryngoscopy were analyzed using χ^2 , Fisher exact, and Wilcoxon rank-sum tests. Associations between the use of video laryngoscopy and adverse TIAEs, severe oxygen desaturation events, number of intubation attempts, and success rates were investigated using χ^2 , Fisher exact, and Wilcoxon rank-sum tests. Logistic regression was performed to determine the independent effect of video laryngoscopy on the outcomes of all adverse TIAEs, severe TIAEs, and severe oxygen desaturation events. In post-hoc analysis, the regression models were adjusted for covariates that significantly differed ($P < .05$) between the video laryngoscopy and conventional laryngoscopy groups in univariable analysis.

Results

Of 805 tracheal intubation encounters performed during the study period, 644 (80%) were performed with conventional

Table I. Comparison of patient, practice, provider characteristics, and outcomes

	Video laryngoscopy (n = 161)	Conventional laryngoscopy (n = 644)	P value
Patient characteristics			
Patient age, d (median, IQR)	40 (10, 82)	15 (1, 30)	<.001
Birth gestational age, wk (median, IQR) (n = 513)	33 (27, 38)	33 (27, 37)	.728
Current weight, kg (median, IQR)	3.0 (1.9, 3.7)	2.6 (1.6, 3.3)	<.001
Male sex	94 (58)	369 (57)	.803
History of a difficult airway	23 (14)	67 (10)	.162
Indication for intubation*			
Respiratory failure	36 (22)	208 (32)	.014
Apnea/bradycardia	18 (11)	95 (15)	.243
Upper airway obstruction	17 (11)	26 (4)	.001
Unplanned extubation	14 (9)	118 (18)	.003
Other indication†	40 (25)	195 (30)	.175
Practice characteristics			
Sedative/analgesic administration	143 (89)	501 (78)	.002
Paralytic administration	129 (80)	284 (44)	<.001
First airway provider			
Pediatric resident	7 (4)	75 (12)	.023
Neonatology fellow	55 (34)	196 (30)	
Neonatology attending	8 (5)	49 (8)	
Other (hospitalist, physician's assistant, nurse practitioner)	91 (57)	324 (50)	
Outcomes			
Any TIAE	10 (6)	124 (19)	<.001
Severe TIAE	3 (2)	30 (5)	.110
Severe oxygen desaturation	70/151 (46)	308/601 (51)	.283
First course number of intubation attempts (median, IQR)	1 (1, 2)	2 (1, 3)	<.001
First intubation attempt success	101 (63)	284 (44)	<.001
First course success	153 (95)	625 (97)	.203

Unless otherwise indicated, values represent patient n (%).

Numbers in bold represent statistically significant values, $P < .05$.

*Each intubation may have more than 1 indication.

†Includes shock, procedural indication, hyperventilation, neurologic weakness, drug administration (including surfactant), and no airway protective reflex.

laryngoscopy and 161 (20%) were performed with video laryngoscopy (Table I). Compared with patients who underwent intubation with conventional laryngoscopy, patients intubated with video laryngoscopy were older (median 40 days, IQR [10-82] vs median 15 days, IQR [1-30], $P < .001$) and larger (median 3.0 kg, IQR [1.9-3.7] vs median 2.6 kg, IQR [1.6-3.3], $P < .001$) at the time of intubation. Upper airway obstruction was a more common indication for intubation using video laryngoscopy. Compared with the conventional laryngoscopy group, the video laryngoscopy group more commonly received sedative/analgesic (89% vs 78%, $P = .002$) and paralytic premedication (80% vs 44%, $P < .001$).

Adverse TIAEs occurred in 134 intubations (17%) during the study period. The most common TIAEs were esophageal intubation with immediate recognition (10% of all encounters) and dysrhythmia including heart rate <60 beats per minute (3% of all encounters) (Table II; available at www.jpeds.com). Adverse TIAEs occurred less frequently in intubations performed with video laryngoscopy (6%) than conventional laryngoscopy (19%), $P < .001$. There was no significant difference in the rate of severe TIAEs or severe oxygen desaturations between groups. Patients in the video laryngoscopy group underwent fewer intubation attempts (median 1, IQR [1-2] vs

Download English Version:

<https://daneshyari.com/en/article/10222211>

Download Persian Version:

<https://daneshyari.com/article/10222211>

[Daneshyari.com](https://daneshyari.com)