



Original Article

Effects of acute gamma irradiation on *in vitro* culture of *Exacum affine* Balf.f. ex RegelMayuree Limtiyayotin,^a Chutiporn Tosri,^b Natnichaphu Sukin,^a Peeranuch Jompuk^{a, b, *}^a Nuclear Technology Research Center, Faculty of Science, Kasetsart University, Bangkok, 10900, Thailand^b Department of Applied Radiation and Isotopes, Faculty of Science, Kasetsart University, Bangkok, 10900, Thailand

ARTICLE INFO

Article history:

Received 17 July 2017

Accepted 28 June 2018

Available online xxx

Keywords:

Exacum affine Balf.f. ex Regel

Gamma irradiation

in vitro culture

ABSTRACT

Exacum affine Balf. f. ex Regel (Persian Violet) is an ornamental plant with small, purple flowers having yellow pollen in their center and it is mostly cultivated as a houseplant. The effect of acute gamma irradiation on *in vitro* cultures of *E. affine* was studied by treatment with gamma radiation using a Mark I Gamma Irradiator at the Nuclear Technology Research Center, Kasetsart University, Thailand with a Cesium-137 source at doses of 0, 10, 20, 30, 40, 50 and 60 Gy and a dose rate of 3.74 Gy/min. After irradiation, subcultures were grown on new Murashige and Skoog (MS) medium; then, after 60 d, the numbers of new branches on surviving plantlets were counted. Samples were subsequently subcultured to new MS medium for morphological study. The median lethal dose was 45 Gy and the 50% decreasing growth rate was 32 Gy. Some variations were observed in irradiated samples in M_1V_2 , such as changes in the number of petals in a flower (four and six petals), flower color (light purple, white), smaller leaves, curled-shaped leaves, variegated leaves, lighter-green leaves and shorter internodes than the control.

Copyright © 2018, Kasetsart University. Production and hosting by Elsevier B.V. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

Introduction

Exacum affine is one of the rare blue-flowering potted plants. In addition to its attractive blue and purple flowers, its popularity is attributable to its fragrance, shiny green foliage, low-mounding habit, ease of shipping and post-harvest shelf life (Riseman, 2007). *Exacum affine* is a species of plant in the family Gentiana-ceae, native to Socotra, Yemen (Miller and Morris, 2004). It is a biennial plant which grows 20–30 cm tall with green ovate leaves and purple stellate flowers with yellow pollen on the stamens in their center. It is commonly known by its commercial name of Persian violet.

Due to the relatively narrow germplasm base and low level of variation found in *E. affine*, mutation breeding has been a viable method to generate genetic variation (Riseman, 2007). Many researchers have shown interest in induced mutation in *E. affine*. Masae et al. (2014) exposed *E. affine* plants to ultra-violet C radiation (UV-C) at 0–32.40 kJ/m² and reported that the median lethal dose (LD₅₀) of shoot apex and axillary explants was 29.7 and 18.9 kJ/m², respectively, and the morphological characteristics of irradiated

samples were a larger size and darker green color compared to the control. A suggested dose for gamma-ray-induced mutation in tissue cultures of *Exacum* sp. is 4 Gy (Shu et al., 2011)

The current research studied the effect of acute gamma irradiation on tissue culture of *E. affine*. Gamma-ray-induced mutations in tissue culture may be another good technique for increasing the genetic variation in *E. affine*.

Materials and methods

Propagation of Exacum affine (Persian Violet) in tissue culture

Pathogen-free plantlets of *E. affine* were cultured on MS medium (Murashige and Skoog, 1962) until enough plantlets were produced for the experiments. Plantlets were multiplied from single-node cuttings transferred to MS medium 90 d before being subjected to radiation treatments.

Acute gamma irradiation

E. affine plantlets were exposed to acute gamma radiation using a Mark I Gamma Irradiator with a ¹³⁷Cs source at the Nuclear Technology Research Center, Kasetsart University, Bangkok, Thailand at doses of 0, 10, 20, 30, 40, 50 and 60 Gy (at a dose rate of

* Corresponding author. Nuclear Technology Research Center, Faculty of Science, Kasetsart University, Bangkok, 10900, Thailand.

E-mail address: fsciprk@ku.ac.th (P. Jompuk).

<https://doi.org/10.1016/j.anres.2018.06.004>

2452-316X/Copyright © 2018, Kasetsart University. Production and hosting by Elsevier B.V. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

Table 1
Survival percentage and growth rate percentage of *E. affine* at 60 d after irradiation.

Radiation dose (Gy)	Plantlet survival (as % of control)	Growth rate (as % of control)
0	100.00 ^{a1}	100.00 ^a
10	84.76 ^{ab}	68.32 ^b
20	93.38 ^{ab}	68.85 ^b
30	70.37 ^{bc}	53.36 ^{bc}
40	56.08 ^{cd}	40.45 ^c
50	44.45 ^{de}	37.08 ^{cd}
60	28.84 ^e	21.15 ^d
F-test	a	a
Coefficient of Variation (%)	20.33	19.21

^a Significant at 1% level.

¹ means within columns followed by a common superscript letters are not significantly different ($p < 0.05$).




Fig. 1. Survival percentage and growth rate percentage at 60 d after irradiation.

3.74 Gy/min). A completely randomized design was used with three replications for each treatment (dose), with 20 plantlets per replication. Following irradiation, the plantlets were then multiplied from two-node cuttings by subculturing on the same medium. After 60 d, for M_1V_1 generation the number of surviving plantlets and the number of new branches were recorded to calculate the median lethal dose ($LD_{50(60)}$) (50% lethal dose at 60 d after irradiation) and the median decreasing growth rate ($GR_{50(60)}$). Desirable variations were recorded and selected in the M_1V_2 generation.

Statistical analysis

The data were analyzed using analysis of variance, after which means were compared using the least significant difference (LSD). The analyses were facilitated by the STAR program (International Rice Research Institute, 2013).

Results and discussion

Effect of acute gamma irradiation on the survival and growth rate of E. affine plantlets

Exposing *E. affine* plantlets raised in aseptic conditions to acute gamma radiation at doses of 0, 10, 20, 30, 40, 50 and 60 Gy showed that 60 d after irradiation, the survival percentages of plantlets (M_1V_1) that had received 10 and 20 Gy of radiation were not different from the control. The percentage of surviving plantlets decreased as the dosage of radiation increased (Table 1). The LD_{50} at 60 d was 45 Gy (Fig. 1) and the effect of irradiation dose on survival percentage was highly significant ($p < 0.01$). The results of the current study differed from Shu et al. (2011) who reported the suggested dose for gamma-ray-induced mutation in tissue cultures of *Exacum* sp. is 4 Gy, but the dose rate in that study was not reported. It is suspected that the difference was due to different dose rates and different ages of plantlets.




Fig. 2. Control and irradiated samples 60 d after irradiation: (A) Control; (B) 10 Gy; (C) 20 Gy; (D) 30 Gy; (E) 40 Gy; (F) 50 Gy; (G) 60 Gy.

Download English Version:

<https://daneshyari.com/en/article/10224952>

Download Persian Version:

<https://daneshyari.com/article/10224952>

[Daneshyari.com](https://daneshyari.com)