

Accepted Manuscript

Diagnosis of PSP-P: Can a newly developed MRPI make the difference?

Florian Krismer, Klaus Seppi

PII: S1353-8020(18)30394-8

DOI: [10.1016/j.parkreldis.2018.09.009](https://doi.org/10.1016/j.parkreldis.2018.09.009)

Reference: PRD 3775

To appear in: *Parkinsonism and Related Disorders*


Received Date: 30 August 2018

Revised Date: 5 September 2018

Accepted Date: 6 September 2018

Please cite this article as: Krismer F, Seppi K, Diagnosis of PSP-P: Can a newly developed MRPI make the difference?, *Parkinsonism and Related Disorders* (2018), doi: <https://doi.org/10.1016/j.parkreldis.2018.09.009>.

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.



Diagnosis of PSP-P: Can a newly developed MRPI make the difference?

Florian KRISMER¹ and Klaus SEPPI.^{1,2}

¹Department of Neurology, Medical University Innsbruck, Innsbruck, Austria

² Neuroimaging Research Core Facility, Medical University Innsbruck, Innsbruck, Austria

Corresponding author:

Klaus Seppi

Department of Neurology

Medical University Innsbruck

Anichstraße 35

A-6020 Innsbruck

Austria

Email: klaus.seppi@tirol-kliniken.at

In early disease stages neurodegenerative parkinsonian disorders overlap considerably in their clinical presentation hampering early clinical differential diagnosis [1]. However, an accurate diagnosis of different parkinsonian disorders is critical not only for the conduct of clinical studies but also for patient counselling since the natural course of these disorders and their treatment strategies vary considerably [2]. Furthermore, interventional treatment trials for degenerative parkinsonian disorders require a correct diagnosis avoiding inclusion of patients with misdiagnoses, especially because there is an increase of disease-modifying trials with disease-specific candidate interventions

Download English Version:

<https://daneshyari.com/en/article/10227110>

Download Persian Version:

<https://daneshyari.com/article/10227110>

[Daneshyari.com](https://daneshyari.com)