



Contents lists available at ScienceDirect

Schizophrenia Research

journal homepage: www.elsevier.com/locate/schres

The nature and timing of social deficits in child and adolescent offspring of parents with schizophrenia: Preliminary evidence for precursors of negative symptoms?

Leslie E. Horton ^{a,*}, Ashley A. Smith ^a, Gretchen L. Haas ^{a,b}

^a Department of Psychiatry, Western Psychiatric Institute and Clinic, University of Pittsburgh Medical Center, Pittsburgh, PA, USA

^b VA Pittsburgh Healthcare System, Pittsburgh, PA, USA

ARTICLE INFO

Article history:

Received 7 May 2014

Received in revised form 27 June 2014

Accepted 3 July 2014

Available online xxxx

Keywords:

High-risk

Schizophrenia

Social skills

Premorbid adjustment

Social anhedonia

Social amotivation

Negative symptoms

ABSTRACT

Children with social dysfunction and a first-degree relative with schizophrenia are at elevated risk for schizophrenia; however, the nature of this dysfunction is unclear. It was hypothesized that familial high-risk (HR) children and adolescents ($n = 17$) would have social skill deficits relative to healthy controls (HC; $n = 35$). HR participants had a bimodal distribution of social skill scores (47% excellent; 53% poor). HR participants had worse social skills, assertion and empathy scores, suggesting possible developmental precursors to the social amotivation domain of negative symptoms. Characterizing HR children's social deficits could assist identification of those at risk for schizophrenia.

© 2014 Elsevier B.V. All rights reserved.

1. Introduction

Children and adolescents who have a first-degree relative with schizophrenia and manifest poor social functioning have 7–8 times greater odds of developing schizophrenia than high-risk and typical children and adolescents with intact social functioning (Tarbox and Pogue-Geile, 2008). A limitation of previous research is that the nature of social dysfunction in children at familial high risk for schizophrenia (HR) has not been well-characterized in terms of types of deficits and developmental timing, and thus is insufficiently specific as an indicator of liability for schizophrenia.

HR adolescents have impairments in academic and social adjustment (Dworkin et al., 1994; Hans et al., 2000; Niemi et al., 2003), peer and romantic relationships (Glatt et al., 2006), social cognition (Eack et al., 2010), and performance-based measures of social behavior (Schiffman et al., 2004; Gibson et al., 2010). However, less is known about specific domains and timing of deficits in younger HR children.

It is well-established in the literature that people with schizophrenia exhibit a variety of social deficits, and that these likely predate onset of illness by several years, possibly as early as childhood (Bearden et al., 2000; Tarbox and Pogue-Geile, 2008; Tsuji et al., 2013). A social deficit of particular relevance to adults with schizophrenia is social amotivation, a core negative symptom of schizophrenia possibly caused by a disruption in reward functioning, and characterized by reduced ability to anticipate and/or experience pleasure. Social amotivation is comprised of asociality and apathy (Foussias and Remington, 2010). However, few studies have attempted to examine whether there are early developmental precursors or similarities to social amotivation in HR children and adolescents. Social amotivation could be trait-like, and a specific aspect of it (i.e., social anhedonia) may be a sensitive and specific indicator of risk for schizophrenia (Kwapil, 1998). Research is needed to clarify whether early developmental precursors can be detected in this population.

We sought to examine the timing and nature of social deficits in familial HR children (i.e. those with a family history of schizophrenia). It was hypothesized that HR children and adolescents would have deficits in social skills and early indicators of social maladjustment, as contrasted with healthy controls (HC); it was further hypothesized that HR participants would experience impairments on a standardized measure of social skills that potentially represent developmental precursors to social amotivation. We selected the Social Skills Rating

* Corresponding author at: University of Pittsburgh Medical Center, 3811 O'Hara St., Pittsburgh, PA 15213, USA. Tel.: +1 412 624 5627; fax: +1 412 624 4496.

E-mail address: hortonle2@upmc.edu (L.E. Horton).

System (SSRS; Gresham and Elliot, 1990), a grade- and sex-normed measure that includes subscales tapping social reciprocity, motivation, and initiative (“empathy” and “assertion” subscales).

2. Methods

2.1. Participants

Participants included 52 (35 HC and 17 HR) individuals ages 5–19 who were successfully age- and sex-matched across conditions (see Table 1). Parents of participants were recruited from community churches and organizations, and advertisements in public areas and mental health services. We obtained written informed consent according to the guidelines of the Institutional Review Board. For both HC and HR participants, structured diagnostic assessments (SCID, supplemented by the K-SADS) were conducted to rule out the presence of current Axis I psychiatric disorders.

2.2. Assessment

Social and academic adjustment was assessed using the Cannon-Spoor Premorbid Adjustment Scale (PAS; Cannon-Spoor et al., 1982). This measure retrospectively assesses levels of social and academic functioning within each of four periods of development: 1) Childhood (5–11 years), 2) Early Adolescence (12–15), 3) Late Adolescence (16–18), and 4) Adulthood (19+). Items are assessed on a 0–6 scale (0 = no dysfunction to 6 = extremely dysfunctional). We assessed IQ using the Peabody Picture Vocabulary Test-III (PPVT-III; Dunn and Dunn, 1997).

Social skills were assessed using the child-report version of the Social Skills Rating System (SSRS; Gresham and Elliot, 1990), a standardized instrument providing grade- and sex-normed scores on 5 domains (Assertion, Empathy, Cooperation, Responsibility and Self-Control). Items are assessed in terms of Frequency from 0 (never) to 2 (very often), and Importance from 0 (not important) to 2 (critical).

We confirmed a putative (self-reported) diagnosis of DSM-IV schizophrenia or schizoaffective disorder in parents of children in the HR group by SCID interview with the parent, and scored lifetime levels (1 to 9) of education and occupational achievement using the Hollingshead 4-Factor Index of Socioeconomic Status (1975).

3. Results

Demographics are presented in Table 1. HR children had greater minority representation and lower parental occupational achievement

Table 1
Demographic characteristics.

| | Healthy controls (HC) n = 35 | High risk (HR) n = 17 |
|--|---------------------------------|--------------------------|
| Age [Mean, range, (SD)] | 12.2, 5–19, (3.9) | 13.9, 9–19, (3.3) |
| % Male | 37% | 41% |
| IQ: PPVT III [Mean (SD)] | 112.3 (14.2) | 104.7 (17.0) |
| Ethnicity* (% Non-White) | 22% | 41% |
| Years of child participants' education | 6.7 (3.9) | 9.4 (3.4) |
| Mother's highest level of education [Mean (SD)] | 5.5 (1.1) | 4.5 (1.7) |
| Father's highest level of education [Mean (SD)] | 5.3 (1.3) | 5.7 (1.2) |
| Mother's highest level of occupation* [Mean (SD)] | 6.3 (1.7) | 4.2 (1.9) |
| Father's highest level of occupation* [Mean (SD)] | 6.5 (2.2) | 4.4 (2.4) |

* Indicates HC and HR groups significantly differed ($p < 0.01$).

as compared to the HC group but did not differ on other key demographics, suggesting successful age- and sex-matching.

As hypothesized, HR subjects had poorer social skills overall ($p < .03$; see Fig. 1). The group effect was sustained in a post-hoc analysis of covariance, controlling for occupational status of parents $F(1,35) = 6.28, p = 0.02$. A striking, albeit unanticipated, finding was that unlike SSRS scores for HC participants, scores for HR participants had a clearly bimodal distribution, with eight participants exhibiting excellent social skills (77th–96th percentile for grade and sex; a ‘High-SS HR’ group) and nine participants exhibiting significantly below normative social skills (2nd–25th percentile; a ‘Low-SS HR’ group) with no individuals scoring in the mid-range (see Fig. 2). High-SS versus Low-SS HR children did not differ in terms of age, sex, education level or parental education and occupation levels. For HR children, poor overall social skills deficits appeared to be driven by specific deficits in “assertion” and “empathy” domains, as compared to other domains ($p < .05$; see Fig. 3).

For all participants combined, there was a correlation between PAS scores and SSRS scores ($r = -0.345, p < .05$). In terms of the timing of social deficits, HR participants also had worse social adjustment on the PAS across developmental stages, including middle childhood [ages 5–8, $t(43) = -2.70, p = .01$]; late childhood [ages 9–11, $t(37) = -2.46, p = .02$]; early adolescence [ages 12–15, $t(24) = -3.29, p = .003$]; and late adolescence [ages 16–19, $t(8) = -2.52, p < .05$]. Notably, these differences were attributable to the subgroup of HR participants with poor social skills, as reflected by results of a repeated measures ANOVA comparing HC, High-SS HR, and Low-SS HR on early and middle childhood PAS scores revealing a trend ($p < .055$) for a group effect and a difference only between the HC and Low-SS HR groups ($p = 0.003$). Visual inspection of scatter plots for early and late adolescent PAS scores (for which sample was by definition restricted to the adolescent subjects), revealed the same pattern for each of these subsequent time periods; i.e. divergent scores for the two HR subgroups, with good PAS social adjustment scores characteristic of the High-SS HR group (and overlapping with scores for the HC group) and poor social adjustment characteristic of the Low-SS HR group.

4. Discussion

Given the finding of high- and low-social skill subgroups among familial HR children, future studies should examine whether other HR offspring samples reveal similarly divergent social skills clusters and, if so,




Fig. 1. Overall SSRS scores: HR and HC participants.

Download English Version:

<https://daneshyari.com/en/article/10307701>

Download Persian Version:

<https://daneshyari.com/article/10307701>

[Daneshyari.com](https://daneshyari.com)