



Short communication

Smooth muscle cell contraction increases the critical buckling pressure of arteries

Danika M. Hayman^{a,1}, Jinzhou Zhang^{b,1,2}, Qin Liu^b, Yangming Xiao^b, Hai-Chao Han^{a,b,*}^a Biomedical Engineering Program, UTSA-UTHSCSA, United States^b Department of Mechanical Engineering, University of Texas at San Antonio, United States

ARTICLE INFO

Article history:

Accepted 21 November 2012

Keywords:

Critical buckling pressure
Mechanical instability
Smooth muscle cell contraction
Vascular tone
Stability

ABSTRACT

Recent *in vitro* experiments demonstrated that arteries under increased internal pressure or decreased axial stretch may buckle into the tortuous pattern that is commonly observed in aging or diseased arteries *in vivo*. It suggests that buckling is a possible mechanism for the development of artery tortuosity. Vascular tone has significant effects on arterial mechanical properties but its effect on artery buckling is unknown. The objective of this study was to determine the effects of smooth muscle cell contraction on the critical buckling pressure of arteries. Porcine common carotid arteries were perfused in an *ex vivo* organ culture system overnight under physiological flow and pressure. The perfusion pressure was adjusted to determine the critical buckling pressure of these arteries at *in vivo* and reduced axial stretch ratios (1.5 and 1.3) at baseline and after smooth muscle contraction and relaxation stimulated by norepinephrine and sodium nitroprusside, respectively. Our results demonstrated that the critical buckling pressure was significantly higher when the smooth muscle was contracted compared with relaxed condition (97.3 mmHg vs 72.9 mmHg at axial stretch ratio of 1.3 and 93.7 mmHg vs 58.6 mmHg at 1.5, $p < 0.05$). These results indicate that arterial smooth muscle cell contraction increased artery stability.

© 2012 Elsevier Ltd. All rights reserved.

1. Introduction

Tortuous arteries are seen in various locations of the vasculature, such as the internal carotid arteries, iliac arteries, coronary arteries, retinal arteries, and conjunctival arteries (Metz et al., 1961; Jakob et al., 1996; Del Corso et al., 1998; Amemiya and Bhutto, 2001; Cheung et al., 2001; Owen et al., 2008; Han, 2012). Arterial tortuosity has been associated with high blood pressure, aging, atherosclerosis, and other pathological changes (Del Corso et al., 1998; Pancera et al., 2000; Han, 2012). Studies have also demonstrated that axial elongation and reduced axial tension could lead to artery tortuosity (Leipzig and Dohrmann, 1986; Jackson et al., 2005), suggesting a possible role of mechanical stress in the development of artery tortuosity. Recent studies, on the other hand, have demonstrated that arteries buckle under increased lumen pressure, weakened wall and decreased axial tension (Han, 2007, 2008, 2011; Liu and Han, 2012). Model analysis and *in vitro* experiments demonstrated that buckled arteries exhibit a tortuous pattern, suggesting buckling as a

possible mechanism for the development of artery tortuosity (Han 2007, 2009, 2012; Lee et al., 2012). Therefore, it is important to understand the behavior of artery buckling.

Computational and experimental studies have demonstrated that the critical buckling pressure, at which the arteries start to deform into wavy shapes, depends on artery dimensions, wall stiffness and axial stretch ratio. Arteries with larger diameters, thicker walls, and higher levels of axial tension all exhibit more stability (Han, 2007; Martinez et al., 2010; Lee et al. 2012). *In vivo*, mechanical properties of the artery wall are also influenced by smooth muscle tone. Vasomotion driven by smooth muscle cell (SMC) contraction and relaxation is one mechanism through which the arteries regulate local blood pressure and flow (Humphrey, 2002). Smooth muscle contraction decreases lumen diameter (vasoconstriction) but increases artery stiffness (Wilkinson and McEniery, 2004; Tremblay et al., 2010). However, it is unclear how SMC tone affects artery stability.

Accordingly, the objective of this study was to determine how SMC tone affects the critical buckling of the artery, by examining the critical buckling pressures of arteries challenged with norepinephrine (NE) and sodium nitroprusside (SNP).

2. Materials and methods

Porcine common carotid arteries were collected from farm pigs post-mortem (6–7 months, 100–150 kg) with the approval by the Texas Department of State

* Corresponding author at: Department of Mechanical Engineering, The University of Texas at San Antonio, San Antonio, TX 78249, United States. Tel.: +1 210 458 4952; fax: +1 210 458 6504.

E-mail address: hchan@utsa.edu (H.-C. Han).

¹ Contributed equally to this work

² On leave from the Department of Cardiovascular Surgery, Xijing Hospital, Fourth Military Medical University, Xi'an, China

Health Service and the University of Texas at San Antonio (UTSA) Institutional Biosafety Committee. Arteries were transported to the lab in ice cold phosphate buffered saline (PBS) supplemented with 1.4% antibiotic/antimycotic (Gibco, Grand Island, NY). Excess loose connective tissue was removed and an inflation test was performed to verify arterial wall integrity. The arteries were tied onto cannulas in a tissue chamber of an *ex vivo* organ culture system (Lee et al., 2010a, 2010b) and perfused with DMEM (Sigma, St. Louis, MO) supplemented with sodium bicarbonate (3.7 g/L), antibiotic/antimycotic (1.4%, Gibco), L-glutamine (1%, Sigma), calf serum (10%, Sigma), HEPES buffer (2.5%, Gibco) and dextran (50–55 g/L, Sigma) to increase viscosity to a physiological level (~4 cP). The arteries were immersed in the same media without dextran and cultured overnight (~12 h, 37 °C, and 5% CO₂) under a flow rate of 160 ml/min and a pressure of 80 ± 2 mmHg to allow the arteries to regain their basal tone. Our previous studies have demonstrated that vasomotor responses were similar under steady or pulsatile pressure after 3 days in organ culture (Hayman et al., 2012).

Two groups of arteries were cultured overnight at a physiological axial stretch ratio of 1.5 ($n=5$, which was at their *in vivo* lengths; Han et al., 2003) and a sub-physiological axial stretch ratio of 1.3 ($n=7$). Buckling tests, as described previously (Martinez et al., 2010; Lee et al., 2012), were performed at the stretch ratios under which the arteries were cultured. Briefly, perfusion pressure in the arteries was gradually reduced to a low level (~15 mmHg) by fully loosening the resistance clamp. The pressure was then gradually increased until buckling occurred and a visually significant deflection (3–8 mm) occurred post-buckling. The buckling test was performed for arteries at baseline, and then repeated after the arteries were contracted by NE (10^{-5} M for ~5 min). It was repeated again after the arteries were relaxed by SNP (10^{-5} M for ~5 min; Hayman, 2011). Our previous studies have demonstrated that arteries were fully contracted or relaxed at these concentrations (Lee et al., 2010a; Hayman et al., 2012). The buckling process was recorded using a Sony DCR-SX63 camcorder. Arterial outer diameter and deflection (displacement of the central axis at the middle of the arterial segments) were measured from images extracted from the video at increments of ~10 mmHg using Image-Pro Plus.

Artery diameters were measured over the range of pressures in the buckling test (70–170 mmHg and 20–120 mmHg for the groups of stretch ratios 1.3 and 1.5, respectively). Both the diameter at a lumen pressure of 80 mmHg and the average over the entire range were used to describe the contraction and relaxation of the arteries. Artery contraction was calculated as the difference between the baseline and the NE-induced contracted diameters divided by the baseline diameter. Artery relaxation was quantified as the difference between the NE-induced contracted diameter and the SNP-induced relaxed diameter divided by the NE-induced contracted diameter. The basal tone was taken as the percent diameter increase from the baseline to the SNP-induced relaxed condition. The critical buckling pressure was taken as the pressure at which the deflection reached 0.5 mm; this point was interpolated from the pressure deflection curve using the TREND function in EXCEL.

The average critical buckling pressure for each group was calculated and statistical significance was determined using a two-way ANOVA followed by a Tukey's post hoc analysis (JMP version 8).

3. Results

Arteries regained their vasomotor tone after being cultured overnight. When pressurized, arterial diameter increased with increasing pressure, at all axial stretch ratios and agonists. Compared to baseline, arterial diameter decreased across all arteries by NE-induced SMC contraction and increased by SNP-induced SMC relaxation (Fig. 1). For the artery group tested under axial stretch ratio 1.3, the contraction and relaxation (mean ± SD, $n=6$) averaged over the entire pressure range (70–170 mmHg) were $6.0 \pm 8.4\%$ and $13.9 \pm 9.0\%$, in response to NE and SNP, respectively ($p < 0.05$), with a basal tone of $10.4 \pm 9.3\%$. At a lumen pressure of 80 mmHg, the contraction and relaxation (mean ± SD, $n=6$) were $4.9 \pm 1.3\%$ and $5.7 \pm 1.2\%$, respectively, in response to NE and SNP, with a basal tone of $5.1 \pm 0.9\%$. Similarly, for the artery group cultured under axial stretch ratio 1.5, the contraction and relaxation (mean ± SD, $n=5$) averaged over the entire pressure range (20–120 mmHg) were $3.7 \pm 1\%$ and $22.7 \pm 7\%$, in response to NE and SNP, respectively ($p < 0.05$), with a basal tone of $18 \pm 9\%$. At a lumen pressure of 80 mmHg, the contraction and relaxation (mean ± SD, $n=5$) were $5.9 \pm 1.1\%$ and $8.1 \pm 1.9\%$ in response to NE and SNP, respectively, with a basal tone of $5.9 \pm 1.7\%$.

During the buckling test, artery deflection was initiated at a critical pressure and gradually increased with increasing lumen




Fig. 1. Comparison of outer diameter change as a function of lumen pressure of representative arteries at baseline, after administration of norepinephrine (NE), and sodium nitroprusside (SNP). The arteries were tested at a stretch ratio of 1.3 (a) and 1.5 (b).

pressure (Fig. 2). NE-induced contraction shifted the pressure–deflection curve to the right while SNP-induced relaxation shifted the curve to the left, indicating that NE-induced SMC contraction increased the critical pressure and SNP-induced SMC relaxation decreased the critical buckling pressure (Fig. 3). The pressure–deflection curves all exhibit a nonlinear accelerated increase immediately at post-buckling but tend to slow down at higher pressures further beyond buckling (Fig. 3a). This trend has been observed previously (Lee, 2011, 2012), so we did not measure the high pressure range for most of these arteries. In comparison, the critical buckling pressures of the arteries at NE-induced contraction were significantly higher than those at SNP-induced relaxation condition, at both axial stretch ratios of 1.3 and 1.5 ($p < 0.05$, Fig. 4). No significant difference was observed between the critical buckling pressures of arteries at the baseline and SNP-induced relaxation condition.

4. Discussion

Our results indicate that contraction of the arteries significantly increases the critical buckling pressure compared with the dilated state. Interestingly, the critical buckling pressure of arteries at baseline showed no significant difference from that at the fully relaxed condition. This suggests that the critical buckling pressure measured *in vitro* in fully relaxed arteries is close to *in vivo* value though it may slightly underestimate the critical buckling pressure *in vivo*.

Download English Version:

<https://daneshyari.com/en/article/10432962>

Download Persian Version:

<https://daneshyari.com/article/10432962>

[Daneshyari.com](https://daneshyari.com)