

## Accepted Manuscript

Optimization of Dendrimer Structure for Sentinel Lymph Node Imaging:  
Effects of Generation and Terminal Group

Yuichiro Niki, Mikako Ogawa, Rie Makiura, Yasuhiro Magata, Chie Kojima


PII: S1549-9634(15)00166-5  
DOI: doi: [10.1016/j.nano.2015.08.002](https://doi.org/10.1016/j.nano.2015.08.002)  
Reference: NANO 1169

To appear in: *Nanomedicine: Nanotechnology, Biology, and Medicine*

Received date: 14 March 2015  
Revised date: 4 August 2015  
Accepted date: 18 August 2015

Please cite this article as: Niki Yuichiro, Ogawa Mikako, Makiura Rie, Magata Yasuhiro, Kojima Chie, Optimization of Dendrimer Structure for Sentinel Lymph Node Imaging: Effects of Generation and Terminal Group, *Nanomedicine: Nanotechnology, Biology, and Medicine* (2015), doi: [10.1016/j.nano.2015.08.002](https://doi.org/10.1016/j.nano.2015.08.002)

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.



# Optimization of Dendrimer Structure for Sentinel Lymph Node Imaging: Effects of Generation and Terminal Group

Yuichiro Niki<sup>a</sup>, Mikako Ogawa<sup>b</sup>, Rie Makiura<sup>c</sup>, Yasuhiro Magata<sup>b</sup>, Chie Kojima<sup>a\*</sup>

<sup>a</sup> Department of Applied Chemistry, Graduate School of Engineering, Osaka Prefecture University, 1-1 Gakuen-cho, Naka-ku, Sakai, Osaka 599-8531, Japan

<sup>b</sup> Medical Photonics Research Center, Hamamatsu University School of Medicine, 1-20-1 Handayama, Higashi-ku, Hamamatsu, Shizuoka 431-3192, Japan

<sup>c</sup> Nanoscience and Nanotechnology Research Center, Research Organization for the 21st Century, Osaka Prefecture University, 1-2 Gakuen-cho, Naka-ku, Osaka 599-8570, Japan

\*Corresponding author: Department of Applied Chemistry, Graduate School of Engineering, Osaka Prefecture University, 1-2 Gakuen-cho, Naka-ku, Sakai, Osaka 599-8570, Japan. Tel.: +81 72 254 8190; Fax: +81 72 254 8190. E-mail: kojima@chem.osakafu-u.ac.jp (C. Kojima).

The authors declare no conflict of interest.

This manuscript was supported by a grant-in-aid from the Japan Society for the Promotion of Science (JSPS) for Scientific Research (C) (No. 26410227).

Word count for abstract: 147

Word count for manuscript: 3488 (Background, Methods, Results, Discussion and Figure legends)

Number of references: 35

Number of figures: 5

Number of tables: 1

Number of Supplementary online-only files: 1

Download English Version:

<https://daneshyari.com/en/article/10436023>

Download Persian Version:

<https://daneshyari.com/article/10436023>

[Daneshyari.com](https://daneshyari.com)