



Short Communication

Associations between self-control and dimensions of nicotine dependence: A preliminary report

Stephen J. Wilson^{*}, R. Ross MacLean

Department of Psychology, The Pennsylvania State University, University Park, PA, 16802, United States

HIGHLIGHTS

- ▶ Self-control has been linked to overall nicotine dependence in those who smoke.
- ▶ Associations between self-control and dimensions of nicotine dependence are unclear.
- ▶ We examined the correlation between self-control and several facets of dependence.
- ▶ Self-control negatively correlated with smoking driven by craving and withdrawal.
- ▶ Self-control positively correlated with the consistency of smoking patterns.

ARTICLE INFO

Keywords:

Craving
Habit
Individual differences
Nicotine dependence
Self-control
Smoking

ABSTRACT

Self-control plays an important role in several health-related behaviors, including cigarette smoking. There is some evidence that individual differences in self-control are negatively associated with overall levels of nicotine dependence but, to our knowledge, finer-grained relationships between these constructs have not been explored. This is an important knowledge gap, as nicotine dependence is thought to be composed of separate dimensions that motivate smoking behavior in relatively unique ways. The goal of this preliminary study was to begin to characterize the potentially nuanced associations between self-control and facets of nicotine dependence using data pooled from two previous studies ($n = 282$). Specifically, we examined the correlation between self-control and the following dimensions of nicotine dependence: compulsion to smoke due to craving and desire to avoid withdrawal symptoms; preference for smoking over other reinforcers; reduced sensitivity to the effects of smoking; consistency of smoking patterns; and smoking behavior that is rigid and immutable. In line with prior research, self-control was negatively correlated with overall levels of dependence. As predicted, however, self-control was differentially associated with distinct dimensions of nicotine dependence. Specifically, self-control was negatively correlated with the compulsion to smoke due to craving and desire to avoid withdrawal symptoms but positively correlated with the consistency of smoking patterns. Given the potential conceptual and clinical importance of such effects, additional research investigating the role(s) that individual differences in self-control play in addiction to cigarettes would be useful.

© 2012 Elsevier Ltd. All rights reserved.

1. Introduction

Self-control, or the capacity to alter one's own responses in the service of goals and standards (Duckworth & Kern, 2011), plays an important role in several health-related behaviors, including the use of cigarettes (de Ridder, Lensvelt-Mulders, Finkenauer, Stok, & Baumeister, 2012). For instance, relative to those with low self-control, children with high self-control are much less likely to smoke cigarettes during adulthood (Moffitt et al., 2011; Welch & Poulton, 2009). Furthermore, research suggests that individual differences in self-control influence the mechanisms underlying cigarette

use among those who smoke. Specifically, recent findings indicate that smokers who are effective at inhibiting prepotent responses and forgoing immediate rewards in favor of larger delayed rewards (and thus who presumably have greater self-control) are less nicotine dependent overall than smokers who have difficulty regulating behavior and resisting temptation (Billieux et al., 2010; Sweitzer, Donny, Dierker, Flory, & Manuck, 2008).

While it appears that individual differences in self-control are negatively associated with global nicotine dependence, to our knowledge, more nuanced relations between these constructs have not been explored. This is an important knowledge gap, as nicotine dependence is thought to be composed of several distinct facets that motivate smoking behavior in unique ways (Piasecki, Piper, & Baker, 2010; Shiffman, Waters, & Hickcox, 2004; West, 2009). One

^{*} Corresponding author. Tel.: +1 814 865 6219; fax: +1 814 863 7002.
E-mail address: sjw42@psu.edu (S.J. Wilson).

instrument designed to assess separate dimensions of nicotine dependence is the Nicotine Dependence Syndrome Scale (NDSS; Shiffman et al., 2004). Specifically, the NDSS measures five core aspects of tobacco addiction: 1) compulsion to smoke due to craving and desire to avoid withdrawal symptoms (*Drive*); 2) preference for smoking over other reinforcers (*Priority*); 3) reduced sensitivity to the effects of nicotine/smoking (*Tolerance*); 4) consistency/regularity of smoking patterns (*Continuity*); and 5) smoking behavior that is rigid and insensitive to contextual factors (*Stereotypy*). Self-control may be more strongly related to some of these dimensions relative to others. In particular, given its importance for overriding strong but undesirable impulses (Muraven & Baumeister, 2000), it seems especially likely that self-control is inversely related to the degree to which smokers experience cravings to smoke.

In addition to regulating impulses, it recently has been proposed that self-control is critical for establishing stable patterns of behavior (de Ridder et al., 2012). This novel perspective emerged from a meta-analysis revealing that self-control is more strongly associated with behaviors that are relatively automatic than with behaviors that are effortful or demanding. The development of highly regular and inflexible smoking patterns is thought to be a core feature of cigarette addiction (Tiffany, 1990). To the extent that self-control facilitates the acquisition of habits, it follows that individual differences in self-control actually may be *positively* related to the automatization of smoking behavior.

The aim of this preliminary study was to examine associations between self-control and facets of nicotine dependence in cigarette smoking adults. Consistent with previous research, we predicted that self-control would correlate negatively with overall dependence. We hypothesized, however, that self-control would relate differentially to conceptually distinct dimensions of dependence. Specifically, we predicted that self-control would correlate negatively with the degree to which smoking is driven by cravings but positively with the consistency and rigidity of smoking patterns.

2. Method

2.1. Participants

Participants ($n = 282$) were drawn from two prior studies. Study 1 (Wilson, Sayette, & Fiez, 2012) examined the effects of motivational status and smoking opportunity on neural responses to a smoking cue and included male and females and smokers who were and who were not motivated to quit smoking. Study 2 (Wilson, Sayette, & Fiez, *in press*) examined the neural mechanisms associated with cognitive strategies for coping with exposure to a cigarette cue and included male smokers who were motivated to quit smoking. For both studies, participants had to report smoking an average of 15–40 cigarettes per day for the past 24 months. Table 1 reports demographic characteristics of the individual and combined samples.

2.2. Materials

2.2.1. NDSS

The NDSS consists of 19 statements related to smoking habits (e.g., “I smoke consistently and regularly throughout the day.”), with

participants rating each item according to how well it describes them using a 6-point scale anchored by 1 (“Not at all true”) and 6 (“Extremely true”) (Shiffman et al., 2004). As described in the introduction, the NDSS is designed to assess five theoretically-derived dimensions of dependence. The NDSS also yields a summary measure of dependence (NDSS-Total; 14 items). The subscales of the NDSS have demonstrated adequate internal consistency ($\alpha = .55-.76$), as has the summary score ($\alpha = .84$) (Shiffman et al., 2004). Scores on the NDSS also have been found to correlate with other measures of dependence, predict cessation-related outcomes, and discriminate dependent from regular, but non-dependent, smokers (Shiffman & Sayette, 2005; Shiffman et al., 2004). The NDSS was scored using the regression-based algorithms described by Shiffman et al. (2004), with higher scores indicating a higher level of dependence ($M = 0$, $SD = 1$ in the normative sample).

2.2.2. Self-Control Scale (SCS)

The SCS consists of 36 self-descriptive items (e.g., “I have a hard time breaking bad habits”) rated on 5-point scale anchored by 1 (“Not at all like me”) and 5 (“Very much like me”) (Tangney, Baumeister, & Boone, 2004). The SCS has demonstrated high internal consistency ($\alpha = .89$) and scores on the measure have been found to correlate with a variety of behaviors thought to require self-control (e.g., regulation of eating behavior; Tangney et al., 2004). A total score indexing self-control was obtained by summing all 36 responses (the possible score ranged from 36 to 180). Higher scores on the SCS indicate higher self-control.

2.3. Procedure

Participants deemed eligible based upon a telephone screening were scheduled for an initial baseline session. They were instructed to smoke ad libitum prior to the session. Upon arrival, participants provided an exhaled carbon monoxide (CO) sample in order to verify smoking status ($CO > 10$ ppm). Participants then completed a battery of questionnaires (including the NDSS and SCS) and behavioral working memory tasks. Data from these additional questionnaires and tasks are not a focus of the present study. For Study 2, participants also were trained to use cognitive coping strategies after completing questionnaires and memory tasks. A subset of participants completed a functional magnetic resonance imaging experiment held on a separate day (reported in Wilson et al., *in press*).

2.4. Statistical analysis

In order to assess the association self-control and overall nicotine dependence, we calculated the Pearson product-moment correlation between total scores on the SCS and NDSS. Additionally, to test the hypothesis that self-control is associated differentially with distinct facets of dependence, we evaluated the correlation between scores on the SCS and scores on each NDSS subscale. The Bonferroni adjustment was used to control for possible inflation of Type I error rate as a result of the number of correlations performed, yielding a corrected alpha of $p < .008$ ($.05/6$).

3. Results

The mean score on the SCS was 114.40 ($SD = 19.82$). Means and standard deviation for scores on the NDSS and correlations between scores on the SCS and NDSS are presented in Table 2. (All correlations that are identified as significant remain so when controlling for the other subscales of the NDSS and for age and smoking rate. For simplicity, only the zero-order correlations are reported.) As predicted, self-control was negatively correlated with overall nicotine dependence (i.e., NDSS total score). As hypothesized, however, self-control was differentially associated with subscales of the NDSS

Table 1
Sample characteristics.

	Full sample ($n = 282$)	Study 1 ($n = 177$)	Study 2 ($n = 105$)
Percent male	71	53	100
Mean age (SD)	31.7 (8.2)	30.2 (7.6)	34.2 (8.6)
Mean years of formal education (SD)	12.5 (2.1)	12.6 (2.2)	12.5 (1.9)
Mean cigarettes/day (SD)	20.3 (5.3)	20.1 (5.1)	20.6 (5.5)

Download English Version:

<https://daneshyari.com/en/article/10443366>

Download Persian Version:

<https://daneshyari.com/article/10443366>

[Daneshyari.com](https://daneshyari.com)