Accepted Manuscript

Early transcutaneous electrical nerve stimulation reduces hyperalgesia and decreases activation of spinal glial cells in mice with neuropathic pain

Hideaki Matsuo, Kenzo Uchida, Hideaki Nakajima, Alexander Rodriguez Guerrero, Shuji Watanabe, Naoto Takeura, Daisuke Sugita, Seiichiro Shimada, Terumasa Nakatsuka, Hisatoshi Baba

PII: S0304-3959(14)00311-X

DOI: http://dx.doi.org/10.1016/j.pain.2014.06.022

Reference: PAIN 9256

To appear in: *PAIN*

Received Date: 29 March 2014 Revised Date: 15 June 2014 Accepted Date: 30 June 2014



Please cite this article as: H. Matsuo, K. Uchida, H. Nakajima, A.R. Guerrero, S. Watanabe, N. Takeura, D. Sugita, S. Shimada, T. Nakatsuka, H. Baba, Early transcutaneous electrical nerve stimulation reduces hyperalgesia and decreases activation of spinal glial cells in mice with neuropathic pain, *PAIN* (2014), doi: http://dx.doi.org/10.1016/j.pain.2014.06.022

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

ACCEPTED MANUSCRIPT

Revised Manuscript PAIN-D-14-12183-R1, Page 1

Early transcutaneous electrical nerve stimulation reduces hyperalgesia and decreases activation of spinal glial cells in mice with neuropathic pain

Hideaki Matsuo ^a, RPT, Kenzo Uchida ^{b,*}, MD, DMSc, Hideaki Nakajima ^b, MD,

DMSc, Alexander Rodriguez Guerrero ^b, MD, DMSc, Shuji Watanabe ^b, MD, Naoto Takeura ^b, MD, Daisuke Sugita ^b, MD, Seiichiro Shimada ^a, RPT, Terumasa Nakatsuka

^c, MD, DMSc and Hisatoshi Baba ^b, MD, DMSc

^a Division of Physical Therapy and Rehabilitation Medicine, University of Fukui

Hospital, ^b Department of Orthopaedics and Rehabilitation Medicine, Faculty of

Medical Sciences, University of Fukui, Matsuoka Shimoaizuki 23, Eiheiji, Fukui

910-1193, Japan

^c Pain Research Center, Kansai University of Health Sciences, Kumatori, Osaka

590-0482, Japan

Number of text pages of the entire manuscript: 55 pages

Number of figures: 12

Number of tables: 0

*Corresponding author. Address: Kenzo Uchida, MD, DMSc, Department of

Orthopaedics and Rehabilitation Medicine, Faculty of Medical Sciences, University of

Fukui, Matsuoka Shimoaizuki 23, Eiheiji, Fukui 910-1193, Japan.

Tel.: +81 776 61 8383; fax: +81 776 61 8125.

E-mail address: kuchida@u-fukui.ac.jp (K. Uchida).

15

Download English Version:

https://daneshyari.com/en/article/10450241

Download Persian Version:

https://daneshyari.com/article/10450241

<u>Daneshyari.com</u>