

Short communication

Voltammetric assay of Guaifenesin in pharmaceutical formulation

I. Tapsoba^b, J.-E. Belgaied^{a,*}, K. Boujlel^b

^a *Institut National des Sciences Appliquées et de Technologies (INSAT), Chimie et Biologie Appliquées,
B.P. 676, 1080 Tunis, Cedex, Tunisia*

^b *Laboratoire de Chimie Analytique et d'Electrochimie, Faculté des Sciences de Tunis, Département de Chimie,
Campus Universitaire, 2029 El Manar, Tunis, Tunisia*

Received 4 May 2004; received in revised form 24 November 2004; accepted 24 November 2004

Available online 11 January 2005

Abstract

The electrochemical oxidation of Guaifenesin in a pharmaceutical formulation containing Guaifenesin has been carried out in Britton–Robinson buffer (BRB) (0.04 mol L^{-1}) on platinum electrode. Guaifenesin exhibits a well-defined irreversible oxidation peak at 0.924 V/ref . The influence of pH on the oxidation of Guaifenesin was studied in BRB (pH range 2–5). A method for the analysis of Guaifenesin in BRB (0.04 mol L^{-1} , pH 2), which allows quantification over the range $20\text{--}60 \mu\text{g mL}^{-1}$, was proposed and successfully applied to the determination of Guaifenesin in syrup with mean recovery and relative standard deviation of 103.3% and 1.32%, respectively.

© 2004 Elsevier B.V. All rights reserved.

Keywords: Guaifenesin; Oxidation; Cyclic voltammetry; Platinum electrode; Irreversible

1. Introduction

Guaifenesin is a drug that reduces the thickness and stickiness of mucus. It is used for short-term relief of dry, non-productive cough and mucus in the breathing passages. Guaifenesin is used to treat symptoms of allergy, colds and upper respiratory infections. Several methods are reported in the literature proposing fast and reliable techniques for the determination of Guaifenesin in various cough-cold formulations [1–7]. Most of the methods reported for the analysis of Guaifenesin in different matrices, alone or with other active substances, rely on the use of chromatographic techniques. Vasudevan et al. developed and validated a method for the analysis of Guaifenesin in the presence of phenylpropanolamine HCl, diphenylpyraline HCl by HPLC on a C_8 column and a mobile phase consisting of acetonitrile–triethylamine (pH adjusted to 3.5 using orthophosphoric acid; 0.5%), (35:65, v/v) at a flow rate of 1.2 mL min^{-1} . Detection was carried out

at 210 nm [8]. Another HPLC method has been developed by Wilcox and Stewart [9] for the simultaneous determination of Guaifenesin pseudoephedrine–dextromethorphan and Guaifenesin–pseudoephedrine in commercially available capsule dosage forms and Guaifenesin–codeine in a commercial cough syrup dosage form. Guaifenesin has been also analysed by spectrophotometric methods [10–12] and Micellar electrokinetic chromatography [13].

Electrochemical methods have proved to be very sensitive for the determination of organic molecules that undergo oxidation or reduction reactions, including drugs and related molecules in pharmaceutical dosage form and biological fluids [14–18]. The advance in experimental electrochemical techniques in the field of analysis of drugs is due to their simplicity, low cost and relatively short analysis time when compared with other techniques. To the best of our knowledge, no electrochemical method was developed for the analysis of Guaifenesin.

The aims of this study are to establish the experimental conditions and to optimize the conditions for the determination of this compound in a pharmaceutical dosage form using cyclic and differential pulse voltammetry (DPV) techniques.

* Corresponding author. Tel.: +216 71704309; fax: +216 71704329.
E-mail address: Jamel.Belgaied@insat.rnu.tn (J.-E. Belgaied).

2. Experimental

2.1. Reagents

Guaifenesin were purchased from Sigma Chemical Co (St. Louis, MO). A stock standard solution of Guaifenesin at $2000 \mu\text{g mL}^{-1}$ was prepared in distilled water. BRB buffer 0.04 M was prepared from phosphoric acid, boric acid and acetic acid of analytical grade in distilled water. Tussipax (an adult syrup produced by Opalia, Tunisia) labeled to contain as active ingredients dextromethorphan bromhydrate and Guaifenesin at 3 and 10 mg mL^{-1} , respectively, and alcohol, sucrose, glycerol, menthol aroma as excipients. Tussipax was purchased from a commercial source.

2.2. Apparatus

Standard three-electrode potentiostatic circuitry was used, employing a PST 050 potentiostat and a high voltage booster HSB 100 (Radiometer Analytical). The counter electrode was a platinum wire ($\Phi = 500 \mu\text{m}$) and the reference electrode was an Ag/AgNO_3 0.1 mol L^{-1} reference electrode. All potentials are given versus the Ag/AgNO_3 0.1 mol L^{-1} . The working electrode was a platinum wire ($\Phi = 250 \mu\text{m}$, $\Phi = 500 \mu\text{m}$ or $\Phi = 1 \text{ mm}$). The system was controlled by Volta Master 4 software (Radiometer Analytical).

2.3. Calibration

Appropriate dilutions of the standard stock solution of Guaifenesin, were performed with water to obtain final concentrations of Guaifenesin 20 – $60 \mu\text{g mL}^{-1}$. The voltammogram of each standard level were recorded. Calibration curves were constructed by plotting peak areas against concentration in $\mu\text{g mL}^{-1}$ and the linear relationships was evaluated by calculation of regression lines by the method of least square.

2.4. Sample solution

Bottles were thoroughly mixed before sampling to ensure homogeneity. An aliquot of the syrup was then transferred to the electrolysis cell and diluted with BRB (0.04 mol L^{-1}). Voltammograms were recorded as described for pure Guaifenesin.

3. Results and discussion

3.1. Voltammetric behaviour

To elucidate the electrode reaction of Guaifenesin, a cyclic voltammogram at platinum electrode was recorded at different pH and at different scan rates. As an example, Fig. 1 shows the cyclic voltammogram of a $1.1 \times 10^{-2} \text{ (mol L}^{-1}\text{)}$ Guaifenesin in BRB (0.04 mol L^{-1} , pH 2) at a scan rate of 100 mV s^{-1} . The voltammogram exhibits one well-defined anodic peak, with no peak on the reverse scan, suggesting the irreversible nature of the electrode reaction.

The effect of potential scan rate, ν , on the peak current and the peak potential of Guaifenesin was evaluated. A linear relationship was observed between $\log I_p$ and $\log \nu$ over the scan range 20 and 200 mV s^{-1} ($r = 0.9978$) and corresponds to equation: $\log I_p (\mu\text{A}) = 0.4586 \log \nu + 1.3536$, (ν in V s^{-1}). The slope of 0.45 is close to the theoretically expected value of 0.5 for a diffusive process [19]. On the other hand, as scan rate was increased, the potential shifted to less positive values as expected for an irreversible reduction process [20]. The dependence of peak potential of a $1.1 \times 10^{-1} \text{ (mol L}^{-1}\text{)}$ Guaifenesin in BRB (0.04 mol L^{-1} , pH 2) on the decimal logarithm of the scan rate was linear, followed the relationship $E_p (\text{V}) = 0.096 \log \nu + 1.023$, (ν in V s^{-1}). The shift of the peak potential to anodic value of about 96 mV per decade of the logarithm of the sweep rate indicates that the electron transfer is the determining step. Under these conditions, the average experimental $n\alpha$ value was 0.31 .




Fig. 1. Cyclic voltammogram of Guaifenesin in BRB (0.04 mol L^{-1} , pH 2) at platinum electrode. Scan rate 100 mV s^{-1} .

Download English Version:

<https://daneshyari.com/en/article/10554357>

Download Persian Version:

<https://daneshyari.com/article/10554357>

[Daneshyari.com](https://daneshyari.com)