



## Evaluation of synthetic naphthalene derivatives as novel chemical chaperones that mimic 4-phenylbutyric acid



Seisuke Mimori<sup>a,\*</sup>, Yukari Koshikawa<sup>a</sup>, Yu Mashima<sup>a</sup>, Katsuyoshi Mitsunaga<sup>b</sup>, Koichi Kawada<sup>c</sup>, Masayuki Kaneko<sup>d</sup>, Yasunobu Okuma<sup>c</sup>, Yasuyuki Nomura<sup>e</sup>, Yasuoki Murakami<sup>a</sup>, Tetsuto Kanzaki<sup>f</sup>, Hiroshi Hamana<sup>a</sup>

<sup>a</sup> Department of Pharmaceutical Chemistry, Faculty of Pharmacy, Chiba Institute of Science, 15-8 Shiomicho, Choshi, Chiba 288-0025, Japan

<sup>b</sup> Faculty of Pharmaceutical Sciences, Toho University, 2-2-1 Miyama, Funabashi, Chiba 274-8510, Japan

<sup>c</sup> Department of Pharmacology, Faculty of Pharmacy, Chiba Institute of Science, 15-8 Shiomi-cho, Choshi, Chiba 288-0025, Japan

<sup>d</sup> Department of Biochemistry, Graduate School of Biomedical & Health Sciences, Hiroshima University, 1-2-3 Kasumi, Minami-ku, Hiroshima 734-8553, Japan

<sup>e</sup> Department of Pharmacology, Faculty of Medicine, Kurume University, 67 Asahimachi, Kurume, Fukuoka 830-0011, Japan

<sup>f</sup> Department of Clinical Medicine, Faculty of Pharmacy, Chiba Institute of Science, 15-8 Shiomicho, Choshi, Chiba 288-0025, Japan

### ARTICLE INFO

#### Article history:

Received 18 October 2014

Revised 22 December 2014

Accepted 24 December 2014

Available online 5 January 2015

#### Keywords:

Chemical chaperone

4-Phenylbutyric acid (4-PBA)

Naphthalene

Parkinson's disease (PD)

Endoplasmic reticulum (ER) stress

### ABSTRACT

The chemical chaperone 4-phenylbutyric acid (4-PBA) has potential as an agent for the treatment of neurodegenerative diseases. However, the requirement of high concentrations warrants chemical optimization for clinical use. In this study, novel naphthalene derivatives with a greater chemical chaperone activity than 4-PBA were synthesized with analogy to the benzene ring. All novel compounds showed chemical chaperone activity, and **2** and **5** possessed high activity. In subsequent experiments, the protective effects of the compounds were examined in Parkinson's disease model cells, and low toxicity of **9** and **11** was related to amphiphilic substitution with naphthalene.

© 2015 Elsevier Ltd. All rights reserved.

Aggregation of insoluble proteins is common to neurodegenerative diseases such as Alzheimer's disease (AD) and Parkinson's disease (PD), involving phosphorylated tau and amyloid- $\beta$  (A $\beta$ ) peptides, and  $\alpha$ -synuclein, respectively. Dysfunctions associated with these aggregates have prompted the discovery of genetic predispositions and inhibitors for use as prophylactics or treatments. In particular, geldanamycin induces molecular chaperones and suppresses neurodegeneration in a drosophila model of PD.<sup>1,2</sup>

Molecular chaperones such as heat shock proteins (Hsps) are required for the functional folding of nascent proteins and the repair of higher-order protein structure. In particular, Hsp70 reduces the overexpression of A $\beta$ , which is the major component of neurofibrillary tangles in AD, and protects neuronal cells from A $\beta$ -dependent neurotoxicity in vivo and in vitro.<sup>3,4</sup> Moreover, inducible Hsp70 is protective against ischemic injury in vivo and in vitro.<sup>5</sup>

Although chemical chaperones are low molecular weight compounds, their actions are similar to those of molecular chaperones, and the well-known molecular chaperone 4-PBA prevents the





Figure 1. Strategy for synthesis of novel chemical chaperones.


aggregation of denatured proteins. In previous studies, 4-PBA has been shown to protect against cerebral ischemic injury and endoplasmic reticulum (ER) stress-induced neuronal cell death,<sup>6,7</sup> and

\* Corresponding author. Tel./fax: +81 479 30 4549.

E-mail address: smimori@cis.ac.jp (S. Mimori).



**Scheme 1.** Reagents and conditions: (a) pyridinium chlorochromate (PCC), methylene chloride, RT, 2–4 h, 64–72%; (b) ethyl diethylphosphonoacetate, NaH, THF, RT, 66–99%; (c) alkaline hydrolysis, EtOH, 10% NaOH, RT, 83–97%; (d) H<sub>2</sub>, 5% palladium carbon, AcOEt, RT, 56–90%; (e) alkaline hydrolysis, EtOH, 10% NaOH, RT, 80–98%; (f) LiAlH<sub>4</sub>, THF, RT, 1.5–3 h, 91–97%.



**Figure 2.** Structures of synthetic naphthalene analogs.

significant neuroprotective effects are shown in mouse models of AD and PD.<sup>8–10</sup> Moreover, 4-PBA has been approved for clinical use as an ammonia scavenger in children with urea cycle disorders.<sup>11</sup> Recently, 4-PBA has also shown remarkable potential as a novel therapeutic agent for type 2 diabetes<sup>12</sup> and familial hypercholesterolemia.<sup>13</sup>

Although these effects are promising, the required doses of 4-PBA are prohibitively high. Thus, we designed synthetic 4-PBA analogs to optimize chemical chaperone activity against denatured proteins.

In a recent study, we showed that chemical chaperone activities and protective effects against ER stress-induced neuronal cell death are dependent on the numbers of fatty acid carbon atoms on 4-PBA derivatives.<sup>14</sup> In further studies, we evaluated the chemical chaperone activities of synthetic methoxy-substituted 4-PBA derivatives.<sup>15</sup> Subsequently, we showed that although 4-PBA inhibits histone deacetylase (HDAC), it acts as a chemical chaperone and protects against ER stress-induced neuronal cell death.<sup>16</sup>

Previous data indicate that the benzene ring of 4-PBA is essential for chemical chaperone activity, suggesting that hydrophobic amino acids are involved in protein aggregation (Fig. 1). Thus, we assumed that the electron clouds of benzene rings interact with the aggregates of denatured proteins, and intended to expand the electron cloud area by synthesizing 4-PBA analogs based on naphthalene.

**Chemistry:** Naphthalene derivatives were synthesized according to a previous study<sup>15</sup> as shown in Scheme 1. Additional details of the synthetic process are presented in Supporting information<sup>15</sup> and all synthesized compounds are shown in Figure 2. All derivatives were characterized using spectrometric methods,<sup>17</sup> and <sup>1</sup>H NMR experiments were performed using a Unity 400 MHz spectrometer. Silica gel column chromatography was conducted using a Merck Silica gel 60 (0.040–0.063; 230–400 mesh ASTM).

**Biology:** Initially, we examined the effects of chemical chaperones on the aggregation of r-LA with BSA in vitro, as previously described.<sup>14,16</sup> All compounds inhibited the aggregation of

Download English Version:

<https://daneshyari.com/en/article/10585850>

Download Persian Version:

<https://daneshyari.com/article/10585850>

[Daneshyari.com](https://daneshyari.com)