



Contents lists available at ScienceDirect

Bioorganic & Medicinal Chemistry Letters

journal homepage: www.elsevier.com/locate/bmcl

Synthesis and biological evaluation of the pirfenidone derivatives as antifibrotic agents



Zhen Ma, Youlu Pan, Wenhai Huang, Yewei Yang, Zunyuan Wang, Qin Li, Yin Zhao, Xinyue Zhang, Zhengrong Shen*

Institute of Materia Medica, Zhejiang Academy of Medical Sciences, Hangzhou 310013, China

ARTICLE INFO

Article history:

Received 31 July 2013

Revised 30 October 2013

Accepted 15 November 2013

Available online 25 November 2013

Keywords:

Pirfenidone

Derivatives

Antifibrotic

IPF

p38

ABSTRACT

A total of 24 pirfenidone derivatives were designed, synthesized and evaluated for their inhibitory activity against the human lung fibroblast cell line MRC-5. These compounds showed the remarkable proliferation inhibition against MRC-5 compared to pirfenidone as the positive control. The possible mechanism of this kind of derivatives as antifibrotic agents was explored. The molecular docking and p38 binding affinity assays demonstrated that the antifibrotic potential of the pirfenidone derivatives was possibly through the inhibition of p38 MAPK signaling pathway. The data from this study suggested that p38 might be a potential therapeutic target for the new generation antifibrotics. All the pirfenidone derivatives are reported here for the first time.

© 2013 Elsevier Ltd. All rights reserved.

Idiopathic pulmonary fibrosis (IPF) is a typical chronic, progressive fibrosing interstitial pneumonia, primarily occurring in older adults, and limited to the lungs. It is characterized by progressive worsening of dyspnea and lung function with a poor prognosis.¹ With a median survival of 3 to 5 years and a 5-year survival rate of 20%, IPF is more mortality than most of the malignant cancers.² The pathogenesis of the IPF is complex and usually involved the interaction among various pathways driven by proinflammatory or profibrogenetic mediators.³ However, the precise mechanisms are still unknown.

More recently, pirfenidone was approved as the first antifibrotic therapy for IPF in Europe and Asia.⁴ To date, pirfenidone is the only orally administered drug that has orphan designation for the treatment of mild to moderate IPF. Its mechanism has been studied extensively and it is generally thought to be a multiple-targets drug against inflammatory, antioxidative stress and antiproliferative processes. For example, pirfenidone reduced inflammatory cytokines such as TNF- α ^{5,6} and downregulated the transcription of key profibrotic growth factors including TGF- β .^{7,8} Beneficial effects have been shown for pirfenidone in the fibrotic disease, including pulmonary, liver, renal, and cardiac muscle fibrosis.⁹

However, some side effects including gastrointestinal upset, fatigue and photosensitivity have been observed in clinical practices for pirfenidone.¹⁰ It was speculated that these adverse symptoms might due to the requirement for high customary doses in order

to counteract the reduced bioactivity by the fast metabolism in human body.¹¹ Thus, considerable efforts have been devoted towards the modification of the pirfenidone scaffold in order to increase the half-time and/or the antifibrotic activity. For example, Chen et al. substituted the 5-methyl group of pirfenidone by trifluoromethyl group or chlorine, and two series of novel (5-substituent)-2(1H)-pyridone compounds were designed with the purpose of overcoming the drawbacks of fast metabolism while maintaining its multi-targeting property. The compounds with 4-methoxy phenyl at the 1-position of pyridone scaffold of pirfenidone showed more antifibrotic activity.¹² Liu and co-workers modified 1-(substituted aryl)-5-trifluoromethyl-2(1H) pyridones with carbohydrate to prepare antifibrotic compounds with higher water solubility.¹³ Wu et al. described a N₁-substituted phenylhydroquinolinone scaffold which can be used for developing novel antifibrotic agents.¹⁴ More recent work by Kossen and co-workers demonstrated 6-oxo-7-phenyl-6,7-dihydro-1H-pyrrolo[2,3-b]pyridine derivatives for treating inflammatory and fibrotic disorders.¹⁵

In order to assess the impact of different side chain on antifibrotic potency in vitro, we introduced a side chain with 4-alkoxy phenyl group including terminal amine to the 1-position of pyridone scaffold (Fig. 1, **6a–9f**). Firstly, we focused on the length of linker between scaffold and amine. Secondly, different amines were selected according to their structure to determine the size of side chain and hydrophobic influence on the biological activity. The potential mechanism of pirfenidone derivatives as antifibrotic agents was also investigated.

* Corresponding author. Tel.: +86 571 88215506; fax: +86 571 88215625.

E-mail address: shenzr601@163.com (Z. Shen).





Figure 1. The structure of pirfenidone and compounds **6a-9f**.



Scheme 1. Reagents and conditions: (a) CuI, K_2CO_3 , DMF, reflux, 74%; (b) 5% Pd/C, THF, H_2 , rt, 78%; (c) $\text{Cl}(\text{CH}_2)_n\text{Br}$, acetone, K_2CO_3 , reflux, 90–95%; (d) NaI, K_2CO_3 , DMF, amine, 70 °C, 40–90%.

Scheme 1 shows the flow for the general preparation of these pirfenidone derivatives. The starting material **1** was obtained as reported.¹⁶ Another starting material **2** was generated by protection of 4-bromophenol with benzyl bromide.¹⁷ Using cuprous iodide as catalyst, **3** was synthesized by the Ullmann coupling reaction of **1** with **2** in dry DMF. The key intermediate **4** was successfully obtained by benzyl deprotection of **3** in THF with 5% Pd/C under H_2 atmosphere. After the reaction between intermediate **4** and alkyl dihalides at reflux temperature, compounds **5a-d** were achieved respectively. Finally, target compounds were obtained by amination reaction of **5a-d** with alkyl or heterocyclic amine.

All target compounds were tested for their cell proliferation inhibitory activity against MRC-5 cell, an ideal in vitro cell model for screening antifibrotic drugs,¹⁸ by MTT assay using pirfenidone as the positive control. DMSO was used as solvent. The results are summarized in **Table 1**.

As shown in **Table 1**, all the synthesized compounds exhibited marked proliferation inhibition against MRC-5 compared to pirfenidone as the positive control. For instance, **8a-f** showed IC_{50} from 1.77 to 3.29 mM, and **9a-f** showed IC_{50} from 1.47 to 4.35 mM, while the corresponding IC_{50} value for pirfenidone was 14.44 mM. Compound **6e** exhibited the most potent inhibitory with approximately 10-fold lower IC_{50} compared to pirfenidone.

In this Letter, we investigated the effect of the modification of substituent at the 1-position of pyridone scaffold on antifibrotic activity. At first, we focused on the length of linker between

Table 1

The structures of pirfenidone derivatives and their inhibitory activities against MRC-5 cells

Compound	n	R	IC_{50} (mM)
6a	2	$-\text{N}(\text{CH}_3)_2$	6.88
6b	2	$-\text{N}(\text{C}_2\text{H}_5)_2$	6.20
6c	2	$-\text{N}$ -pyrrolidin-1-yl	3.78
6d	2	$-\text{N}$ -piperidin-1-yl	2.48
6e	2	$-\text{N}$ -4-methyl-piperazin-1-yl	1.36
6f	2	$-\text{N}$ -morpholin-4-yl	7.64
7a	3	$-\text{N}(\text{CH}_3)_2$	5.22
7b	3	$-\text{N}(\text{C}_2\text{H}_5)_2$	2.10
7c	3	$-\text{N}$ -pyrrolidin-1-yl	3.00
7d	3	$-\text{N}$ -piperidin-1-yl	2.55
7e	3	$-\text{N}$ -4-methyl-piperazin-1-yl	4.36
7f	3	$-\text{N}$ -morpholin-4-yl	5.28
8a	4	$-\text{N}(\text{CH}_3)_2$	1.80
8b	4	$-\text{N}(\text{C}_2\text{H}_5)_2$	2.16
8c	4	$-\text{N}$ -pyrrolidin-1-yl	1.77
8d	4	$-\text{N}$ -piperidin-1-yl	3.29
8e	4	$-\text{N}$ -4-methyl-piperazin-1-yl	2.33
8f	4	$-\text{N}$ -morpholin-4-yl	3.03
9a	5	$-\text{N}(\text{CH}_3)_2$	2.98
9b	5	$-\text{N}(\text{C}_2\text{H}_5)_2$	4.35
9c	5	$-\text{N}$ -pyrrolidin-1-yl	1.47
9d	5	$-\text{N}$ -piperidin-1-yl	3.67
9e	5	$-\text{N}$ -4-methyl-piperazin-1-yl	3.41
9f	5	$-\text{N}$ -morpholin-4-yl	2.59
Pirfenidone			14.44

scaffold and amine. A series of compounds with two methylenes, three methylenes, four methylenes and five methylenes spacer were designed and synthesized. As the IC_{50} values of these new compounds were on one order of magnitude, the length of linker from two to five methylenes demonstrated no significant impact on antifibrotic activity. Furthermore, different amines were selected according to their structure to determine the size of side chain and hydrophobic influence on the biological activity. Replacement with different heterocyclic amine including pyrrolidin-1-yl group, piperidin-1-yl group, (4-methyl-piperazin)-1-yl and morpholin-4-yl group was carried out. A view on inhibitory data of compounds **6c-9f** showed that both the carbon number of heterocyclic rings and the hydrophobicity of heterocyclic amine had little effect on antifibrotic activity. For example, **7c** with a pyrrolidin-1-yl group, **7d** with a piperidin-1-yl group, **7e** with a

Download English Version:

<https://daneshyari.com/en/article/10592805>

Download Persian Version:

<https://daneshyari.com/article/10592805>

[Daneshyari.com](https://daneshyari.com)