

Novel 2-aminothiazonaphthalimides as visible light activatable photonucleases: effects of intercalation, heterocyclic-fused area and side chains

Zhigang Li,^a Qing Yang^b and Xuhong Qian^{a,c,*}

^aState Key Laboratory of Fine Chemicals, Dalian University of Technology, PO Box 158, Zhongshan Road, Dalian 116012, China

^bDepartment of Bioscience and Biotechnology, Dalian University of Technology, Dalian 116012, China

^cShanghai Key Laboratory of Chemical Biology, East China University of Science and Technology, Shanghai 200237, China

Received 24 December 2004; revised 15 February 2005; accepted 17 February 2005

Abstract—A new family of 2-aminothiazonaphthalimides with different side chains as novel intercalative and visible light activatable photonucleases, was designed, synthesized and quantitatively evaluated. The order of their photocleaving abilities was parallel to that of their intercalative properties. The compound with linear heterocyclic-fused chromophore could intercalate into and photocleave DNA more efficiently than the one with angular heterocyclic-fused chromophore. **B**₂, the most efficient compound, caused obvious DNA damage at 1 μM. Mechanism experiment showed that superoxide anion was involved.

© 2005 Elsevier Ltd. All rights reserved.

Photonucleases hold great promise in molecular biology and medicinal chemistry. Some of them have been proven to be valuable tools in biotechnology including therapeutic agents,¹ structural probes and photofootprinting agents as well.² Triggered by ultra-violet or visible light, such agents can cause significant damage on DNA with or without site selectivity.³

Many naphthalimide derivatives were well-known photonucleases³ or anticancer drugs.⁴ Our previous study has demonstrated that the presence of sulfur rather than its oxo-counterpart promoted the DNA intercalating and photocleaving ability.^{3f} Based on this knowledge, we expected fusing novel thio-heterocyclic group into the naphthalimide skeleton would be an efficient strategy.

2-Aminothiazole moiety, proven its value in medicinal chemistry, has been successfully applied in dopamine agonists, such as B-HT920, PD118440 and pramipexole, the widely used anti-Parkinsonian agent.⁵ In these cases, 2-aminothiazole group was assumed to interact with nu-

cleic acid to form a stable DNA–drug complex with the aid of hydrogen bonds formed between the amino group and the sugar–phosphate chain. Herein, a new family of 2-aminothiazonaphthalimides, **A**_{1–2} and **B**_{1–2} (Fig. 1), was synthesized by fusing a 2-aminothiazole group to different positions of the naphthalimide skeleton. Two aminoalkyl side chains varying in length were incorporated into the parent structures not only to increase the affinity interaction with DNA^{4,6} but also to elucidate the influence of them on biological activities. Furthermore, the comparison of effects of structural features of such agents on photocleaving activities could offer novel insights for rational molecular design.





Figure 1. Structures of novel 2-aminothiazonaphthalimides.

Keywords: Plasmid DNA; Photocleaver; Intercalation; Heterocycles.

* Corresponding author. Tel.: +86 411 83673466; fax: +86 411 83673488; e-mail addresses: xhqian@dlut.edu.cn; xhqian@ecust.edu.cn



Scheme 1. Synthesis of 2-aminothiazonaphthalimides. Reagents and conditions: (a) NaNO_3 , H_2SO_4 , 0°C , 1 h, 85% yield; (b) SnCl_2 , concentrated HCl , 80°C , 2 h, 80% yield; (c) KSCN , Br_2 , AcOH , 48 h, 95% yield; (d) corresponding amine, ethanol, reflux 2 h.

The synthesis of these compounds was outlined in **Scheme 1**. Nitration of the starting material, 1,8-naphthalic anhydride with sodium nitrate in concentrated H_2SO_4 at 0°C for 1 h, 3-nitro-1,8-naphthalic anhydride was obtained and then reduced by Tin(II) chloride dehydrate in concentrated hydrochloric acid to give 3-amino-1,8-naphthalic anhydride. Upon cyclization of 3-amino-1,8-naphthalic anhydride by a reported procedure,⁷ in which 3-amino-1,8-naphthalic anhydride was treated with potassium thiocyanate and bromine in acetic acid, two 2-aminothiazonaphthalic anhydride isomers were obtained by filtration in high yield. Without further purification, the obtained anhydride mixtures were condensed with the corresponding amine in ethanol to form the corresponding naphthalimide mixtures, **A**_{1–2} and **B**_{1–2}, with the ratio of 1:10. After separation with careful column chromatography, each pure targeted product was obtained. All of their structures were confirmed by ^1H NMR, HRMS and IR.⁸

The UV–vis and fluorescent data for these compounds were measured and shown in **Table 1**. It was found that they had slight differences in both absorption wavelength (around 420 nm) and emission wavelength (around 475 nm). The activating factor was visible light (420 nm) rather than UV light, indicating the safe manipulation towards them and thus in turn their prevalent advantage as biological tools.

The Scatchard binding constants between compounds, **A**₁, **A**₂, **B**₁, **B**₂ and CT-DNA (in 30 mM Tris–HCl buffer, pH 7.5) were determined using fluorescence technique method (**Fig. 2**)⁹ and were shown in **Table 1**. The intercalating abilities of **B**_n were stronger than those of their isomers, **A**_n ($n = 1, 2$), indicating that the compound with linear heterocyclic-fused chromophore was more easily intercalated into DNA than the one with angular chromophore. Furthermore, the intercalating abilities of




Figure 2. Fluorescence spectra before and after interaction of compound **B**₂ and calf thymus DNA. Curves F and F-CT corresponded to compound **B**₂ before and after being mixed with DNA. Numbers 1–4 indicated the concentration of **B**₂, 5, 10, 20, 40 μM , respectively. DNA applied was 50 μM (bp).

X₂ were stronger than those of **X**₁ (**X** = **A**, **B**), suggesting the importance of the length of aminoalkyl chain serving as DNA groove binder and/or external electrostatic binder. In our cases, the chain with three methylene units between two nitrogen atoms significantly enhanced the intercalating abilities of the aminothiazonaphthalimides. That might be the reason accounting for the order of intercalating ability: **A**₂ > **A**₁, **B**₂ > **B**₁.

The photocleavage activities of **A**₁, **A**₂, **B**₁, **B**₂, were evaluated using closed supercoiled pBR322 DNA under photo-irradiation with a transilluminator (400 nm) at a distance of 20 cm at 0°C for 3 h under aerobic conditions and analyzed on a 1% agarose gel. The cleavage efficiency was defined by the degree of conversion of the supercoiled pBR322 DNA (form I) to relaxed circular DNA (form II). It was apparent that these compounds photocleaved the supercoiled DNA with different activities by the order: **B**₂ > **A**₂ > **B**₁ > **A**₁ (**Fig. 3a**), which was parallel to that of their intercalative properties. Further the experiment showed that **B**₂, the most efficient compound, nicked supercoiled DNA (form I, DNA in 200 μM base pair) to give a 29% relaxed circular form (form II) at a concentration as low as 1 μM , while the concentration for giving 100% form II (**Fig. 3b**) was 100 μM under identical conditions. Also the cleaving activity of **B**₂ increased remarkably with prolonging photo-irradiation time (**Fig. 3c**), indicating it was a time-dependent process. As no damage was observed in the absence of light (lane 3), it was thus believed that the visible light functioned as a trigger to initiate the DNA strand scission. These newly designed naphthalimides possessing 2-aminothiazole showed much high photobiological activities by comparison with their parent compound without 2-aminothiazole.^{3f}

Table 1. Spectral data,^{a,b} Scatchard binding constants of these compounds

Compds	UV λ_{max} (lg ϵ)	FL λ_{max} (Φ)	Scatchard binding constants ($\times 10^4 \text{ M}^{-1}$)
A ₁	421 (3.50)	475.5 (0.003)	1.98
A ₂	422 (3.73)	478.3 (0.004)	2.87
B ₁	421 (3.72)	475.5 (0.003)	2.56
B ₂	421 (3.81)	478.4 (0.004)	4.73

^a In absolute DMSO.

^b With quinine sulfate in sulfuric acid as quantum yield standard ($\Phi = 0.55$).

Download English Version:

<https://daneshyari.com/en/article/10598122>

Download Persian Version:

<https://daneshyari.com/article/10598122>

[Daneshyari.com](https://daneshyari.com)