



# Extraction optimization and bioactivity of polysaccharides from *Aspergillus fumigatus* AF1



Xianchun Jin<sup>a,\*</sup>, Yu Ning<sup>b</sup>

<sup>a</sup> School of Sciences, Henan Agricultural University, 450002, China

<sup>b</sup> School of Life Sciences, Henan Agricultural University, 450002, China

## ARTICLE INFO

### Article history:

Received 15 February 2013

Received in revised form 30 March 2013

Accepted 10 April 2013

Available online 17 April 2013

### Keywords:

Polysaccharides

Response surface methodology

*Aspergillus fumigatus*

Antitumor

## ABSTRACT

*Aspergillus fumigatus* was used to produce polysaccharide using sodium hydroxide pretreated rice straw as the sole carbon source by submerged fermentation. Response surface methodology (RSM) was used to optimize the extraction of polysaccharide from the mycelia of *A. fumigatus* (PAF). A three-level, three-factor Box–Behnken design (BBD) was applied for experimental design and analyzed the results to obtain the optimal extraction parameters. RSM analysis indicated good correspondence between experimental and predicted values. The optimal conditions for polysaccharide extraction were extraction temperature 50 °C, extraction time 6 h, ratio of dried mycelia mass to water 0.03. PAF showed excellent antitumor activity *in vitro* and exert no damage to the normal cells. PAF also showed prominent antitumor activity *in vivo* against S180 and increased the spleen and thymus index significantly.

© 2013 Elsevier Ltd. All rights reserved.

## 1. Introduction

In recent years, a plethora of polysaccharides have been isolated from the microorganisms by extraction (Chen, Wang, Zhang, & Huang, 2012; Gan, Ma, Jiang, Xu, & Zeng, 2011) as many polysaccharides showed a variety of bioactivities such as antitumor, antioxidant and anti-inflammatory effects (Cengiz, Bektas, & Mustafa, 2008; Dentinger, Ammirati, & Both, 2010; He et al., 2012; Mao, Mao, & Meng, 2013; Smiderle et al., 2012). Many polysaccharides have been extracted from various edible fungi such as *Lentinus edodes* (Zhang et al., 2010), *Flammulina velutipes* (Shi, Yang, Guan, Zhang, & Zhang, 2012), *Gomphidius rutilus* (Wang, Zhang, Wang, & Wang, 2013). However, there is few reports about the bioactivity of *Aspergillus fumigatus* (Costachel, Coddeville, Latgé, & Fontaine, 2005), as it is usually be thought as the most prevalent airborne fungal pathogen causing fatal invasive infections in immunocompromized patients over the past decade (Latgé, 1999) though Tekaiia and Latgé thought that *A. fumigatus* should be a saprophyte (Tekaiia & Latgé, 2005).

In recent years, there has been an unprecedented increase in interest in the more efficient utilization of the agro-residues because their application provides an alternative way to reduce the production costs and solve many environmental hazards (Qijun et al., 2008). Rice straw, a by-product of rice production, is discharged as an agricultural waste. 18.4 million tons of rice straw

were produced every year in China (Bi, Wang, Wang, Gao, & Wang, 2011), most of which was incinerated, causing much serious air pollution. With the strengthening of environmental legislation, incinerating straw was banned, alternative method of using straw should be found. Using straw to produce enzymes and other valuable products may be such alternative. In the present study, *A. fumigatus* was used to degrade NaOH pretreated rice straw to produce polysaccharide from its mycelia by submerged fermentation as submerged cultures of fungi can give rise to higher productions of mycelial biomass and EPS in a more compact space, within a shorter period of time, and with less chance for contamination (Cho, Oh, Chang, & Yun, 2006). At the same time, response surface methodology was used to optimize the polysaccharide extraction conditions, and the antitumor activity of the polysaccharide both *in vitro* and *in vivo* was also studied.

## 2. Materials and methods

### 2.1. Microorganism and culture conditions

*A. fumigatus* AF1 was isolated from moldy rice straw collected from Chenzhou China. The preservation and sporulation of the strain was done on PDA. To prepare the inoculum, spores from the slant were suspended in 2 mL of 0.9% NaCl ( $10^7$ – $10^8$  spores/mL) and transferred into a 250 mL conical flask containing 50 mL of modified Czapek medium (g/L): sucrose 10, FeSO<sub>4</sub> 0.01, KCl 0.5, MgSO<sub>4</sub>·7H<sub>2</sub>O 0.5, K<sub>2</sub>HPO<sub>4</sub> 1.0, NaNO<sub>3</sub> 3.0. Fungal cells were cultured in an orbital shaker (150 rpm) at 40 °C for 20 h before the mycelium pellets was used for inocula. The composition of

\* Corresponding author. Tel.: +86 377 63554844; fax: +86 377 63558130.

E-mail addresses: [xchjin@163.com](mailto:xchjin@163.com), [xchjin@gmail.com](mailto:xchjin@gmail.com) (X. Jin).

**Table 1**  
Coded values and experimental range of extraction variables used in experimental design.

| Variables                              | Symbols | Coded level of variable |       |      |
|----------------------------------------|---------|-------------------------|-------|------|
|                                        |         | –1                      | 0     | 1    |
| Extraction temperature (°C)            | $X_1$   | 50                      | 70    | 90   |
| Ratio of dried mycelia to water (g/mL) | $X_2$   | 0.03                    | 0.065 | 0.10 |
| Extraction time (h)                    | $X_3$   | 2                       | 4     | 6    |

fermentation medium was 2.0 g of different concentration of NaOH solution pretreated rice straw (2.0 g milled rice straw was mixed with 38.0 mL NaOH solution, autoclaved at 121 °C for 15 min. After cooling to room temperature, the reaction mixture was adjusted to given pH with 10.0 mol/L sulfuric acid and then 50% (w/v) separately autoclaved ammonium sulfate solution was added as the nitrogen source (1% of final concentration)) in 250 mL conical flask. Five milliliters of *A. fumigatus* AF1 was inoculated into each 250 mL flask then cultured at 40 °C for 24 h.

## 2.2. Extraction of crude polysaccharides from *A. fumigatus* AF1 mycelia

As the fermentation medium turned to a light yellow liquid in 24 h, *A. fumigatus* mycelia obtained by filtration with a sintered glass crucible (G-3) was washed thoroughly with distilled water. The mycelia were dried at 30 °C. Then, it was grinded and stored in desiccators at room temperature until use. The dried mycelia powders were extracted with distilled water in a 100 mL centrifuge tubes. The centrifuge tube was held in a water bath and exposed to extract polysaccharides for a different time at varying temperatures in different ratios of dried mycelia to water.

## 2.3. Determination of the yield of polysaccharides from *A. fumigatus* AF1 mycelia

After extraction for a given time, the extracted slurry was centrifuged at  $10,000 \times g$  for 10 min, and the supernatant was collected and concentrated under reduced pressure at 40 °C. The solution was precipitated with three volumes of absolute ethanol, and the precipitate was collected by filtration and redissolved in distilled water. After removal of proteins by the Sevag method, the polysaccharide solution was dialyzed extensively against distilled water and lyophilized. Prior to use, the polysaccharide was dissolved in water and diluted into a series of concentrations as indicated. The carbohydrate content of the solution was determined by the phenol–sulfuric acid method. The isolated polysaccharide was referred to as PAF.

## 2.4. Experimental design

Based on the preliminary results, a three-level–three-factor BBD was used to determine the best combination variables for extraction of polysaccharides from *A. fumigatus* mycelia. Table 1 shows the range of each factor.

Quadratic models considered as response surface model for predicting the optimal points were expressed according to Eq. (1)

$$Y = \beta_0 + \beta_1 X_1 + \beta_2 X_2 + \beta_3 X_3 + \beta_{12} X_1 X_2 + \beta_{13} X_1 X_3 + \beta_{23} X_2 X_3 + \beta_{11} X_1^2 + \beta_{22} X_2^2 + \beta_{33} X_3^2 \quad (1)$$

where  $Y$  is the dependent variable,  $\beta_0$  is the model constant,  $\beta_i$ ,  $\beta_{ij}$  and  $\beta_{ij}$  are the model coefficients. They represent the linear, quadratic and interaction effects of the variables. Analysis of the experimental design data and calculation of predicted responses

was carried out using Design Expert software (version 7.0.0, USA). Additional confirmation experiments were subsequently conducted to verify the validity of the statistical experimental design.

## 2.5. Antitumor activity of PAF

### 2.5.1. In vitro antitumor activity assay

Inhibition activities *in vitro* of crude PAF on such cell lines as human gastric cancer Sarcoma180, human hepatoma cell line (HepG2), human breast adenocarcinoma cell line (MCF-7) and normal human liver cell line (L-02) were evaluated using the MTT method. Briefly, cells in a density of  $1.0 \times 10^5$  cells/mL in the RPMI-1640 medium supplemented with 10% FBS, 100 U/mL penicillin and 100 µg/mL streptomycin were pipetted into 96-well flat-bottom plate (100 µL/well). After 12 h of incubation at 37 °C in a humidified 5% CO<sub>2</sub> incubator, non-adherent cells were removed by washing three times with RPMI-1640 medium. Then, fresh medium (100 µL/well, control group) or test sample (100 µL/well, crude PAF at different final concentration) was added to each well, and the cells were incubated for 24 h. After incubation, MTT solution (10 µL/well, 5 mg/mL) was added to each well, and the plate was incubated for an additional 4 h at 37 °C. Finally, 100 µg/L of 10% SDS in 0.01 M HCl was added to each well and the plate was kept overnight for the dissolution of formazan crystals. The absorbance of each well at 570 nm was measured with an ELISA plate reader (BioTek Instruments Inc., Winooski, VT, USA). The inhibition rate was calculated according to the following formula:

$$\text{Inhibition rate (\%)} = \frac{(A_b - A_s)}{A_b} \times 100$$

where  $A_s$  and  $A_b$  are the absorbance of treated cells and untreated cells.

### 2.5.2. In vivo antitumor activity assay

Animal care and handling were according to the Committee for the Purpose of Control and Supervision of Experiments on Animal's guidelines after research project approval by the Institutional Animal Ethics Committee.

BALB/C mice (male, 7–9 weeks old, weighing  $20 \pm 1.0$  g) were purchased from the Animal Center of Tongji Medical College, Huazhong University of Science and Technology, Wuhan, China.

Each of polysaccharides sample as well as control agents was tested on an individual group of animals consisting of 6 BALB/C mice.

Sarcoma180 tumor cells ( $5 \times 10^6$  cells/mL) were transplanted subcutaneously into the right groin of the mice. After 24 h of tumor inoculation, PAF samples were dissolved in 0.9% aqueous NaCl, then injected intraperitoneally to mice in experimental group every other day for 20 days. The equivalent volume of 0.9% aqueous NaCl was injected intraperitoneally in the negative control group. The mice were killed on the next day of the last injection, and the tumors, spleens, thymuses and livers were excised and weighed.

The inhibition ratio  $\xi$  and enhancement ratio of body weight  $f$  were calculated as follows:

$$\xi = \left[ \frac{(W_c - W_t)}{W_c} \right] \times 100\%$$

$$f = \left[ \frac{(W_a - W_b)}{W_b} \right] \times 100\%$$

$W_c$  is the average tumor weight of the negative control group.  $W_t$  is the average tumor weight of PAF group;  $W_b$  and  $W_a$  are the body weight of mice before and after the assay.

The indexes of spleen, liver and thymus were defined as the ratio of corresponding organ's weight (mg) to the body weight (g).

Download English Version:

<https://daneshyari.com/en/article/10601514>

Download Persian Version:

<https://daneshyari.com/article/10601514>

[Daneshyari.com](https://daneshyari.com)