



Short communication

Alcohol consumption patterns in HIV-infected adults with alcohol problems

Nicolas Bertholet^{a,b,c,*}, Debbie M. Cheng^{a,d}, Jeffrey H. Samet^{a,e}, Emily Quinn^f, Richard Saitz^{a,g}^a Clinical Addiction Research and Education (CARE) Unit, Section of General Internal Medicine, Department of Medicine, Boston Medical Center and Boston University School of Medicine, 801 Massachusetts Avenue, 2nd Floor, Boston, MA 02118-2605, United States^b Clinical Epidemiology Center, Institute of Social and Preventive Medicine, Centre Hospitalier Universitaire Vaudois and University of Lausanne, Bugnon 17, Lausanne 1005, Switzerland^c Alcohol Treatment Center, Centre Hospitalier Universitaire Vaudois and University of Lausanne, Mont-Paisible 16, Lausanne 1005, Switzerland^d Department of Biostatistics, Boston University School of Public Health, 801 Massachusetts Avenue, 3rd Floor, Boston, MA 02118-2605, United States^e Department of Social and Behavioral Sciences, Boston University School of Public Health, 801 Massachusetts Avenue, 4th Floor, Boston, MA 02118-2605, United States^f DM-STAT Inc., 1 Salem Street Malden, MA 02148-5236, United States^g Youth Alcohol Prevention Center, Boston University School of Public Health, 580 Harrison Avenue, 3rd Floor, Boston, MA 02118-2605, United States

ARTICLE INFO

Article history:

Received 23 October 2009

Received in revised form 16 May 2010

Accepted 25 May 2010

Available online 25 June 2010

Keywords:

Alcohol

HIV

Drinking patterns

ABSTRACT

Objective: To understand patterns of alcohol consumption and baseline factors associated with favorable drinking patterns among HIV-infected patients.**Methods:** We studied drinking patterns among HIV-infected patients with current or past alcohol problems. We assessed drinking status in 6-month intervals. Based on National Institute on Alcohol Abuse and Alcoholism guidelines a favorable drinking pattern was defined as not drinking risky amounts at each assessment or decreased drinking over time. All other patterns were defined as unfavorable. Logistic regression models were used to identify baseline factors associated with a favorable pattern.**Results:** Among 358 subjects, 54% had a favorable drinking pattern with 44% not drinking risky amounts at every assessment, and 11% decreasing consumption over time. Of the 46% with an unfavorable pattern, 4% drank risky amounts each time, 5% increased, and 37% both decreased and increased consumption over time. Current alcohol dependence and recent marijuana use were negatively associated with a favorable pattern, while older age and female gender, and having a primary HIV risk factor of injection drug use were positively associated with a favorable pattern.**Conclusion:** Many HIV-infected adults with alcohol problems have favorable drinking patterns over time, and alcohol consumption patterns are not necessarily constant. Identifying HIV-infected adults with a pattern of risky drinking may require repeated assessments of alcohol consumption.

© 2010 Elsevier Ireland Ltd. All rights reserved.

1. Introduction

Alcohol use is common among human immunodeficiency virus (HIV)-infected adults and has negative health consequences. Over a third of HIV-infected veterans drink amounts associated with health consequences (Samet et al., 2007a,b). In another study, almost half of HIV-infected patients initiating primary care reported a high probability of having an alcohol use disorder (had two of more positive answers to the CAGE screening tool) (Mayfield et al., 1974; Samet et al., 2004b). Unhealthy alcohol use (the spectrum from drinking risky amounts through alcohol dependence) is more prevalent among HIV-infected patients than it is in the general population (Lefevre et al., 1995; Conigliaro et al., 2003; Samet et al., 2004b; Chander et al., 2006). Drug and unhealthy alcohol use have been linked to HIV-disease progression, HIV risk behaviors,

and decreased adherence to antiretroviral therapy (Samet et al., 2007a,b). In adults with alcohol use disorders, social and personal factors (e.g., male gender, major depression, heroin use, cocaine use, divorce, and less education) have been linked to a worse prognosis (Moss et al., 2007). The environment has an impact on the course of drinking too (e.g., social pressure to drink has been linked with relapse among individuals with alcohol dependence) (Zywiak et al., 2006a,b).

There is sufficient evidence to date (though not specifically among those with HIV infection) to conclude that brief counseling in the primary care setting for nondependent unhealthy alcohol use can lead to a decrease in drinking (Bertholet et al., 2005; Kaner et al., 2007), including among injection drug users (Stein et al., 2002). Nevertheless, alcohol use disorders are often chronic conditions characterized by recurrent episodes, and few prospective studies have explored alcohol consumption over time. Since the treatment of HIV infection requires longitudinal care, it gives clinicians repeated opportunities to address alcohol use. A greater understanding of consumption over time in HIV-infected individuals with alcohol use disorders could help clinicians and researchers

* Corresponding author at: Alcohol Treatment Center, Mont-Paisible 16, 1011 Lausanne, CHUV, Switzerland. Tel.: +41 21 314 73 51; fax: +41 21 314 05 62.

E-mail address: nberthol@bu.edu (N. Bertholet).

better address these problems. Therefore, in a prospective cohort of HIV-infected adults with current or past alcohol problems (HIV-Longitudinal Interrelationships of Viruses and Ethanol [HIV-LIVE]), we studied patterns of alcohol consumption and factors associated with those patterns.

2. Methods

Participants were recruited between August 2001 and July 2003 with follow-up every 6 months through 2005. Recruitment occurred from a previous cohort study (HIV-Alcohol Longitudinal Cohort), an intake clinic for HIV-infected patients, HIV primary care and specialty clinics, a homeless respite facility, a methadone program, study flyers, and referrals from physicians, other participants and social service agencies (Samet et al., 2004a). Individuals were eligible if they had a documented HIV antibody test by ELISA (confirmed by Western blot) and ≥ 2 positive answers to the CAGE alcohol screening questionnaire (Mayfield et al., 1974) or an alcohol use disorder diagnosis (lifetime) by a study physician clinical assessment. Participants were fluent in either English or Spanish. Exclusion criteria were cognitive impairment (score of < 21 on the Mini-Mental State Examination) and inability to provide informed consent (Folstein et al., 1975; Smith et al., 2006). The study was approved by the Institutional Review Boards of Boston Medical Center and Beth Israel Deaconess Medical Center. Study participants who attended the baseline assessment and at least 2 follow-up visits (total of 3 or more assessments) were included in this analytic sample.

2.1. Outcome

The primary outcome of this study was having a favorable drinking pattern over time. At baseline and at each follow-up visit, alcohol consumption was assessed using a validated calendar method (30-day Timeline FollowBack) (Sobell and Sobell, 1995). Participants were classified at each assessment as abstinent, drinking below or drinking above risky amounts (as defined by the National Institute on Alcohol Abuse and Alcoholism [≥ 5 drinks/occasion or > 14 drinks/week for men; 4 or 7 drinks, respectively, for women and persons aged 65 and over]). Longitudinal drinking patterns were summarized as favorable or unfavorable. A favorable drinking pattern was defined as not drinking risky amounts at every observed study visit (i.e., abstinent or consistently drinking below risky amounts) or a decrease in the observed drinking over time (e.g., from risky to not risky) with no observed increases. All other drinking patterns were defined as unfavorable (i.e., consistently drinking risky amounts, increase in drinking from not risky to risky amounts, or intermittent risky drinking).

In addition, since it is uncertain whether risky amounts as defined in the general population are appropriate definitions for HIV-infected adults, we evaluated the secondary outcome continuous abstinence, defined as reporting abstinence at every study visit.

2.2. Factors associated with drinking patterns

Factors of interest and potential confounders of these associations were assessed at baseline (defined *a priori* based on published literature and clinical experience). Marital status, homelessness (Kertesz et al., 2003), age, gender, race/ethnicity, primary HIV risk factor at the time of infection (injection drug use, men sex with men, heterosexual sex), recent heroin, cocaine and marijuana use (past 12 months), any attendance at Alcoholics Anonymous (AA) meetings (past 6 months), and whether or not the individual spent time with people who drink alcohol (a measure of social pressure to drink) were self-reported. Health-related quality of life was summarized using the Mental Component Summary (MCS) and Physical Component Summary (PCS) scores of the 12-item Short-Form Health Survey (Delate and Coons, 2000). Past 6-month and lifetime diagnosis of alcohol dependence were assessed using the Composite International Diagnostic Interview (World Health Organization, 1996).

We hypothesized that marital status, homelessness, AA attendance, and lack of social pressure to drink would be associated with more favorable drinking pattern; worse health-related quality of life and recent drug use would be associated with an unfavorable pattern. Other variables were considered potential confounders.

2.3. Analysis

We determined the frequency and proportion of each drinking pattern based on the observed data for each person. Multiple logistic regression models were fit to identify baseline factors associated with a favorable alcohol consumption pattern across time. The models adjusted for all factors of interest and potential confounders, none of which were highly correlated (Spearman $r > 0.40$). The Hosmer–Lemeshow test was used to assess model goodness-of-fit. All analyses were conducted using two-sided tests and a significance level of 0.05. Analyses were performed using SAS software (version 9.1; SAS Institute, Cary, NC).

Table 1

Baseline characteristics of a prospective cohort of adults with HIV-infection and current or past alcohol problems [$n = 358$].

Age, mean (SD)	41.6 (7.4)
Female, n (%)	89 (24.9)
Race/ethnicity, n (%)	
Black	157 (43.9)
White	119 (33.2)
Latino	63 (17.6)
Other	19 (5.3)
Homelessness, n (%)	84 (23.5)
Currently married, n (%)	22 (6.1)
Mental Component Summary score (MCS), mean (SD)	39.8 (11.4)
Physical Component Summary score (PCS), mean (SD)	43.5 (9.8)
Recent drug use (past 12 months), n (%)	
Heroin use	88 (24.6)
Cocaine use	163 (45.5)
Marijuana use	133 (37.2)
Drinks per day, last 30 days, mean (SD)	1.8 (5.5)
Attended AA meeting, past 6 months, n (%)	203 (56.7)
Alcohol dependence diagnosis (current), n (%)	35 (9.8)
Alcohol abuse diagnosis (lifetime), n (%)	61 (17.0)
Alcohol dependence diagnosis (lifetime), n (%)	245 (68.4)
Baseline drinking status, n (%)	
At risk drinking	114 (31.8)
“Moderate” drinking	41 (11.5)
Abstinent	203 (56.7)
Spending time with people who drink alcohol (Social pressure to drink alcohol), n (%)	219 (61.2)
Primary HIV risk factor, at the time of infection, n (%)	
Injection drug use	184 (51.5)
Men sex with men	82 (23.0)
Heterosexual sex	91 (25.5)

SD = Standard deviation. Homelessness was defined as at least one night on the street or in a shelter in the past 6 months. AA = Alcoholics Anonymous. Alcohol consumption (drinks per day) was assessed using the Timeline FollowBack method. The presence of a current (past 6 months) alcohol dependence, and a lifetime diagnosis of alcohol abuse or dependence was assessed using the Composite International Diagnostic Interview (CIDI). Risky drinking was defined as > 14 drinks/week or ≥ 5 drinks on an occasion for men, > 7 drinks/week or ≥ 4 drinks on an occasion for women and persons \geq age 65 years. Moderate drinking was defined as drinking alcohol but below risky drinking limits. A drink was defined as 12–14 g of ethanol, as in the amount in the U.S. in one 12 ounce beer, one 5 ounce glass of wine, or 1.5 ounces of 80 proof liquor. Social pressure to drink alcohol: Subjects were asked how many of the people they spend time with currently drink alcohol (none, a few, about half, most, all). Answers were later dichotomized (none vs. other).

3. Results

Of the 400 participants in the HIV-LIVE prospective cohort study, 358 (90%) completed at least 3 study visits and were included in the present analyses. Median follow-up was 3.4 years. The baseline characteristics of the study sample are presented in Table 1. Over the course of follow-up, 54.5% had a favorable drinking pattern with 43.9% consistently drinking below risky limits at every assessment (70.7% of the latter group abstained), and 10.6% decreasing consumption over time. Of the 45.5% with an unfavorable drinking pattern, 3.6% drank risky amounts at each assessment, 4.7% increased their drinking to risky amounts and 37.2% had intermittent risky drinking.

In the logistic regression model (Table 2), among the factors of interest, recent marijuana use had a negative association with a favorable drinking pattern. Current alcohol dependence (past 6 months) was also negatively associated with an unfavorable pattern. Older age, female gender and having a primary HIV risk factor of injection drug use at the time of infection (compared to heterosexual sex) were associated with a favorable drinking pattern.

Older age and female gender were also associated with continuous abstinence while marijuana use, cocaine use, social pressure to

Download English Version:

<https://daneshyari.com/en/article/1070524>

Download Persian Version:

<https://daneshyari.com/article/1070524>

[Daneshyari.com](https://daneshyari.com)