

Brief original article

Incidence and temporal trends of childhood type 1 diabetes between 1975 and 2012 in Navarre (Spain)



Luis Forga Llenas^{a,*}, María José Goñi Iriarte^a, Koldo Cambra Contin^b, Berta Ibáñez Beroiz^b, María Chueca Guendulain^c, Sara Berrade Zubiri^c

^a Endocrinology Department, Complejo Hospitalario de Navarra, 31008 Pamplona, Navarra, Spain

^b Navarrabiomed, Fundación Miguel Servet, Red de Investigación en Servicios Sanitarios en Enfermedades Crónicas (REDISSEC), 31008 Pamplona, Navarra, Spain

^c Paediatric Endocrinology, Complejo Hospitalario de Navarra, 31008 Pamplona, Navarra, Spain

ARTICLE INFO

Article history:

Received 23 April 2014

Accepted 17 June 2014

Available online 4 August 2014

Keywords:

Type 1 diabetes

Incidence

Trend

Epidemiology

ABSTRACT

Objective: To determine trends in the incidence of type 1 diabetes in Navarre (Spain) between 1975 and 2012 by age and sex.

Patients and methods: The study population comprised residents of Navarre under 15 years of age. A Poisson regression model was fitted to analyze changes in the incidence over time, adjusted by year of diagnosis, age group and sex.

Results: A total of 494 patients were registered, representing an adjusted incidence rate of 13.2/100,000 person-years. The annual relative increase in the incidence rate was 3.7%. The highest incidence was found in the group aged 10–14 years. The incidence among boys aged 10–14 tended to be higher than that in girls of the same age.

Conclusions: Since the year 2000, the incidence of type 1 diabetes among persons younger than 15 years in Navarre has been very high and has quadrupled over the last four decades.

© 2014 SESPAS. Published by Elsevier España, S.L.U. All rights reserved.

Incidencia y tendencia temporal de la diabetes tipo 1 en la infancia, entre 1975 y 2012, en Navarra (España)

RESUMEN

Objetivo: Determinar la tendencia en la incidencia de diabetes tipo 1 en Navarra entre 1975 y 2012 por edad y sexo.

Pacientes y métodos: La población objeto de estudio comprende a los residentes en Navarra menores de 15 años de edad. Para analizar la evolución de la incidencia a lo largo del tiempo, se ha utilizado un modelo de regresión de Poisson ajustado por año de diagnóstico, grupo de edad y sexo.

Resultados: Se han diagnosticado 494 pacientes, lo que supone una incidencia ajustada de 13,2/100.000 personas-año. El incremento relativo anual en la tasa de incidencia ha sido del 3,7%. El grupo de edad con mayor incidencia fue el de 10 a 14 años. En este mismo grupo, la incidencia en niños tiende a ser mayor que en niñas.

Conclusiones: Desde el año 2000, la incidencia de diabetes tipo 1 en menores de 15 años, en Navarra, es muy alta y se ha cuadruplicado en las últimas cuatro décadas.

© 2014 SESPAS. Publicado por Elsevier España, S.L.U. Todos los derechos reservados.

Palabras clave:

Diabetes tipo 1

Incidencia

Tendencia

Epidemiología

Introduction

The incidence of type 1 diabetes (T1D) in children under 15 ranges from 0.1/100,000 person-years in Venezuela to 57.6/100,000 in Finland.¹ This variation could reflect differences in genetic susceptibility between populations, different exposure to environmental risk factors² or differences in features and quality registers.³ In Spain, the incidence is very high (20.6)¹ and, between

Autonomous Communities, the highest incidence rate is twice that of the lowest.³

The highest incidence appears in the 10–14 age group, coinciding with puberty.⁴ In some populations, however, the 5–9 age group has the highest incidence.⁵ Furthermore, a shift to younger age of onset has been described⁶ and, in some countries, such as Sweden,⁷ Germany⁸ and Belgium,⁹ it has been debated whether there has truly been an increase in incidence in recent years, or just an earlier onset of the disease.

Regarding the differences in incidence between the sexes before the age of 15, it appears that in countries with a high overall incidence of T1D, it is more common in boys, whereas in low-incidence countries, it is higher among girls.¹⁰ In some registers, no difference

* Corresponding author.

E-mail addresses: lfor gall@cfnavarra.es, luisforgallenas@gmail.com (L. Forga Llenas).

Table 1
Incidence of type 1 diabetes (number of cases per 100,000, per year) by age and sex. Navarre 1975–2012.

| Group age (years) | Cases | People/year | Incidence | 95% CI | <i>p</i> ^a |
|-------------------|-------|-------------|-----------|-----------|-----------------------|
| 0–4 | 94 | 1,156,439 | 8.1 | 6.5–9.7 | 0.607 |
| Male | 46 | 596,559 | 7.7 | 5.5–9.9 | |
| Female | 48 | 559,880 | 8.5 | 6.1–11.0 | |
| 5–9 | 148 | 1,225,186 | 12.0 | 10.1–14.0 | 0.343 |
| Male | 82 | 631,106 | 12.9 | 10.2–15.8 | |
| Female | 66 | 594,080 | 11.1 | 8.4–13.8 | |
| 10–14 | 252 | 1,302,921 | 19.3 | 16.7–21.7 | 0.051 |
| Male | 145 | 669,793 | 21.6 | 18.1–25.1 | |
| Female | 107 | 633,128 | 16.9 | 13.7–20.1 | |
| <15 | 494 | 3,684,546 | 13.4 | 12.2–14.6 | 0.094 |
| Male | 273 | 1,897,458 | 14.3 | 12.7–16.1 | |
| Female | 221 | 1,787,088 | 12.3 | 10.7–14.0 | |

^a Results of the test comparing incidence between males and females.

is found between the sexes⁹ or, where these differences appear, they start at the age of ten.¹¹

In order to determine whether there has been an increase in the incidence and/or a change in the age of onset in Navarre, this paper describes the changes over the past four decades in the incidence in children under 15 years, including characteristics in terms of age group and sex.

Subjects and methods

The study population comprises the residents of Navarre under 15 years of age. Information on patients diagnosed between 01/01/1975 and 31/12/1991 was obtained as described in the first study conducted in Navarra.¹² Between 01/01/1992 and 31/12/2008; we used the same methods, using Primary Care computerized information systems as main secondary source since 2001. Information on patients diagnosed from 01/01/2009 to 31/12/2012 was obtained as described in our last publication.¹³

Statistical analysis

Incidence rates expressed per 100,000 person-years at risk during the study period, by age group, sex and time period were calculated using data from censuses and records of Navarre (Source: Spanish National Institute of Statistics). Incidence rates were adjusted to a standard population consisting of equal number of children in each of six subgroups defined by age group (0–4, 5–9 and 10–14 years) and sex. Four-year time periods were considered for calculations, except for the first and the last one, which were of five years. Incidence rates were adjusted to a standard population consisting of equal number of children in each of six subgroups defined by age group (0–4, 5–9 and 10–14 years) and sex.

Confidence intervals were estimated at 95%, assuming an underlying Poisson distribution. A Chi-square test was used to compare incidence between groups (independence test), A Poisson

regression model was used to analyze changes in incidence since 1975, adjusted for year of diagnosis, age group and sex, from which rate ratios were obtained, together with their 95% confidence intervals. Interaction terms were also included to assess whether time trend differed among age groups or sex, and removed if not significant. For statistical analysis, IBM SPSS Statistics 20 and R 2.13.1 were used.

This study has been reviewed and approved by the Navarre Research Ethics Committee.

Results

A total of 494 new cases of T1D aged under 15 (273 boys and 221 girls) were recorded, equivalent to a crude incidence rate of 13.4 per 100,000 inhabitants per year (95% CI: 12.2–14.6) and an adjusted incidence rate of 13.2 (95% CI: 12.0–14.3) (Table 1). The age group with the highest incidence was the 10–14 year-olds ($p < 0.001$). The incidence in boys, 14.3 (95% CI: 12.7–16.1), was higher than in girls, 12.3 (95% CI: 10.7–14.0), but no significant differences ($p = 0.094$) were observed except the trend in the 10–14 age group, for which the statistical significance was marginal ($p = 0.051$) (Table 1).

Adjusted incidences for four or five-year periods are shown in Table 2. The highest incidence was recorded in 2004–2007 (21.1; 95% CI: 16.2–26.1) and the lowest, between 1975 and 1979 (4.6; 95% CI: 2.9–6.2) (Table 2). The overall rate of diagnosis increases markedly up to the period 2000–2003, before stabilizing between 20 and 21 cases per 100,000 until the end of the study period.

The results of the Poisson regression model show an annual relative increase of 3.7% (95% CI: 2.9–4.5%) in the incidence rate ($p < 0.001$). Compared to the group under 5 years, the incidence rate in children aged 5–9 is 1.53 (95% CI: 1.18–1.99), and that of children aged 10–14 is 2.50 (95% CI: 1.98–3.18). The rate ratio for boys versus girls is 1.16 (95% CI: 0.97–1.39). We did not find the time trend to be different between age groups (interaction term p -value = 0.912) or sex (interaction term p -value = 0.745).

Table 2
Crude and adjusted incidence rate of type 1 diabetes (number of cases per 100,000 person-years) by time period. Navarre 1975–2012.

| Period | Cases | People/year | Crude incidence | | Adjusted incidence | |
|-----------|-------|-------------|-----------------|-----------|--------------------|-----------|
| | | | Rate | 95% CI | Rate | 95% CI |
| 1975–1979 | 29 | 630,456 | 4.6 | 2.9–6.3 | 4.6 | 2.9–6.2 |
| 1980–1983 | 48 | 483,523 | 9.9 | 7.1–12.7 | 9.8 | 7.0–12.6 |
| 1984–1987 | 45 | 441,642 | 10.1 | 7.2–13.2 | 9.9 | 7.0–12.8 |
| 1988–1991 | 61 | 384,287 | 15.8 | 11.9–19.9 | 14.4 | 10.8–17.9 |
| 1992–1995 | 48 | 333,886 | 14.3 | 10.3–18.4 | 14.0 | 10.0–18.0 |
| 1996–1999 | 36 | 304,741 | 11.8 | 7.9–15.7 | 11.4 | 7.7–15.2 |
| 2000–2003 | 63 | 305,547 | 20.6 | 15.5–25.7 | 20.8 | 15.6–25.9 |
| 2004–2007 | 70 | 334,443 | 20.9 | 16.0–25.8 | 21.1 | 16.2–26.1 |
| 2008–2012 | 94 | 466,021 | 20.1 | 16.1–24.2 | 20.7 | 16.5–24.9 |

Note: 1975–1979 and 2008–2012 span five years while the remaining periods span four years.

Download English Version:

<https://daneshyari.com/en/article/1073251>

Download Persian Version:

<https://daneshyari.com/article/1073251>

[Daneshyari.com](https://daneshyari.com)