

Original

Heavy drinking and alcohol-related injuries in college students

Lucía Moure-Rodríguez^{a,b,*}, Francisco Caamaño-Isorna^a, Sonia Doallo^c, Pablo Juan-Salvadores^a,
Montserrat Corral^c, Socorro Rodríguez-Holguín^c, Fernando Cadaveira^c^a Department of Preventive Medicine, University of Santiago de Compostela, Spain, (CIBERESP)^b Department of Preventive Medicine, Complejo Hospitalario Universitario de Ourense, Spain^c Department of Preventive Medicine, University of Santiago de Compostela, Spain

ARTICLE INFO

Article history:

Received 3 December 2013

Received in revised form 9 February 2014

Accepted 10 February 2014

Available online 13 April 2014

Keywords:

Alcohol Drinking

Epidemiology

Injuries

Adolescents

Cohort Studies

Primary Prevention

ABSTRACT

Objective: The main objective of this study is to evaluate the effect of heavy drinking on alcohol-related injuries.**Material and methods:** We carried out an open cohort study among university students in Spain (n = 1,382). Heavy drinking and alcohol-related injuries were measured by administering AUDIT questionnaires to every participant at the ages of 18, 20, 22 and 24. For data analysis we used a Multilevel Logistic Regression for repeated measures adjusting for consumption of alcohol and cannabis.**Results:** The response rate at the beginning of the study was 99.6% (1,369 students). The incidence rate of alcohol-related injuries was 3.2 per 100 students year. After adjusting for alcohol consumption and cannabis use, the multivariate model revealed that a high frequency of heavy drinking was a risk factor for alcohol-related injuries (Odds Ratio = 3.89 [95%CI: 2.16 – 6.99]). The proportion of alcohol-related injuries in exposed subjects attributable to heavy drinking was 59.78% [95%CI: 32.75 – 75.94] while the population attributable fraction was 45.48% [95%CI: 24.91 – 57.77].**Conclusion:** We can conclude that heavy drinking leads to an increase of alcohol-related injuries. This shows a new dimension on the consequences of this public concern already related with a variety of health and social problems. Furthermore, our results allow us to suggest that about half of alcohol-related injuries could be avoided by removing this consumption pattern.

© 2013 SESPAS. Published by Elsevier España, S.L.U. All rights reserved.

Consumo intensivo y lesiones relacionadas con el alcohol en estudiantes universitarios

RESUMEN

Objetivo: El objetivo principal del estudio es evaluar el efecto del consumo intensivo de alcohol sobre las lesiones relacionadas con esta droga.**Material y métodos:** Se ha realizado un estudio de cohorte abierta entre universitarios en España (n = 1.382). El consumo intensivo y las lesiones relacionadas con el alcohol se midieron mediante la administración del cuestionario AUDIT a cada uno de los participantes a las edades de 18, 20, 22 y 24 años. Para analizar los datos se utilizó la Regresión Logística Multinivel para medidas repetidas ajustando por consumo de alcohol y de cannabis.**Resultados:** La tasa de respuesta al comienzo del estudio fue 99,6% (1.369 estudiantes). La tasa de incidencia de lesiones relacionadas con el alcohol fue de 3,2 por 100 estudiantes año⁻¹. Tras ajustar por consumo de alcohol y de cannabis, el modelo multivariante revela que la alta frecuencia de consumo intensivo fue un factor de riesgo para las lesiones relacionadas con el alcohol (Odds Ratio = 3,89[95%CI:2,16 – 6,99]). La proporción de lesiones relacionadas con el alcohol en expuestos atribuible al consumo intensivo fue 59,78% [95%CI: 32,75 – 75,94] mientras que la fracción atribuible poblacional fue 45,48% [95%CI: 24,91 – 57,77].**Conclusión:** Podemos concluir que el consumo intensivo conduce a un aumento de las lesiones relacionadas con el alcohol. Esto muestra una nueva dimensión de las consecuencias de esta preocupación social que ya se ha relacionado con variedad de problemas sociales y de salud. Además los resultados nos permiten sugerir que aproximadamente la mitad de las lesiones relacionadas con el alcohol podrían evitarse eliminando este patrón de consumo.

© 2013 SESPAS. Publicado por Elsevier España, S.L.U. Todos los derechos reservados.

Palabras clave:

Consumo de Bebidas Alcohólicas

Epidemiología

Lesiones

Adolescentes

Estudios de cohortes

Prevención Primaria

Introduction

In Spain, alcohol consumption is part of the culture. The traditional pattern consists of regular consumption associated with meals, family celebrations, and social events.¹⁻² However, this drinking pattern is increasingly being replaced among young

* Corresponding author.

E-mail address: lucia.moure@rai.usc.es (L. Moure-Rodríguez).

people by intermittent weekend consumption, with episodes of heavy drinking.³ Heavy drinking is characterized by the consumption of large amounts of alcohol in a short time, and brings blood alcohol concentrations to .08 g/dl or above. In Spain, heavy drinking is generally practiced outdoors, at public spaces where large numbers of youths meet. Noise, littering, and even destruction of public property are the main negative consequences attributed by society to this drinking pattern.⁴ Hospital admissions due to alcohol poisoning are other consequences, often out of public sight. However, in Spain the perceived risk of heavy drinking is quite low compared to the perceived risk of the consumption of other drugs, even in small doses.³

Chronic alcohol abuse has well-established effects on morbidity and mortality.^{5,6,7} However, heavy drinking may also increase the risk of alcohol-related health issues.⁸ A recent transversal study has estimated that alcohol-related injuries are more frequent among university students who indulge in a heavy drinking pattern of consumption than in those who do not participate in this pattern of drinking (43% compared with 10% occurrence of alcohol-related injury).⁹ Even though there are important differences between populations, we think this association can also be apply in our context.

To date, no follow-up studies have examined the association between heavy drinking and alcohol-related injuries adjusted for consumption of alcohol in Mediterranean countries.¹⁰ Therefore the aim of this study is to evaluate the effect of heavy drinking on the incidence of alcohol-related injuries.

Material and methods

Design, Population and Sample

An open cohort analysis was conducted within the framework of a cohort study designed to evaluate the neuropsychological and psychophysiological consequences of alcohol use. The study was carried out between November 2005 and May 2012 among students at the University of Santiago de Compostela (Spain). We performed a cluster sampling. From each one of the 33 university schools, at least one of the freshman year classes was randomly selected (a total of 53 classes). The number of classes selected on each university school was proportional to its number of students. All students present in the class on the day of the survey were invited to participate in the study ($n = 1,382$). Abstinent subjects were excluded from the study. The study was approved by the university ethics committee (October 2004).

Data Collection Procedures

Participants were evaluated via a self-administered questionnaire in the classroom in November 2005 and again in November 2007. Students that provided their phone number were further evaluated by phone at 4.5- and 6.5-year follow-up. On all four occasions, alcohol consumption and alcohol-related injuries were measured using the Galician validated version of the Alcohol Use Disorder Identification Test (AUDIT).^{11,12} We decided to use the AUDIT because it is widely considered one of the best screening tests for alcohol abuse; it's transnational; it has been use with college population and it has also been use specifically for measuring heavy drinking. At baseline and at the 2-year follow-up, participants responded to additional questions about age of onset of alcohol consumption and about cannabis consumption.

Definition of variables

Dependent variable

Alcohol-related injuries at 20, 22 and 24 years old. Dichotomous variable. Question 9 of the AUDIT: "Have you or someone else been injured as a result of your drinking? No; Yes, but not during the last year; and Yes, during the last year".

Independent variables

Heavy drinking at 18, 20, and 22 years old. Question 3 of the AUDIT: "How often do you have 6 or more alcoholic drinks on a single occasion? Never; Less than once a month; At least once a month; At least once a week; Daily or Almost daily". The categories At least once a month; At least once a week and Daily or Almost daily were recategorized to More frequently.

Cannabis consumption and number of alcoholic drinks on a typical day (Question 2 of the AUDIT) were also considered as independent variables, because it is known that both variables can result in a lower risk perception and decreased attention, which may consequently lead to injury. Adjusting by both variables, we can therefore identify the specific effect associated to heavy drinking. Frequency of alcohol consumption (Question 1 of the AUDIT) was also considered in order to remove the abstinent subjects.

Cannabis consumption at 18, and 20 years old. This variable was measured with the question "Do you consume cannabis when you go out? Never; Sometimes; Most of the times; Always". The categories Most of the times and Always were recategorized to Usually.

Number of alcoholic drinks on a typical day at 18, 20, and 22 years old. Question 2 of the AUDIT: "How many alcoholic drinks do you have on a typical day when you are drinking? 1 or 2; 3 or 4; 5 or 6; 7 to 9; 10 or more".

Frequency of alcohol consumption at 18, 20, and 22 years old. Question 1 of the AUDIT: "How often do you have a drink containing alcohol? Never; Monthly or less; Two to four times a month; Two to three times a week; Four or more times a week".

Statistical analysis

The follow-up was structured in 3 periods: 11/2005- 11/2007 (2 years); 11/2007 – 05/2010 (2.5 years); and 05/2010 – 05/2012 (2 years). Because of the open cohort design of the study, the conditions of subjects may have changed during follow-up. While the 9th AUDIT question refer to alcohol related injuries suffered in the past, the independent variables about alcohol consumption and cannabis consumption were referred about the present. Therefore, the variable "alcohol-related injuries" measured in 11/2007, 05/2010, and 05/2012 was considered as the effect of both the number of alcoholic drinks on a typical day (Question 2 of the AUDIT) and the heavy drinking (Question 3 of the AUDIT) having occurred in 11/2005, 11/2007 and 05/2010 respectively. Since we have only two measures of cannabis use (2005 and 2007), we considered the alcohol-related injuries in 11/2007 as the effect of the cannabis use in 11-2005 and both alcohol related injuries at 05/2010 and 05/2012 as the effect of cannabis use at 11/2007. At each stage of the study the subjects than answered Never to the first question of the AUDIT were excluded.

We used multilevel logistic regression for repeated measures to obtain adjusted Odds Ratios for Alcohol-related injuries. The university school and the class were considered as randomized variables. The follow-up time was included as an offset term. Data were analyzed using Generalized Linear Mixed Models from the SPSS 20.0.

Download English Version:

<https://daneshyari.com/en/article/1073499>

Download Persian Version:

<https://daneshyari.com/article/1073499>

[Daneshyari.com](https://daneshyari.com)