

Short report

Sex differences in cognitive ageing: Testing predictions derived from life-history theory in a dioecious nematode

Martyna K. Zwoinska^{a,*}, Niclas Kolm^b, Alexei A. Maklakov^a^a Department of Animal Ecology, Evolutionary Biology Centre, Uppsala University, Norbyvägen 18 D, 752 36 Uppsala, Sweden^b Department of Zoology/Ethology, Stockholm University, Svante Arrhenius väg 18 B, 106 91 Stockholm, Sweden

ARTICLE INFO

Article history:

Received 20 June 2013

Received in revised form 5 September 2013

Accepted 30 September 2013

Available online 9 October 2013

Section Editor: T.E. Johnson

Keywords:

Ageing

Caenorhabditis

Learning

Life-history

Sex differences

Trade-off

ABSTRACT

Life-history theory maintains that organisms allocate limited resources to different traits to maximize fitness. Learning ability and memory are costly and known to trade-off with longevity in invertebrates. However, since the relationship between longevity and fitness often differs between the sexes, it is likely that sexes will differentially resolve the trade-off between learning and longevity. We used an established associative learning paradigm in the dioecious nematode *Caenorhabditis remanei*, which is sexually dimorphic for lifespan, to study age-related learning ability in males and females. In particular, we tested the hypothesis that females (the shorter-lived sex) show higher learning ability than males early in life but senesce faster. Indeed, young females outperformed young males in learning a novel association between an odour (butanone) and food (bacteria). However, while learning ability and offspring production declined rapidly with age in females, males maintained high levels of these traits until mid-age. These results not only demonstrate sexual dimorphism in age-related learning ability but also suggest that it conforms to predictions derived from the life-history theory.

© 2013 Elsevier Inc. All rights reserved.

1. Introduction

Learning ability is a key behavioural trait that affects survival and reproduction of both sexes but there is mounting evidence that learning and memory are costly in vertebrates and invertebrates alike (Dukas, 1998; Kotschal et al., 2013; Mery and Kawecki, 2003; Snell-Rood et al., 2011). Specifically, learning ability has been shown to trade-off with longevity (Burger et al., 2008; Lagasse et al., 2012; Mery and Kawecki, 2004; Plaçais and Preat, 2013) and reproduction (Snell-Rood et al., 2011). The most striking example of a cost of learning is a symmetrical evolutionary trade-off between learning ability and longevity in the fruit fly *Drosophila melanogaster* (Burger et al., 2008). It was found that experimental populations selected for improved learning suffered a considerable reduction in longevity, while populations selected for increased longevity suffered a massive reduction in learning ability (Burger et al., 2008). Furthermore, learning and memory were also shown to incur constitutive fitness costs in terms of delayed reproduction and lower fecundity in butterflies (Snell-Rood et al., 2011), and reduced larval competitive ability in *D. melanogaster* (Mery and Kawecki, 2003).

Because males and females often resolve their survival vs. reproduction trade-offs differently (Bonduriansky et al., 2008; Maklakov and

Lummaa, 2013; Trivers, 1972; Williams, 1957), it is likely that the trade-off between survival and learning will also be resolved differently between the sexes. Applying the logic of a symmetrical evolutionary trade-off (Burger et al., 2008), increased sex-specific investment in learning ability should result in reduced longevity in a given sex. However, the longer-lived sex would experience stronger selection late in life because of a larger number of surviving individuals in older ages (Williams, 1957) such that it should maintain its level of performance for a longer period of time and senesce slower than the shorter-lived sex. This basic pattern can be modified when fitness costs and benefits of learning differ between the sexes.

Despite the evidence that learning ability is likely to trade-off with longevity and the widespread occurrence of sexual dimorphism in lifespan across taxa (Clutton-Brock and Isvaran, 2007), little is known about sex differences in age-related learning across species and whether such differences are in line with predictions derived from life-history theory. This is unfortunate because there is broad cross-disciplinary interest in cognitive differences between the sexes, as well as in sex-specific ageing. Animal models that are accessible for experimental manipulation may prove particularly valuable in linking sexual dimorphism in major life-history traits to sex differences in life-long learning performance.

Here, we used a positive associative learning paradigm in the dioecious nematode worm *Caenorhabditis remanei*, which exhibits sexual dimorphism in longevity (McCulloch and Gems, 2003) and reproductive ageing, to test whether sexes differ in age-related learning performance. Specifically, we tested whether the shorter-lived sex (females in *C. remanei*) exhibits improved learning ability early in life, and whether

* Corresponding author at: Ageing Research Group, Department of Animal Ecology, Evolutionary Biology Centre, Uppsala University, Norbyvägen 18 D, 752 36 Uppsala, Sweden. Tel.: +46 18 471 2613.

E-mail addresses: martyna.zwoinska@ebc.uu.se (M.K. Zwoinska), niclas.kolm@ebc.uu.se (N. Kolm), alexei.maklakov@ebc.uu.se (A.A. Maklakov).

the longer-lived sex (males) shows slower ageing in learning. *Caenorhabditis* nematodes can learn in associative and non-associative paradigms, and form short- and long-term memories. Since learning and memory are known to be actively regulated by highly-conserved longevity pathways in these animals, they emerge as a convenient system for the study of trade-offs between cognitive and life-history traits (Kauffman et al., 2010; Stein and Murphy, 2012).

2. Materials and methods

2.1. Worm strain and culture

We used an outbred SP8 strain of *C. remanei*, provided by N. Timmermeyer from the Department of Biology at Tuebingen University, Germany. This strain harbours substantial amounts of genetic variation for fitness and life-history traits (Chen and Maklakov, 2012). Worm cultures were recovered from freezing and maintained under standard cultivation conditions (Stiernagle, 2006) for 2–3 generations before being used in experiments. Age-synchronized populations were obtained by isolating eggs through a hypochlorite treatment (Stiernagle, 2006). From the first day of adulthood onwards, animals were transferred to new plates every day to prevent mixing adults with progeny.

2.2. Longevity and reproduction assays

For the longevity assay, virgin worms (L4 larvae stage) of the focal sex were placed together with the corresponding number of virgin worms of the opposite sex (10 worms of the focal sex per plate, $n = 4$ plates for each sex). The sex ratio was adjusted throughout the study period. In the female reproductive assay ($n = 13$ females), a virgin

female was placed with a single male for about 24 h. Each day a female was transferred to a new plate with a fresh male. The male reproductive assay ($n = 12$ males) was conducted by allowing a single male to mate with 8 virgin females for 3 h. After this time the male was removed and females were allowed to lay eggs for another 3 h. Two days later the progeny, by this time mainly in the L4 stage, were counted.

2.3. Behavioural assays

Behavioural assays were conducted at days 2, 5 or 8 of adulthood, which roughly corresponds to young, middle-aged and elderly adults in this population (Chen and Maklakov, 2012), and followed the protocol from Kauffman et al. (2010) (for details see Electronic Supplementary Material). Prior to testing, males and females were separated and left overnight on Petri plates with nematode growth medium (NGM) and bacterial food source (*Escherichia coli* OP50). Three behavioural responses were quantified: motility, naïve chemotaxis to butanone, and post-conditioning chemotaxis to butanone. For all assays, animals were placed on 92 mm NGM plates (no food) on the place of origin and allowed to move freely for 1 h. After this time, assays were terminated by inverting the plate over a few drops of chloroform and then worms were hand-counted. The number of worms per assay ranged from 50 to 151. Motility was quantified by calculating the motility index (MI) according to the following formula: $MI = \frac{n_{\text{left the place of origin}}}{(n_{\text{total on a plate}} - n_{\text{dead on the place of origin}})}$ (we subtracted all worms that have been accidentally killed or injured during testing, this was not, however, a big number), where n represents the number of worms. Naïve chemotaxis to butanone was estimated by placing naïve worms on the place of origin at equal distance from 10% butanone and ethanol (control) odour patches at the opposite sides of the plate (Fig. S1). 1.5 μ l




Fig. 1. (a) Survival and (b) reproduction of males and females *C. remanei* from the first day of adulthood onwards. For illustrative purpose offspring production is presented as the mean of the number of larvae per individual for a given day divided by the highest mean of larvae produced per day; \pm SEM. Solid lines, circles – females and dotted lines, triangles – males.

Download English Version:

<https://daneshyari.com/en/article/10736855>

Download Persian Version:

<https://daneshyari.com/article/10736855>

[Daneshyari.com](https://daneshyari.com)