



Contents lists available at ScienceDirect

## Biochemical and Biophysical Research Communications

journal homepage: [www.elsevier.com/locate/ybbrc](http://www.elsevier.com/locate/ybbrc)

## Hepatic maturation of human iPS cell-derived hepatocyte-like cells by ATF5, c/EBP $\alpha$ , and PROX1 transduction



Daiki Nakamori <sup>a, 1</sup>, Kazuo Takayama <sup>a, b, c, 1</sup>, Yasuhito Nagamoto <sup>a, c</sup>, Seiji Mitani <sup>a</sup>, Fuminori Sakurai <sup>a, d</sup>, Masashi Tachibana <sup>a</sup>, Hiroyuki Mizuguchi <sup>a, c, e, f, \*</sup>

<sup>a</sup> Laboratory of Biochemistry and Molecular Biology, Graduate School of Pharmaceutical Sciences, Osaka University, Osaka 565-0871, Japan

<sup>b</sup> The Keihanshin Consortium for Fostering the Next Generation of Global Leaders in Research (K-CONNEX), Kyoto University, Kyoto 606-8302, Japan

<sup>c</sup> Laboratory of Hepatocyte Regulation, National Institute of Biomedical Innovation, Health and Nutrition, Osaka 567-0085, Japan

<sup>d</sup> Laboratory of Regulatory Sciences for Oligonucleotide Therapeutics, Clinical Drug Development Project, Graduate School of Pharmaceutical Sciences, Osaka University, Osaka 565-0871, Japan

<sup>e</sup> iPS Cell-based Research Project on Hepatic Toxicity and Metabolism, Graduate School of Pharmaceutical Sciences, Osaka University, Osaka 565-0871, Japan

<sup>f</sup> Global Center for Medical Engineering and Informatics, Osaka University, Osaka 565-0871, Japan

### ARTICLE INFO

#### Article history:

Received 25 November 2015

Accepted 1 December 2015

Available online 9 December 2015

#### Keywords:

Human induced pluripotent stem cells

Hepatocyte

Drug metabolism

ATF5

c/EBP $\alpha$

PROX1

### ABSTRACT

Hepatocyte-like cells differentiated from human iPS cells (human iPS-HLCs) are expected to be utilized in drug development and research. However, recent hepatic characterization of human iPS-HLCs showed that these cells resemble fetal hepatocytes rather than adult hepatocytes. Therefore, in this study, we aimed to develop a method to enhance the hepatic function of human iPS-HLCs. Because the gene expression levels of the hepatic transcription factors (activating transcription factor 5 (ATF5), CCAAT/enhancer-binding protein alpha (c/EBP $\alpha$ ), and prospero homeobox protein 1 (PROX1)) in adult liver were significantly higher than those in human iPS-HLCs and fetal liver, we expected that the hepatic functions of human iPS-HLCs could be enhanced by adenovirus (Ad) vector-mediated ATF5, c/EBP $\alpha$ , and PROX1 transduction. The gene expression levels of *cytochrome P450 (CYP) 2C9, 2E1, alpha-1 antitrypsin, transthyretin, Na<sup>+</sup>/taurocholate cotransporting polypeptide*, and *uridine diphosphate glucuronosyl transferase 1A1* and protein expression levels of CYP2C9 and CYP2E1 were upregulated by ATF5, c/EBP $\alpha$ , and PROX1 transduction. These results suggest that the hepatic functions of the human iPS-HLCs could be enhanced by ATF5, c/EBP $\alpha$ , and PROX1 transduction. Our findings would be useful for the hepatic maturation of human iPS-HLCs.

© 2015 The Authors. Published by Elsevier Inc. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

**Abbreviations:** Human iPS-HLCs, Human induced pluripotent stem cells-derived hepatocyte-like cells; HLCs, Hepatocyte-like cells; ATF5, Activating transcription factor 5; c/EBP $\alpha$ , CCAAT/enhancer-binding protein alpha; PROX1, Prospero homeobox protein 1; 3 TFs, The three transcription factors (ATF5, c/EBP $\alpha$ , and PROX1); Ad, Adenovirus; CYP, cytochrome P450;  $\alpha$ AT, alpha-1 antitrypsin; TTR, transthyretin; NTCP, Na<sup>+</sup>/taurocholate cotransporting polypeptide; UGT1A1, Uridine diphosphate glucuronosyltransferase 1A1; HNF, Hepatocyte nuclear factor; FGF, Fibroblast growth factor; FBS, Fetal bovine serum; BMP, Bone morphogenetic protein; HCM, Hepatocyte culture medium; EGF, Epidermal growth factor; HGF, Hepatocyte growth factor; OsM, Oncostatin M; BSA, Bovine serum albumin; FOXA2, Forkhead box protein A2; AhR, Aryl hydrocarbon receptor; CAR, Constitutive androstane receptor; PPAR $\alpha$ , Peroxisome proliferator-activated receptor alpha; LXR, Liver X receptor.

\* Corresponding author. Laboratory of Biochemistry and Molecular Biology, Graduate School of Pharmaceutical Sciences, Osaka University, 1-6 Yamadaoka, Suita, Osaka 565-0871, Japan.

E-mail address: [mizuguch@phs.osaka-u.ac.jp](mailto:mizuguch@phs.osaka-u.ac.jp) (H. Mizuguchi).

<sup>1</sup> These two authors contribute equally to the work.

### 1. Introduction

The liver is an important organ in term of the absorption, distribution, metabolism, and excretion of drugs. Therefore, to develop safe and effective drugs, primary human hepatocytes (PHHs) are widely used in the early stage of drug development. However, alternative hepatocyte resources for drug screening are required because PHHs are costly and not abundantly available. Hepatocyte-like cells (HLCs) that are generated from human induced pluripotent stem (iPS) cells are expected to be used in drug screening in place of PHHs. We previously reported that the stage-specific transient transduction of hepatic transcription factors mediated by adenovirus (Ad) vector was useful for promoting hepatic differentiation [1–4]. However, the drug metabolism capacity of human iPS-HLCs is still lower than that of PHHs [4]. In addition, Baxter

et al. have shown by means of phenotypic and functional analyses that the human iPS-HLCs better mimic fetal hepatocytes rather than adult hepatocytes [5]. Therefore, it is necessary to promote the hepatic maturation of human iPS-HLCs.

Recently, Du et al. showed that metabolically functioning human induced hepatocytes (hiHeps) can be converted from fibroblasts by overexpressing the hepatic fate conversion factors hepatocyte nuclear factor 1 alpha, 4 alpha, and 6 (HNF1 $\alpha$ , HNF4 $\alpha$ , HNF6, respectively) [6]. In addition, they demonstrated that hiHeps could be further matured by overexpressing activating transcription factor 5 (ATF5), CCAAT/enhancer-binding protein alpha (c/EBP $\alpha$ ), and prospero homeobox protein 1 (PROX1). Consistently, all the hepatic transcription factors (ATF5, c/EBP $\alpha$ , and PROX1) are known to play an important role in the maturation of hepatocytes and maintenance of hepatic function. Specifically, it is known that the level of cytochrome P450 (CYP) 2B6 expression in hepatocytes is regulated by ATF5, and also cooperatively regulated with the hepatic nuclear receptor, constitutive androstane receptor (CAR) [7]. The hepatic functions, such as albumin secretion and urea production, are positively regulated by c/EBP $\alpha$  [8]. It has also been reported that PROX1 plays a key role in the metabolic maturation of hepatocytes, as well as FOXA2 [9]. Considering these results together, we expected that the hepatic functions of human iPS-HLCs could be improved by the transduction of three transcription factors, ATF5, c/EBP $\alpha$ , and PROX1.

In this study, we attempt to produce matured human iPS-HLCs by Ad vector-mediated ATF5, c/EBP $\alpha$ , and PROX1 (3TFs) transduction. To investigate whether the hepatic functions were enhanced, we analyzed the expression of hepatocyte markers in the Ad-3TFs-transduced human iPS-HLCs. Moreover, the protein expression levels of CYP2C9, CYP2E1, and  $\alpha$ -1 antitrypsin ( $\alpha$ AT) were examined by western blotting and immunofluorescent staining.

## 2. Materials and methods

### 2.1. Human iPS cells

Human iPS cells generated from the human embryonic lung fibroblast cell line MCR5 were provided from the JCRB Cell Bank (Dotcom, JCRB Number: JCRB1327). Human iPS cells were maintained on a feeder layer of mitomycin C-treated mouse embryonic fibroblasts (Millipore) with ReproStem (ReproCELL) supplemented with 10 ng/ml fibroblast growth factor (FGF) 2 (KATAYAMA Kogyo Kagaku).

### 2.2. Hepatocyte differentiation

Before the initiation of hepatocyte differentiation, human iPS cells were dissociated into clumps by using dispase (Roche Diagnostics) and plated onto BD Matrigel Basement Membrane Matrix Growth Factor Reduced (Becton, Dickinson and Company). These cells were cultured in the mouse embryo fibroblasts-conditioned medium for 3–4 days. The differentiation protocol for the induction of definitive endoderm cells, hepatoblast-like cells, and human iPS-HLCs was based on our previous reports with some modifications [10]. Briefly, in the definitive endoderm differentiation, human iPS cells were cultured with the L-Wnt3A-expressing cell (CRL2647; ATCC)-conditioned RPMI1640 medium (Sigma) containing 100 ng/mL Activin A (R&D Systems), 1% GlutaMAX (Invitrogen), 0.2% fetal bovine serum (FBS), and 1  $\times$  B27 Supplement Minus Vitamin A (Invitrogen) for 4 days. For the induction of hepatoblasts, the definitive endoderm cells were cultured with RPMI1640 medium containing 30 ng/mL bone morphogenetic protein 4 (BMP4) (R&D Systems) and 20 ng/mL

FGF4 (R&D Systems), 1% GlutaMAX, and 1  $\times$  B27 Supplement Minus Vitamin A for 5 days. To perform the hepatocyte differentiation, the hepatoblast-like cells were cultured with RPMI1640 medium containing 20 ng/mL hepatocyte growth factor (HGF) (R&D Systems), 1% GlutaMAX, and 1  $\times$  B27 Supplement Minus Vitamin A for 5 days. Finally, the cells were cultured with hepatic maturation medium (hepatic maturation medium consists of Hepatocyte Culture Medium (HCM; Lonza, without epidermal growth factor (EGF)) containing 20 ng/mL oncostatin M (OsM) and 3% GlutaMAX) for 11 days. To promote hepatic maturation, human iPS-HLCs were transduced with 1000 VP/cell of Ad-ATF5, Ad-c/EBP $\alpha$ , and Ad-PROX1 (total 3000 VP/cell) for 1.5 h on day 25 and were cultured with hepatic maturation medium for 5 days.

### 2.3. Ad vector

Ad vectors were constructed by an improved *in vitro* ligation method [11,12]. The human ATF5 and PROX1 gene (accession number NM\_001193646 and NM\_001270616, respectively) were amplified by PCR using primers: ATF5 Fwd 5'-TGctagaCCACCATGTCCTGCGGACCTG -3' and ATF5 Rev 5'-AAGcgccgcCTAGCAGCTACGGGTCTCTG -3'; PROX1 Fwd 5'-AGtcagaCCACCATGCTGACCATGACAGCAC -3' and PROX1 Rev 5'-TAGcgccgcCTACTCATGAAGCAGCTCTG -3'. The human ATF5 and PROX1 gene was inserted into pHMEF5 [13], which contains the human elongation factor-1 $\alpha$  (EF-1 $\alpha$ ) promoter, resulting in pHMEF-ATF5 and -PROX1, respectively. The pHMEF-ATF5 and -PROX1 was digested with I-CeuI/PI-SceI and ligated into I-CeuI/PI-SceI-digested pAdHM41-K7 [14], resulting in pAd-ATF5 and pAd-PROX1, respectively. The human EF-1 $\alpha$  promoter-driven LacZ- and c/EBP $\alpha$ -expressing Ad vectors (Ad-LacZ and Ad-c/EBP $\alpha$ , respectively) were constructed previously [15,16]. Ad-LacZ, Ad-ATF5, Ad-c/EBP $\alpha$ , and Ad-PROX1, each of which contains a stretch of lysine residue (K7) peptides in the C-terminal region of the fiber knob for more efficient transduction of human iPS cells, definitive endoderm cells, and hepatoblast-like cells were used in this study. The vector particle (VP) titer was determined by using a spectrophotometric method [17].

### 2.4. Real-time RT-PCR

Total RNA was isolated from human iPS cells and their derivatives using ISOGENE (NIPPON GENE). cDNA was synthesized using 500 ng of total RNA with a Superscript VILO cDNA synthesis kit (Invitrogen). Real-time RT-PCR was performed with SYBR Green PCR Master Mix (Applied Biosystems) using a StepOnePlus real-time PCR system (Applied Biosystems). Relative quantification was performed against a standard curve and the values were normalized against the input determined for the housekeeping gene, *glyceraldehyde 3-phosphate dehydrogenase (GAPDH)*. The primer sequences used in this study are described in Table S1. Fetal liver and Adult liver total RNA were purchased from Clontech and BioChain, respectively.

### 2.5. Western blotting

The human iPS cell-derivatives were homogenized with RIPA Lysis and Extraction Buffer (Thermo Fisher Scientific) containing a protease inhibitor mixture (Sigma). After being frozen and thawed, the homogenates were centrifuged at 15,000 g at 4  $^{\circ}$ C for 10 min, and the supernatants were collected. The lysates were subjected to SDS-PAGE on 7.5% polyacrylamide gel, and then transferred onto polyvinylidene fluoride membranes (Millipore). After the reaction was blocked with 1% skim milk in TBS containing 0.1% Tween 20 (Sigma) at room temperature for 1 h, the membranes were

Download English Version:

<https://daneshyari.com/en/article/10749637>

Download Persian Version:

<https://daneshyari.com/article/10749637>

[Daneshyari.com](https://daneshyari.com)