

Accepted Manuscript

Hepatic fibrosis and angiogenesis after bile duct ligation are endogenously expressed vasohibin-1 independent

Yutaka Furutani, Yumi Shiozaki-Sato, Mitsuko Hara, Yasufumi Sato, Soichi Kojima



PII: S0006-291X(15)30003-6

DOI: [10.1016/j.bbrc.2015.05.074](https://doi.org/10.1016/j.bbrc.2015.05.074)

Reference: YBBRC 33972

To appear in: *Biochemical and Biophysical Research Communications*

Received Date: 12 May 2015

Accepted Date: 22 May 2015

Please cite this article as: Y. Furutani, Y. Shiozaki-Sato, M. Hara, Y. Sato, S. Kojima, Hepatic fibrosis and angiogenesis after bile duct ligation are endogenously expressed vasohibin-1 independent, *Biochemical and Biophysical Research Communications* (2015), doi: 10.1016/j.bbrc.2015.05.074.

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

Hepatic fibrosis and angiogenesis after bile duct ligation are endogenously expressed vasohibin-1 independent

Yutaka Furutani^a, Yumi Shiozaki-Sato^a, Mitsuko Hara^a, Yasufumi Sato^b, and Soichi Kojima^a

^aMicro-Signaling Regulation Technology Unit, RIKEN Center for Life Science Technologies, Saitama, 2-1 Hirosawa, Wako, Saitama 351-0198, Japan.

^bDepartment of Vascular Biology, Institute of Development, Aging and Cancer, Tohoku University, 4-1 Seiryō-machi, Aoba-ku, Sendai 980-8575, Japan.

*Correspondence should be addressed to S.K. (skojima@riken.jp)

Download English Version:

<https://daneshyari.com/en/article/10751076>

Download Persian Version:

<https://daneshyari.com/article/10751076>

[Daneshyari.com](https://daneshyari.com)