



Pre-clinical evaluation of cinobufotalin as a potential anti-lung cancer agent



Sheng Kai¹, Jia-huan Lu¹, Ping-ping Hui, Hui Zhao*

Department of Geriatrics, Shanghai Changning Center Hospital, Shanghai 200336, China

ARTICLE INFO

Article history:

Received 26 August 2014

Available online 6 September 2014

Keywords:

Lung cancer
Cinobufotalin
Cyclophilin D
Mitochondrial permeability transition pore (mPTP)
Cell death

ABSTRACT

Lung cancer is a major cause of cancer-related mortality in the United States and around the world. Due to the pre-existing or acquired chemo-resistance, the current standard chemotherapy regimens only show moderate activity against lung cancer. In the current study, we explored the potential anti-lung cancer activity of cinobufotalin *in vivo* and *in vitro*, and studied the underlying mechanisms. We demonstrated that cinobufotalin displayed considerable cytotoxicity against lung cancer cells (A549, H460 and HTB-58 lines) without inducing significant cell apoptosis. Our data suggest that mitochondrial protein cyclophilin D (Cyp-D)-dependent mitochondrial permeability transition pore (mPTP) opening mediates cinobufotalin-induced non-apoptotic death of lung cancer cells. The Cyp-D inhibitor cyclosporine A (CsA), the mPTP blocker sanglifehrin A (SfA), and Cyp-D shRNA-silencing significantly inhibited cinobufotalin-induced mitochondrial membrane potential (MMP) reduction and A549 cell death (but not apoptosis). Using a mice xenograft model, we found that cinobufotalin inhibited A549 lung cancer cell growth *in vivo*. Thus, cinobufotalin mainly induces Cyp-D-dependent non-apoptotic death in cultured lung cancer cells. The results of this study suggest that cinobufotalin might be further investigated as a novel anti-lung cancer agent.

© 2014 Elsevier Inc. All rights reserved.

1. Introduction

Lung cancer, mainly non-small cell lung cancers (NSCLCs), is a major cause of cancer-related mortality in the United States and around the world [1–4]. Surgery resection is available only for the early stage of lung cancers [2,5]. Majority of NSCLC or other lung cancer patients are presenting with advanced/metastatic malignancies at time of diagnosis, and chemotherapy are important treatment options [5]. Due to the pre-existing or acquired chemo-resistance, the current standard chemotherapy regimen only showed moderate activity against lung cancers. Thus, the search more efficient agents are extremely important and urgent.

Traditional Chinese medicine has become an important source for novel chemotherapeutic compounds [6,7]. Cinobufotalin, a cardiotonic steroids or bufadienolides [8,9], is extracted from the skin secretions of the traditional Chinese medicine giant toads (*Chan Su*) [8]. Cinobufotalin has been used as a cardiotonic, diuretic and a hemostatic agent [9]. Although numerous reports have displayed

anti-tumor activity and chemotherapeutic enhancing properties of many bufadienolides (i.e. bufalin) in a wide spectrum of cancer cells [10], the possible anti-tumor ability of cinobufotalin has not been extensively studied [8,9]. Its potential role in lung cancer cells is un-known. More importantly, the underlying mechanisms remain to be explored.

Mitochondrial permeability transition pore (mPTP) is a key regulator of cell death [11,12]. It is a channel complex composed of voltage-dependent anion channel (VDAC, in the outer membrane), adenine nucleotide translocator (ANT, in the inner membrane), and cyclophilin D (Cyp-D, inside the mitochondrial matrix) [11,13]. With cytotoxic stimuli, Cyp-D will associate with ANT in the inner membrane to promote mPTP opening, which leads to mitochondrial depolarization, mitochondria swelling, Ca²⁺ release, and eventually cell death [14,15]. In this study, we found that Cyp-D and mPTP are also important mediators of cinobufotalin-induced lung cancer cell death.

2. Materials and methods

2.1. Chemicals and reagents

Cinobufotalin, sanglifehrin A (SfA) and cyclosporine A (CsA) were obtained from Sigma Chemicals (St. Louis, MO); Z-VAD-fmk

* Corresponding author. Address: Department of Geriatrics, Shanghai Changning Center Hospital, No. 1111, Xian-xia Road, Changning District, Shanghai 200336, China. Fax: +86 21 62909911 2407.

E-mail address: huizhaovip@126.com (H. Zhao).

¹ Equal contributors.

(ZVAD) was purchased from Calbiochem (Shanghai, China). Antibodies against Cyp-D and tubulin were purchased from Santa Cruz Biotech (Santa Cruz, CA).

2.2. Cell culture

A549, H460 and HTB-58 human lung cancer cell lines were purchased from Cell Bank of Chinese Academy of Medical Sciences (Shanghai, China). Cells were cultured in DMEM medium supplemented with 10% fetal bovine serum (FBS, Hyclone, Shanghai, China) with antibiotics, and incubated at 37 °C in a humidified air atmosphere containing 5% CO₂.

2.3. Cell survival assay

Cell survival was evaluated by the tetrazolium dye (MTT, Sigma) assay as reported [16]. Cells were seeded into 96-well plates and then treated. MTT (5 mg/ml in PBS) was added into the wells and incubated for 3 h. After the medium was carefully aspirated, DMSO (150 µl, Sigma) was added to each well. The plates were gently agitated until the color reaction was uniform and the OD₄₉₀ was determined using a microplate reader. Media-only treated cells served as 100% cell viability, and the relative survival was defined as absorbance of treated wells divided by that of controls. For each treatment, 5-wells were included.

2.4. Cell death assay

The number of dead cancer cells (Trypan blue positive) after treatment was counted, and the percentage (%) of dead cells was calculated by the number of the Trypan blue positive cells divided by the total number of the cells, which was automatically recorded by a cell counter (Roche, Shanghai, China).

2.5. Cell apoptosis assay

Cells were seeded into 6-well plates, and treatments were initiated when cells were 60–70% confluent. Tested reagents were added into the wells and incubated for indicated time points, and cells were then harvested by digesting with trypsin/EDTA (Gibco, NY). After fixed with cold 70% ethanol, cells were stained with propidium iodide (PI) solution (5 mg/mL, Invitrogen, CA) and Annexin V (1 mg/mL, Invitrogen, CA) for 30 min in 37 °C. Samples (10,000 cells/sample) were tested on a Becton Dickinson FACS calibur (Becton, Dickinson & Co, Mountain View, CA). Annexin V percentage was used to reflect cell apoptosis. Annexin V positive/PI negative and Annexin V positive/PI positive cells were combined as apoptotic cells, see [16].

2.6. Hoechst 33342 staining of apoptotic cells

Cells were plated at 50,000 per well in 6-well cell culture plates with glass slides (Corning Incorporated, USA) and cultured overnight. After treatment, cells were fixed with 3.7% formaldehyde in PBS for 10 min and stained with Hoechst 33342 solution (10 µg/mL, Sigma). The slides were washed twice in PBS and fixed onto the microscopic slide. The cell nuclei images were taken with a Zeiss fluorescence microscope (Zeiss, Nanjing, China), and the number of apoptotic cells was manually counted in 5 random fields for at least 500 cells from each group. The apoptotic nuclei showed characteristic condensation of the chromatin, see [16].

2.7. Western-blot

Cells were harvested with trypsin/EDTA, and PBS-washed cell pellets were treated with HEPES lysis buffer (30 mM HEPES, 1% Tri-

ton X-100, 10% glycerol, 5 mM MgCl₂, 25 mM NaF, 1 mM EDTA and 10 mM NaCl). Equal amounts of protein extracts were loaded onto sodium dodecyl sulfate (SDS)-polyacrylamide gels and ran at 100 mV for 80 min, followed by transferring to PVDF membranes at 100 mV for 30 min at room temperature. The membranes were probed with indicated primary antibodies at room temperature for 180 min. As secondary antibodies, goat anti-rabbit/mouse labeled with horseradish peroxidase (HRP, Santa Cruz Biotechnology, Santa Cruz, CA) were added. Blots were developed using a chemiluminescence detection system. Blot intensity was quantified through the ImageJ software (NIH).

2.8. Cyp-D stable knockdown

Cyp-D stable knockdown in A549 cells were based on the lentiviral infection procedure as reported by Dr. Bi [16,17]. Cyp-D and equal loading (tubulin) expression in the infected cells was tested by Western blots. For Cyp-D shRNA infection assay, two non-overlapping shRNAs against Cyp-D cDNA were utilized to insure results consistency, and stable clones were selected through puromycin [16,17].

2.9. Mitochondrial membrane potential (MMP) assay

MMP was measured through JC-10 dye (Invitrogen, Carlsbad, CA) [18]. The dye forms aggregates in the mitochondrial membrane, exhibiting orange fluorescence in normally resting cells. When the membrane potential is decreasing, the monomeric JC-10 will form in the cytosol, which exhibits green fluorescence. Thus, the intensity of green fluorescence is detected as indicator of MMP reduction [19]. Briefly, after treatment, A549 cells were stained with 5.0 µg/ml of JC-10 for 15 min in the dark. Cells were then washed twice with warm PBS, and resuspended in fresh culture medium and read immediately on a microplate reader with an excitation filter of 485 nm. The OD value of treatment group was normalized to that of medium-treated control group.

2.10. Animals

Male nude mice (4–6 weeks old, BALB/c) were purchased from the Institute of Laboratory Animal Science, Chinese Academy of Medical Sciences (Shanghai, China). All of the experiments were performed in accordance with the Guidelines for Care and Use of Experimental Animals of Experimental Animal Research Committee of authors' affiliation.

2.11. A549 tumor xenograft model in nude mice

Male nude mice were introduced to establish xenograft tumor model of A549 cells as previously described [16]. A549 cells (five million cells in 0.1 mL of culture medium) were subcutaneously injected at the right thigh of nude mice, and treatment was started when the tumors reached an average volume of 200–300 mm³. Animals were randomized into 3 groups with 10 mice each group: (a) vehicle; (b) 1.0 mg/kg of cinobufotalin; (c) 5.0 mg/kg of cinobufotalin. Cinobufotalin was injected intraperitoneally (i.p.) twice daily for 1 weeks. The mice were examined daily for toxicity/mortality relevant to treatment, and the tumor was measured with a caliper once a week for up to 5 weeks. Mice body weight and mice survival (at week 5) were also recorded. The tumor volume (in mm³) was calculated by the formula: volume = (width)² × length / 2, and the tumor growth curve was presented.

Download English Version:

<https://daneshyari.com/en/article/10753904>

Download Persian Version:

<https://daneshyari.com/article/10753904>

[Daneshyari.com](https://daneshyari.com)