



Contents lists available at ScienceDirect

Biochemical and Biophysical Research Communications

journal homepage: [www.elsevier.com/locate/ybbrc](http://www.elsevier.com/locate/ybbrc)

## RmKK, a tissue kallikrein inhibitor from *Rhipicephalus microplus* eggs

Patrícia A. Abreu, Tatiane S. Soares, Diego S. Buarque, Ricardo S. Torquato, Aparecida S. Tanaka \*

Department of Biochemistry, Universidade Federal de São Paulo, São Paulo, SP, Brazil

### ARTICLE INFO

#### Article history:

Received 21 April 2014

Available online xxxxx

#### Keywords:

*Rhipicephalus microplus*

Eggs

Tissue kallikrein inhibitor

Serine protease inhibitor

### ABSTRACT

*Rhipicephalus microplus* is an important ectoparasite that is responsible for transmission of anaplasmosis and babesiosis to cattle. Tissue kallikrein inhibitors might play an important role in *R. microplus* eggs. In the present work, we purified and characterized, a tissue kallikrein inhibitor presents in *R. microplus* eggs (RmKK), a protein which contains two Kunitz domain in tandem. Purified inhibitor was confirmed by amino terminal determination and its dissociation constant ( $K_i$ ) for bovine trypsin and porcine pancreatic kallikrein were 0.6 nM and 91.5 nM, respectively. Using a cDNA library from *R. microplus* midgut, we cloned the cDNA fragment encoding mature RmKK and expressed the protein in *Pichia pastoris* system. Recombinant RmKK was purified by ion exchange chromatography and presented molecular mass of 16.3 kDa by MALDI-TOF analysis. Moreover, RmKK showed a tight binding inhibition for serine proteases as bovine trypsin ( $K_i = 0.2$  nM) and porcine pancreatic kallikrein (PPK) ( $K_i = 300$  nM). We performed, for the first time, the characterization of a tissue kallikrein inhibitor presents in *R. microplus* eggs, which the transcript is produced in the adult female gut. BmKK seems to be the strongest PPK inhibitor among all BmTIs present in the eggs and larvae (Andreotti et al., 2001; Sasaki et al., 2004). This data suggests that BmKK may participate in the development of tick egg and larvae phase.

© 2014 Elsevier Inc. All rights reserved.

### 1. Introduction

*Rhipicephalus microplus* is one of the most important ectoparasites in the world, inflicting economic losses in cattle production. *R. microplus* infestations cause reductions in cattle's weight and milk production [3]. Moreover, this tick species transmits several pathogens, including anaplasmosis [4] and babesiosis parasites, which result in neurological problems and death of cattle [5].

The study of new important molecules for *R. microplus* can help in the development of methods to control infestation by this tick. Ticks are a rich source of inhibitors of serine proteases, particularly those belonging to the family BPTI – Kunitz [2]. Several inhibitors of the Kunitz type have been studied and examples of these are: TAP (tick anticoagulant peptide) [6] ornithodorin [7], BmTI-A [8], RSTI [9], boophilin, chymotrypsin inhibitor [10] and savignin [11], among others. However, the role of some proteins from the Kunitz family remains to be elucidated and it includes the tissue kallikrein inhibitors from tick's presents in the pool of BmTIs [1,2].

It becomes important to study inhibitors of kallikreins, since these enzymes are involved in the main physiologic regulatory systems of animals, i.e. blood coagulation, and fibrinolysis, blood

fluidity, vascular growth, and inflammatory process and interferes in the renin–angiotensine system [12]. Therefore, kallikreins are very important molecules in the homeostasis of the animals among them mammals including cattle, which is the host of *R. microplus*.

Furthermore, serine proteases, such as tissue kallikrein, have been related to proteolysis in *R. microplus* eggs [13]. Thus, serine protease inhibitors are important candidates for studying this process in eggs. In attempt to investigate the activity of serine protease inhibitors, we performed the purification and characterization of a Kunitz double-domain tissue kallikrein inhibitor found in *R. microplus* eggs, named RmKK. In addition, it was produced the recombinant RmKK in *Pichia pastoris* system. Finally, we provide an interesting discussion about the possible traffic of RmKK from midgut to the eggs and its role in the *R. microplus* eggs.

### 2. Materials and methods

#### 2.1. Ticks

*R. microplus* adults were kindly provided by Dr. Itabajara Silva Vaz Junior from Federal University of Rio Grande do Sul (UFRGS). The adults were maintained at 27 °C to produce eggs. Eggs were used to prepare the crude extract. In addition, adults were dissected to obtain midguts to perform RNA extraction and cDNA preparation.

\* Corresponding author. Address: Department of Biochemistry, UNIFESP-EPM, Rua Três de Maio 100, 04044-020 São Paulo, SP, Brazil. Fax: +55 11 5572 3006.

E-mail address: [tanaka.bioq@epm.br](mailto:tanaka.bioq@epm.br) (A.S. Tanaka).

## 2.2. Purification of the tissue kallikrein inhibitor from *R. microplus* eggs

The eggs obtained from adult ticks were homogenized in 0.025 M Tris–HCl buffer pH 8.0 containing 0.02 M NaCl, afterwards the extract was centrifuged during 15 min, at 12,000 rpm, 4 °C and the supernatant was separated. The supernatant was applied to a trypsin–Sephacrose column previously equilibrated with 0.050 M Tris–HCl buffer pH 8.0. The fractions eluted with 0.5 M KCl pH 8.0 were immediately, neutralized with 1 M Tris–HCl buffer pH 8.0. Solution containing eluted proteins was dialyzed against 0.05 M sodium acetate buffer pH 5.5. Following, we performed an ion exchange chromatography on a Resource S column connected to an AKTA Purifier System, previously equilibrated with the same buffer. The proteins were eluted by a linear NaCl gradient (0–1 M), with flow rate of 1 mL/min for 60 min. The fractions containing inhibitory activity were pooled. All purification steps were analyzed by SDS–PAGE electrophoresis using 15% polyacrylamide gels [14]. The purified protein was also used for N-terminal amino acid determination by Edman degradation [15] using a PPSQ-23 protein sequencer (Shimadzu).

## 2.3. Cloning of RmKK DNA fragment into pPIC9 expression vector

Nucleotide sequence encoding RmKK (accession number TC6491) was presented in the *R. microplus* gene index project ([http://comp-bio.dfci.harvard.edu/tgi/cgi-bin/tgi/gimain.pl?gudb=b\\_microplus](http://comp-bio.dfci.harvard.edu/tgi/cgi-bin/tgi/gimain.pl?gudb=b_microplus)), which was made available by Dr. Felix D. Guerrero from Knippling Bushland US Livestock Insect Research Laboratory – USA. The DNA fragment encoding the inhibitor was amplified by PCR using a cDNA preparation of *R. microplus* midgut as template in 50 µL reaction volume containing 5 pmol of gene-specific primers, 200 µM dNTPs, 1.5 mM MgCl<sub>2</sub>, and 5 U Taq DNA polymerase (Fermentas). PCR conditions were: 94 °C for 5 min, 94 °C for 45 s, 50 °C for 45 s and 72 °C for 45 s 35 cycles. DNA final extension was carried out at 72 °C for 10 min. The primers used were: RmKKFW 5′-CCGCTCGAGA AAAGAGTCTTCTCAATGTGACACTG-3′ and RmKKRev 5′-TTTTCCT TTTGCGGCGCTTAGCGCGGGAATGAC-3′. The PCR product was purified from 1% agarose gel using a QIAEX II Gel Extraction kit (Qiagen, Hilden, Germany), followed by digestion with XhoI and NotI restriction enzymes and cloning into pPIC9 vector, which was previously digested with the same enzymes.

## 2.4. Production of recombinant RmKK (rRmKK)

After cloning, the construction pPIC9-RmKK was linearized with SacI restriction enzyme and *P. pastoris* GS115 yeast strain was transformed with the linearized plasmid using a Gene Pulser II electroporation system (Bio-Rad) according to manufacturer's instructions. Transformed yeasts were randomly selected, screened and the large-scale heterologous protein expressions were performed. *P. pastoris* grown up in BMGY medium [1% yeast extract (p/v), 2% peptone (p/v), 100 mM potassium phosphate, pH 6.0, 1.34% YNB (p/v), 40 µM biotin, 1% glycerol (p/v)]. After that, recombinant RmKK was produced by incubation of the yeast in BMMY medium [1% yeast extract (p/v), 2% peptone (p/v), 100 mM potassium phosphate, pH 6.0, YNB 1.34% (p/v), 40 µM biotin, 0.5% methanol (v/v)]. This procedure was carried out under stirring at 210 rpm at 30 °C for six days and the induction of protein expression was carried out by addition of methanol to a final concentration 0.5% (v/v) at intervals of 24 h.

## 2.5. Purification of recombinant RmKK

The supernatant of yeast culture containing RmKK was applied to a HiTrap Q column, connected to an AKTA™ purifier system (GE Healthcare) previously equilibrated with 0.05 M Tris–HCl pH 8.0.

The protein was eluted with a linear NaCl gradient (0–0.5 M) in 0.05 M Tris–HCl pH 8.0 containing 0.5 M NaCl at a flow rate of 1.5 mL/min for 60 min. The elution profile was monitored by absorbance at 280 nm and fractions were tested regarding inhibitory activity towards bovine trypsin using synthetic substrate. This purification step was analyzed by SDS–PAGE electrophoresis [10].

## 2.6. Mass spectrometry analysis of rRmKK

The rRmKK was purified by reverse phase chromatography in a C<sub>18</sub> Sephasil column connected to an AKTA Purifier System. The column was pre-equilibrated in 0.1% trifluoroacetic acid (TFA) and the proteins eluted with a linear acetonitrile gradient (0–90%), with flow rate of 1 mL/min for 60 min. rRmKK molecular mass was determined by MALDI TOF mass spectrometer (Bruker Autoflex®) operating in linear mode and calculated from the *m/z* peaks in the charge-distribution profiles of the multiple charged ions.

## 2.7. Dissociation constant (*K<sub>i</sub>*) determination

Different concentrations of the native and recombinant inhibitor were pre-incubated with bovine trypsin and porcine kallikrein in 0.1 M Tris–HCl buffer, pH 8.0 containing 0.15 M NaCl and 0.1% Triton X-100, at 37 °C. The residual enzyme activity was measured by incubation with chromogenic substrates, Tosyl-Gly-Pro-Arg-pNA and 2266 (D-Val-Leu-Arg-pNA), for bovine trypsin and porcine pancreatic kallikrein, respectively. Finally, dissociation constants were calculated by fitting the steady-state velocities to the Morrison equation ( $V_i/V_o = 1 - \{E_t + I_t + K_i - [(E_t + I_t + K_i)^2 - 4E_t I_t]^{1/2}\} / 2et$ ) for tight-binding inhibitors using a nonlinear regression analysis [16].

## 2.8. Determination of tissue-specific expression using PCR

PCR reactions were performed using cDNAs preparations from midgut, ovary, hemocyte, salivary gland and fat body of engorged *R. microplus*. The amplicons were analyzed on 1% (p/v) agarose gel.

# 3. Results

## 3.1. Purification and characterization of a tissue kallikrein inhibitor from *R. microplus* eggs

The native RmKK was purified by affinity chromatography using a trypsin–Sephacrose column (Fig. 1A) followed by ion exchange chromatography using a Resource S column (Fig. 1B) and reverse phase chromatography in a C<sub>8</sub> Sephasil column (Fig. 1C). The N-terminal sequencing of purified inhibitor provided 10 amino acid residues (VLLNVTLPV).

## 3.2. Nucleotide sequence analysis of RmKK

Nucleotide sequence encoding RmKK was found in the *R. microplus* gene index project (see Section 2.3.). Using the nucleotide sequence accession number TC6491 the DNA fragment coding RmKK was cloned and sequenced. Nucleotide sequence encoding RmKK presented 411-bp ORF translated to a Kunitz double domain protein with 137 amino acid residues. The amino acid sequence analysis revealed the presence of 10 cysteine residues and the protein presented a calculated molecular weight of 16.7 kDa and isoelectric point (pI) of 7.54.

Download English Version:

<https://daneshyari.com/en/article/10755063>

Download Persian Version:

<https://daneshyari.com/article/10755063>

[Daneshyari.com](https://daneshyari.com)