



Neuronal cells but not muscle cells are resistant to oxidative stress mediated protein misfolding and cell death: Role of molecular chaperones



Arunabh Bhattacharya^{a,c,1}, Rochelle Wei^{a,1}, Ryan T. Hamilton^a, Asish R. Chaudhuri^{a,b,d,*}

^a Barshop Institute for Longevity and Aging Studies, University of Texas Health Science Center at San Antonio, 15355 Lambda Drive, San Antonio, TX 78245, United States

^b Department of Biochemistry, University of Texas Health Science Center at San Antonio, 7703 Floyd Curl Drive, San Antonio, TX 78229, United States

^c Cellular and Structural Biology, University of Texas Health Science Center at San Antonio, 7703 Floyd Curl Drive, San Antonio, TX 78229, United States

^d South Texas Veterans Health Care System, San Antonio, TX 78229, United States

ARTICLE INFO

Article history:

Received 9 March 2014

Available online 28 March 2014

Keywords:

Protein misfolding
Heat shock proteins
Oxidative stress
Skeletal muscle
Neuron
Cell viability

ABSTRACT

Our recent study in a mouse model of familial-Amyotrophic Lateral Sclerosis (f-ALS) revealed that muscle proteins are equally sensitive to misfolding as spinal cord proteins despite the presence of low mutant CuZn-superoxide dismutase, which is considered to be the key toxic element for initiation and progression of f-ALS. More importantly, we observed differential level of heat shock proteins (Hsp's) between skeletal muscle and spinal cord tissues prior to the onset and during disease progression; spinal cord maintains significantly higher level of Hsp's compared to skeletal muscle. In this study, we report two important observations; (i) muscle cells (but not neuronal cells) are extremely vulnerable to protein misfolding and cell death during challenge with oxidative stress and (ii) muscle cells fail to mount Hsp's during challenge unlike neuronal cells. These two findings can possibly explain why muscle atrophy precedes the death of motor neurons in f-ALS mice.

© 2014 Elsevier Inc. All rights reserved.

1. Introduction

Protein misfolding and aggregation are considered as hallmark for number of neurodegenerative diseases including Alzheimer's disease (AD), Parkinson's disease (PD) and Amyotrophic Lateral Sclerosis (ALS) [1]. In order to maintain the functional structure of proteins and enzymes, several strategies have evolved including the molecular chaperone network to respond to chronic cellular stress. Molecular chaperones namely heat shock proteins (Hsp's) act as catalysts for proper folding of misfolded or unfolded proteins [1]. The importance of Hsp's in regulation of protein misfolding and aggregation in neurodegenerative diseases is demonstrated by observations that (i) overexpression of selective Hsp's delays the progression of disease in animal models of neurodegeneration

Abbreviations: f-ALS, familial-Amyotrophic Lateral Sclerosis; BisANS, 4,4'-dianilino-1,1'-binaphthyl-5,5'-disulfonic acid, dipotassium salt; Hsp, heat shock protein; tBHP, *tert*-butyl hydroperoxide.

* Corresponding author at: Barshop Institute for Longevity and Aging Studies, University of Texas Health Science Center at San Antonio, Texas Research Park Campus, 15355 Lambda Drive, San Antonio, TX 78245-3207, United States. Fax: +1 210 562 6110.

E-mail address: chaudhuri@uthscsa.edu (A.R. Chaudhuri).

¹ Equal contribution.

<http://dx.doi.org/10.1016/j.bbrc.2014.03.097>

0006-291X/© 2014 Elsevier Inc. All rights reserved.

[1–3] and (ii) Hsp's are colocalized with aggregated proteins in diseases such as AD and PD [1]. Moreover, Hsp's have recently been shown to trigger the solubilization of toxic protein aggregates [1]. Together, these findings demonstrate the essential role of Hsp's in the regulation of proteostasis in the central nervous system (CNS). In comparison, nothing is known about protein misfolding and its regulation by Hsp's in skeletal muscle although several studies in the literature demonstrate that selective overexpression of Hsp's in cultured cells and intact animals reduce myotube/muscle atrophy [4–7]. Studies from our group have consistently shown that misfolding of skeletal muscle cytosolic proteins is a consistent finding in mouse models that exhibit significant muscle atrophy [8–10].

While recently studying protein misfolding in spinal cord and skeletal muscle (two tissues primarily affected in ALS) from wild-type (WT) mice and G93A mouse model of familial-ALS (f-ALS), we made an interesting observation: the expression of all the Hsp's studied (40, 60, 70 and 90) were strikingly lower in skeletal muscle versus spinal cord in the same WT animals [11]. This observation would suggest that there is less likelihood for the repair of misfolded proteins in skeletal muscle versus spinal cord and could possibly translate into increased susceptibility of muscle cells to oxidative stress/damage mediated cell death versus cells present

in the CNS such as neurons. Therefore, in this study we asked (i) if muscle cells are more vulnerable to oxidative stress induced misfolding and cell death, compared to neuronal cells, (ii) if these findings are linked to the differential response of protective molecular chaperones. Here we first show that skeletal muscle cytosolic proteins from WT mice show a higher kinetic of protein misfolding in response to *in vitro* oxidative and heat stress, compared to spinal cord cytosolic proteins. Next we show that cultured muscle cells are highly susceptible to oxidative stress mediated protein misfolding and cell death compared to neuronal cells, which is intimately linked to their inability to induce a Hsp response, unlike in neuronal cells. Together, these findings suggest for the first time that differential level of Hsp's and their response are possible determinants for the integrity of neuronal cells (versus muscle cells) in the event of an oxidative challenge.

2. Materials and methods

2.1. Animals and cell lines

4-mo-old C57BL/6 WT mice were used for measuring the expression of Hsp's and *in vitro* cell free-system assay. All procedures for handling animals were reviewed and approved by the Institutional Animal Care and Use Committee of University of Texas Health Science Center at San Antonio. 88NPC cells were grown with N-2 medium (containing DMEM/F-12, penicillin, gentamicin, B27, transferring, insulin and β FGF) in Matrigel-coated 100 mm dishes. C2C12 cells were grown in 100 mm dishes with DMEM high glucose + 20% FBS + 100 U/ml Penicillin + 100 μ g/ml Streptomycin.

2.2. Measurement of change in protein surface hydrophobicity in a cell-free system

Skeletal muscle and spinal cord tissues from 4-mo-old WT mice were homogenized in buffer containing 50 mM Tris, pH 7.4 buffer containing 1 mM MgSO_4 and protease cocktail inhibitors followed by centrifugation at 100,000g for 1 h at 4 °C. Cytosolic proteins (100 μ g) were incubated for (i) 1 h in *tert*-butyl hydroperoxide (tBHP) at 37 °C or for (ii) 1 h at 25 °C, 37 °C, and 42 °C. After the incubation, a reaction mixture containing 50 μ g (1 mg protein/ml) proteins was set up for photo-labeling under UV light with 4,4'-dianilino-1,1'-binaphthyl-5,5'-disulfonic acid, dipotassium salt (BisANS, 0.1 mM) as described earlier [8,10]. Equal amounts of BisANS-labeled proteins (10 μ g) were subjected to 12% SDS-PAGE and BisANS fluorescence captured under 365 nm UV light with an AlphaImage™ 3400 and followed by overnight staining of gels in SYPRO Ruby to normalize for protein loading. SYPRO Ruby fluorescence was captured on a Typhoon 9400 Variable Mode Imager (Amersham Biosciences) under the settings 610 BP 30, 520 PMT, 532 nm.

2.3. Measurement of change in protein surface hydrophobicity in cultured cells

Cells were stressed with 0, 50, 100, and 200 μ M tBHP for 12 h in serum-free medium supplemented with 2% BSA. Cells were washed in ice-cold PBS, harvested in 50 mM Tris, pH 7.4 buffer containing 1 mM MgSO_4 , protease cocktail inhibitors and centrifuged at 100,000g for 1 h at 4 °C. A reaction mixture of 50 μ g (1 mg protein/ml) proteins with BisANS (0.1 mM) was set up for photo-labeling under UV light as described earlier [8,10]. Equal amounts of BisANS-labeled proteins (10 μ g) were subject to 12% SDS-PAGE, BisANS fluorescence captured and normalized to SYPRO Ruby as described above.

2.4. Heat shock proteins in 88NPC and C2C12 cells

Cells were stressed with 0, 50, 100, and 200 μ M tBHP for 12 h in serum-free medium supplemented with 2% BSA. Cells were washed in ice-cold PBS and harvested in 20 mM potassium phosphate buffer pH 7.4 containing 0.5 mM MgCl_2 , 1 mM EDTA and protease inhibitors. Cells were sonicated and centrifuged at 100,000g for 1 h at 4 °C and equal amount of proteins (20 μ g) were subject to 12% SDS-PAGE followed by western blot using anti-rabbit primary antibodies against Hsp-60 and -70, and β -Actin (loading control). HRP secondary antibodies were used to visualize the distribution of hsp's.

2.5. 88NPC and C2C12 WST-1 cell viability

Cells were plated into a 96-well plate at 30,000 cells/well in serum-free medium supplemented with 2% BSA and allowed to recover and attach for 24 h. Thereafter, cells were stressed with 0, 50, 100, and 200 μ M tBHP for 12 h in serum-free medium supplemented with 2% BSA. Upon completion of tBHP incubation, the plates were spun down at 1000g for 2 min and old medium was aspirated. 100 μ l of fresh serum-free medium supplemented with 2% BSA and 5 μ l of WST-1 reagent was added to each well and incubated at 37 °C for 3.5 h. The absorbance at 450–630 nm was read on a microplate reader.

3. Results

Our recent report showed that at any given time point, the expression of molecular chaperones, namely Hsp's differs considerably between spinal cord and skeletal muscle tissues from WT and G93A mouse model of familial-ALS [11]. The expression of all the Hsp's (Hsp-40, -60, -70 and -90) tested were significantly higher in spinal cord versus skeletal muscle [11]. We hypothesized that this dramatic difference in molecular chaperones could possibly explain the surprising finding that protein misfolding pattern is quite similar between spinal cord and muscle tissues of G93A mice, despite considerable difference in the expression of the mutant G93A protein between these tissues [11]. Because G93A overexpression in mice is known to increase endogenous oxidative stress [12–14] and level of Hsp's differ between spinal cord and skeletal muscle, here we first asked if spinal cord proteins in general are more resistant to stress induced misfolding, compared to skeletal muscle proteins in an *in vitro* assay in cytosolic protein extracts. We used 4-mo-old WT mice to perform these experiments that exhibit ~15- and 5-fold higher expression of Hsp-70 and Hsp-60, respectively, in spinal cord versus skeletal muscle tissues (Fig. 1A and B). Using the BisANS photo-labeling approach, we found that in response to oxidative stress (tBHP), the kinetics of hydrophobic domain exposure is slower in spinal cord proteins compared to skeletal muscle proteins (Fig. 2A and B). In response to heat shock stress, the difference between spinal cord and skeletal muscle proteins with respect to the kinetics of hydrophobic domain exposure is even more dramatic. Skeletal muscle proteins progressively lost their hydrophobic pockets with increasing temperature (~60% decrease at 42 °C compared to 25 °C), whereas, spinal cords proteins were completely resistant to unfolding (Fig. 2C and D). These results clearly indicate that the cellular level of Hsp's could be a critical component in our findings as the assays were performed in cytosolic extracts prepared in the presence of protease inhibitors, thus ruling out the effect of proteasome. Although our findings were interesting, we still didn't know if they translated into any functional consequence. To address this question, we first determined the global status of protein folding in C2C12 myoblasts and NPCs exposed to varying concentrations of tBHP. The data in

Download English Version:

<https://daneshyari.com/en/article/10755701>

Download Persian Version:

<https://daneshyari.com/article/10755701>

[Daneshyari.com](https://daneshyari.com)