



## Neuronal nitric oxide synthase is phosphorylated in response to insulin stimulation in skeletal muscle

Kathryn Hinchee-Rodriguez, Neha Garg<sup>1</sup>, Priya Venkatakrisnan<sup>2</sup>, Madeline G. Roman<sup>3</sup>, Martin L. Adamo, Bettie Sue Masters, Linda J. Roman\*

Department of Biochemistry, The University of Texas Health Science Center at San Antonio, 7703 Floyd Curl Dr., San Antonio, TX 78229-7760, USA

### ARTICLE INFO

#### Article history:

Received 30 April 2013

Available online 14 May 2013

#### Keywords:

Nitric oxide synthase  
Type 2 Diabetes (T2DM)  
Insulin signaling  
Myotubes  
Skeletal muscle

### ABSTRACT

Type 2 Diabetes (T2DM) is the seventh leading cause of death in the United States, and is quickly becoming a global pandemic. T2DM results from reduced insulin sensitivity coupled with a relative failure of insulin secretion. Reduced insulin sensitivity has been associated with reduced nitric oxide synthase (NOS) activity and impaired glucose uptake in T2DM skeletal muscle. Upon insulin stimulation, NO synthesis increases in normal adult skeletal muscle, whereas no such increase is observed in T2DM adults. Endothelial NOS is activated by phosphorylation in the C-terminal tail in response to insulin. Neuronal NOS (nNOS), the primary NOS isoform in skeletal muscle, contains a homologous phosphorylation site, raising the possibility that nNOS, too, may undergo an activating phosphorylation event upon insulin treatment. Yet it remains unknown if or how nNOS is regulated by insulin in skeletal muscle. Data shown herein indicate that nNOS is phosphorylated in response to insulin in skeletal muscle and that this phosphorylation event occurs rapidly in C2C12 myotubes, resulting in increased NO production. *In vivo* phosphorylation of nNOS was also observed in response to insulin in mouse skeletal muscle. These results indicate, for the first time, that nNOS is phosphorylated in skeletal muscle in response to insulin and in association with increased NO production.

Published by Elsevier Inc.

### 1. Introduction

Type 2 Diabetes mellitus (T2DM) is characterized by insulin resistance and impaired insulin-stimulated nitric oxide (NO) generation in both endothelium and skeletal muscle. NO is an important gaseous regulatory molecule, produced by isoforms of nitric oxide synthase (NOS). NO acts variably as a vasodilator [1,2], neurotransmitter [3] or cytotoxic agent [4], depending upon its source and location of production. NOS enzymes synthesize NO from L-arginine and molecular oxygen [5], utilizing electrons donated by NADPH. The homeostatic regulation of NO synthesis is crucial in biological systems, as the generation of insufficient amounts of NO can result in deleterious outcomes, such as increased blood pressure due to uncontrolled vasoconstriction. In contrast, overproduction of NO can yield toxic sequelae, such as endotoxic shock.

Neuronal NOS (nNOS) and endothelial NOS (eNOS) knockout mice demonstrate insulin resistance, suggesting that NO produced by nNOS plays an important role in mediating insulin action [6,7]. However, the direct effect of insulin on skeletal muscle nNOS is unknown. One prior study by Kashyap et al. [8], using euglycemic/hyperinsulinemic clamps, showed that NO production in muscle biopsies is significantly reduced in obese, T2DM human subjects compared to obese, non-diabetic controls. Moreover, in the insulin-resistant subjects, insulin treatment failed to evoke increases in NO in skeletal muscle, whereas insulin stimulated a robust increase in NO in the control subjects [8]. These results suggest that insulin action in skeletal muscle involves NO production, thus implicating mechanisms regulating NO bioavailability. These mechanisms must be elucidated to understand the pathogenesis of muscle insulin resistance, which is an important component of the pathology of T2DM.

Phosphorylation plays a critical role in the potentiation of NO generation by vascular eNOS, which is activated by insulin-dependent AKT-catalyzed phosphorylation at eNOS Ser1177 [9,10], potentially explaining the vasodilatory effects of insulin. While the role of eNOS in insulin action and its inactivation as a component of insulin resistance in the vasculature has been well-documented [2,11–13], the metabolic role of NOS in other tissues, such as skeletal muscle, remains elusive. The predominant NOS

\* Corresponding author. Fax: +1 210 567 6984.

E-mail address: [roman@uthscsa.edu](mailto:roman@uthscsa.edu) (L.J. Roman).

<sup>1</sup> Present address: Department of Genetics and The Paul F. Glenn Laboratories for the Biological Mechanisms of Aging, Harvard Medical School, Boston, MA, USA.

<sup>2</sup> Present address: Integrity Bio, Inc., Camarillo, CA 93012, USA.

<sup>3</sup> Present address: Barshop Institute for Longevity and Aging Studies, The University of Texas Health Science Center at San Antonio, 15355 Lambda Drive, San Antonio, TX 78245-3207, USA.

isoform in skeletal muscle is nNOS, several splice variants of which, i.e., nNOS $\mu$ , nNOS $\beta$  and nNOS $\alpha$  [14,15], may be present. In this report, we do not distinguish among the nNOS variants, hereafter collectively referred to as skeletal muscle nNOS.

The rapid increase in NO production observed in the human insulin clamp studies suggests that post-translational modification of skeletal muscle nNOS occurs in direct response to insulin rather than as insulin-regulated transcription [8]. Since skeletal muscle nNOS contains a homologous residue to the AKT-phosphorylated serine in eNOS (Ser1177), it was possible that nNOS could be phosphorylated in response to insulin in skeletal muscle. There is evidence of nNOS phosphorylation in response to insulin in brain: prior immunoblot studies in rat hypothalamus demonstrated an increase in nNOS Ser1416 phosphorylation [16], the rat equivalent of Ser1412 in mouse nNOS. *In vitro* kinase assays in rat brain nucleus *Tractus solitarii* additionally showed nNOS is phosphorylated at this residue in an insulin-dependent manner [17].

To examine nNOS phosphorylation by insulin treatment, the murine muscle C2C12 cell line, which can be differentiated from myoblasts to myotubes, was used to probe putative Ser1412 phosphorylation on the C-terminal tail of skeletal muscle nNOS in response to stimulation by insulin. Our results show that insulin stimulation resulted in significantly increased phosphorylation of skeletal muscle nNOS in C2C12 myotubes, as well as a concomitant increase in NO levels. *In vivo*, insulin-stimulated nNOS phosphorylation was also observed in WT mouse C57BL/6J muscle tissue, further validating that insulin-mediated phosphorylation of nNOS is an important component of the insulin-signaling cascade.

## 2. Materials and methods

### 2.1. Cell culture and treatment

Murine-derived muscle cell line C2C12 (Cell Biology Labs) myoblasts were grown to near 100% confluency in high glucose (25 mM) DMEM (Sigma–Aldrich) with 10% fetal bovine serum (Gibco), hereafter called complete media. Myoblasts were then switched to high glucose DMEM media, 2% horse serum (Gibco), hereafter called horse serum media. After four days in serum starvation, cells were completely differentiated to myotubes, and were treated with either vehicle (3  $\mu$ M HCl) or 100 nM insulin (Sigma Aldrich) from 0 to 60 min.

### 2.2. Animals

Wild-type male C57BL/6J mice were fed *ad libitum* a standard lab chow (Harlan, 11.5 kcal% fat) and maintained in micro-isolator cages, 5 to a cage, on a 12 h dark/light cycle. Mice (4–6 months of age) were given intraperitoneal injections of either vehicle (PBS) or 5 mU/g body weight insulin (Novo Nordisk). Ten minutes after insulin injection, mice were sacrificed and quadriceps femoris muscles were collected for analysis.

All animal studies have been approved by Institutional Animal Care and Use Committee (IACUC), University of Texas Health Science Center at San Antonio, San Antonio, TX, USA. Animals are housed in an Association for Assessment and Accreditation of Laboratory Animal Care, International (AAALAC) accredited facility with full veterinary support. The facility is operated in compliance with the Public Law 89-544 (Animal Welfare Act) and Amendments, Public Health Services Policy on Humane Care and Use of Laboratory Animals (PHS Policy) and The Guide for the Care and Use of Laboratory Animals.

### 2.3. NO detection

The NO-specific fluorescent dye 4,5-diaminofluorescein diacetate (DAF-2 DA; Sigma) was used to measure NOS activity indirectly. C2C12 myoblasts were seeded onto 6-well plates, maintained in complete media and differentiated into myotubes with horse serum-containing media. For fluorescent microscopy imaging, myotubes were treated with either vehicle or 100 nM insulin in horse serum media for 60 min, rinsed 2 $\times$  with PBS, then treated with 5  $\mu$ M DAF-2 DA in PBS + glucose for 30 min. After rinsing, clear  $\alpha$ -MEM (Sigma–Aldrich) was applied to the living cells, which were then imaged with at 10 $\times$  using a Zeiss Axioscope 2 HBO 100 with excitation wavelength at 495 nm and emission at 515 nm. For quantitative analysis of DAF-2 DA fluorescence, myotubes were treated with 5  $\mu$ M DAF-2 DA in PBS + glucose for 30 min, rinsed 2 $\times$  with PBS, then treated with either vehicle or 100 nM insulin in horse serum media for 60 min. As a positive control, cells treated with 450  $\mu$ M S-nitroso-N-acetyl-D,L-penicillamine (SNAP, Cayman Chemical). After rinsing, PBS was applied to the cells, and measurements were made in a Tecan plate reader with excitation wavelength at 475 nm and emission at 550 nm.

### 2.4. Immunoblot analyses

Cell lysates from C2C12 cells were prepared by collecting cells in PBS with phosphatase and protease inhibitors (Thermo-Fisher) and Benzonase, and then homogenized. For mouse quadriceps muscle lysates, after an overnight fast, mice were given an intraperitoneal injection of 5 mU /g body weight of insulin (Novo Nordisk) or an equivalent volume of sterile saline. Ten min post injection, quadriceps muscle was collected and frozen in liquid nitrogen, then whole tissue homogenates were prepared. For all immunoblots, thirty micrograms of lysates were electrophoresed in gradient SDS–PAGE (2–15%; Bio-Rad) and proteins were transferred onto PVDF membrane (Millipore). Membranes were probed with antibodies to phospho-S1417 nNOS (recognizes mouse nNOS at Ser1412, Abcam), total nNOS (BD Biosciences), phospho-S473 AKT (Cell Signaling Technology), total AKT (Cell Signaling Technology) and/or GAPDH (Cell Signaling Technology), as indicated, followed by a 1 h incubation in horseradish-peroxidase linked secondary antibody (Santa Cruz). Detection was by Immobilon Western Chemiluminescent HRP Substrate (Millipore). Immunoblots were analyzed using ImageJ [18].

## 3. Results/discussion

### 3.1. nNOS is phosphorylated in response to insulin in C2C12 myotubes

To determine whether nNOS is phosphorylated in skeletal muscle in response to insulin, the murine-derived *in vitro* muscle cell line C2C12 was used as a model system. Control and insulin-treated C2C12 cell lysates were probed for nNOS phosphorylated on residue S1412 via immunoblot analysis. As shown in Fig. 1A, insulin treatment resulted in increased nNOS phosphorylation at Ser1412 in a time-dependent manner, with significant ( $P < 0.05$ ) phosphorylation occurring at 15, 30 and 60 min post-insulin addition. Fig. 1B shows the ratio of phosphorylated nNOS to total nNOS, clearly demonstrating the significant increase of this event, particularly at the 30 min time point. Phosphorylation of nNOS occurred concomitantly with activation of the insulin-signaling pathway, shown by the increase in phospho-AKT at Ser473 by insulin treatment (Fig. 1A). This result suggests that nNOS phosphorylation is a consequence of activated insulin signaling in C2C12 myotubes.

Download English Version:

<https://daneshyari.com/en/article/10759356>

Download Persian Version:

<https://daneshyari.com/article/10759356>

[Daneshyari.com](https://daneshyari.com)