



Identification of p21 (CIP1/WAF1) as a direct target gene of HIC1 (Hypermethylated In Cancer 1)

Vanessa Dehennaut¹, Ingrid Loison, Gaylor Boulay², Capucine Van Rechem², Dominique Leprince^{*}

CNRS-UMR 8161, «Institut de Biologie de Lille», Université de Lille Nord de France, Institut Pasteur de Lille, IFR 142, 1 rue Calmette, BP447, 59017 Lille Cedex, France

ARTICLE INFO

Article history:

Received 30 October 2012

Available online 22 November 2012

Keywords:

HIC1

p21

Tumor suppressor

DNA damages

Transcription

ABSTRACT

The tumor suppressor gene *HIC1* (*Hypermethylated In Cancer 1*) encodes a transcriptional repressor involved in the regulation of growth control and DNA damage response. We previously demonstrated that *p57Kip2*; a member of the CIP/KIP family of CDK (cyclin dependent kinase) inhibitors (CKI); is a direct target gene of HIC1 in quiescent cells. Here we show that ectopic expression of HIC1 in MDA-MB-231 cells or its overexpression in BJ-Tert fibroblasts induces decreased mRNA and protein expression of p21 (CIP1/WAF1) another member of this CKI family that plays essential roles in the p53-mediated DNA damage response. Conversely, knock-down of endogenous HIC1 in BJ-Tert through RNA interference up-regulates p21 in basal conditions and further potentiates this CKI in response to apoptotic etoposide-induced DNA damage. Through promoter luciferase activity and chromatin immunoprecipitation (ChIP), we demonstrate that HIC1 is a direct transcriptional repressor of *p21*. Thus, our results further demonstrate that HIC1 is a key player in the regulation of the DNA damage response.

© 2012 Elsevier Inc. All rights reserved.

1. Introduction

HIC1 (*Hypermethylated In Cancer 1*), a tumor suppressor gene frequently hypermethylated or deleted in numerous cancers, encodes a transcriptional repressor [1–3]. The HIC1 protein is composed of three main functional domains: a BTB/POZ protein–protein interaction domain (Broad complex, Tramtrack and Bric à brac/POX viruses and zinc finger) in the N-terminal part of the protein, a central region and a C-terminal domain containing five Krüppel-like C₂H₂ zinc fingers. These zinc fingers allow the specific binding of the protein to specific DNA sequences consisting of a 5'-(C/G)NG(C/G)GGGCA(C/A)CC-3' centered on a GGCA motif and named HIC1 responsive elements (HiRE) [4]. The central region of HIC1 contains two short phylogenetically conserved motifs: (i) GLDLSKK, allowing the recruitment of the co-repressor CtBP (C-terminal Binding Protein) [5,6] and (ii) MK³¹⁴HEP, whose lysine is competitively acetylated or SUMOylated [7] to regulate the interaction between HIC1 and MTA1; a member of the NuRD (Nucleosome Remodeling and histone Deacetylase) complex [8]. To date only 11 direct target genes of HIC1 have been described including *p57kip2* [8] a cyclin-dependent kinase inhibitor belonging to the

CIP/KIP family, which also includes p21(CIP1/WAF1) and p27(KIP1) [9]. Several studies have highlighted a central role for HIC1 in the regulation of the DNA damage response and more particularly through the existence of a complex regulatory loop between HIC1, p53 and the deacetylase SIRT1 [10]. First HIC1 is a direct target gene of p53 [1,11,12], the master regulator of the DNA damage-induced checkpoints which, depending on the damage intensity, transactivates various growth inhibitory or apoptotic genes. Among these are *p21* [13], promoting G1 growth arrest and *SIRT1* [14] that in a feed-back loop deacetylates and inactivates p53. HIC1 directly represses the transcription of *SIRT1* [15] whereas SIRT1 by deacetylating the lysine 314 of HIC1 increases its transcriptional repression potential [7]. In this study, we show that *p21* is a new direct target gene of HIC1 thus adding a new step to the p53-HIC1-SIRT1 regulatory loop and further implicating HIC1 as an essential element in the regulation of the DNA damage response.

2. Materials and methods

2.1. Cell lines and plasmids

HEK 293T, U2OS, MDA-MB-231 and BJ-Tert cells were maintained in Dulbecco modified Eagle medium (Invitrogen) supplemented with 10% fetal calf serum, non-essential amino acids and gentamycin. Cells were cultured at 37 °C in water-saturated 5% CO₂ atmosphere. The pcDNA3-FLAG-HIC1 has been previously described [6].

^{*} Corresponding author. Fax: +33 3 20 87 11 11.

E-mail address: dominique.leprince@ibl.fr (D. Leprince).

¹ Present address: CNRS-UMR 8576, Unité de Glycobiologie Structurale et Fonctionnelle, Université de Lille 1, Bâtiment C9, 59655 Villeneuve d'Ascq Cedex, France.

² Present address: Harvard Medical School and Massachusetts General Hospital Cancer Center, Charleston, MA 02129, USA.

2.2. Vector and retroviral infection

Retroviral infection of BJ-Tert fibroblasts with the pBABE-Puro-FLAG-HIC1 [16] were performed as previously described [17].

2.3. Small Interfering RNA

BJ-Tert fibroblasts were reverse-transfected with Lipofectamine RNAiMax (Invitrogen) according to manufacturer's instructions using 10 nM small interfering RNA targeting HIC1 (HIC1 siGENOME SMART Pool M-006532-01, Dharmacon), MTA1 (MTA1 siGENOME SMART Pool M-004127-02, Dharmacon), or a scrambled control sequence (si Ctrl; siGENOME RISC free control siRNA, Dharmacon) as previously described [17] 48 h later, cells were treated with 80 μ M etoposide for 16 h and harvested for RNA/protein extraction [16,17].

2.4. Luciferase repression assays

The pGL3 p21 promoter construct has been kindly provided by Olivier Rohr (University of Strasbourg, France) and has been previously described [18].

HEK 293T or U2OS cells were transfected in OptiMEM (Invitrogen) by the PEI (Euromedex) method in 12-well plates with 500 ng of DNA. Cells were transfected for 6 h and then incubated in complete fresh medium. 48 h after transfection, cells were lysed in Luc assay buffer (25 mM glycyl glycine [pH 7.8], 15 mM MgSO₄, 4 mM EGTA, 1% Triton X-100). Luciferase and β -galactosidase activities were measured by using, respectively, beetle luciferine (Promega) and the Galacto-light kit (Tropix) with a Berthold chemiluminometer. After normalization to the β -galactosidase activity, the data were expressed as the Luc activity relative to the activity of pGL3-Luc with empty control vector, which was given an arbitrary value of 1. The results represent the mean values and standard deviations from 3 independent experiments.

2.5. Quantitative RT-PCR

Total RNA was reverse transcribed using random primers and MultiScribe™ reverse transcriptase (Applied Biosystems). Real-time PCR analysis was performed by Power SYBR Green (Applied Biosystems) in a MX3005P fluorescence temperature cycler (Stratagene) according to the manufacturer's instructions. Results were normalized with respect to 18S RNA used as internal control. The primers used are as follows: HIC1 forward 5'-CGACGACTACAAG AGCAGCAGC-3' and reverse 5'-CAGGTTGTACCGAAGCTCTC -3'; p21 forward 5'-GACTCTCAGGGTCGAAAACG-3' and reverse 5'-GGC TTCCTCTTGGAGAAGATCA-3' and 18S forward 5'-GGCGCCCC CTCGATGCTCTTAG-3' and reverse 5'-GCTCGGGCCTGCTTTGAACA CTCT-3'.

2.6. Chromatin immunoprecipitation

BJ-Tert cells were fixed by adding formaldehyde directly into the cell plate to a final concentration of 1% for 15 min at 37 °C. Adding glycine to a final concentration of 0.125 M stopped the cross-linking. After 5 min at 37 °C, cells were lysed directly in the plates by resuspension in cell lysis buffer (5 mM PIPES pH 8, 85 mM KCl, 0.5% NP-40) for 5 min. Then, the samples were pelleted, resuspended in 100 μ l of nuclei lysis buffer (50 mM Tris-HCl pH 8, 10 mM EDTA, 0.2% SDS), and sonicated to chromatin with an average size of 250 bp using a BioRuptor (Diagenode, Liege, Belgium). 20 μ g of chromatin was transferred to a tube containing 4.8 μ g of anti-HIC1 [19] antibody or IgG control coated to magnetic protein A beads (Millipore) and 900 μ l of IP buffer (16.7 mM Tris-HCl pH 8, 1.2 mM EDTA, 167 mM NaCl, 0.01% SDS, 1.1% Triton X-100) and

incubated for 6 h on a rotator at 4 °C. Immune complexes were washed once in IP buffer, once in TSE buffer (20 mM Tris pH 8, 2 mM EDTA, 500 mM NaCl, 1% Triton X-100, 0.1% SDS), once in LiCl buffer (100 mM Tris pH 8, 500 mM LiCl, 1% deoxycholic acid, 1% NP40) and twice in TE buffer (10 mM Tris/HCl, pH 8, 10 mM EDTA). Each wash lasted for 3 min on a rotator at room temperature. ChIP complexes were eluted in 150 μ l of elution buffer (10 mM Tris/HCl, pH 7.5, 50 mM NaCl, 1% SDS) containing 50 μ g/ml of proteinase K and incubated 1 h at 55 °C. Then samples were treated with 133 μ g/ml of RNase A for 30 min at 37 °C. Finally, the supernatant was recovered and incubated for 3 h at 68 °C. DNAs as well as 5% input were purified on Nucleobond Extract II (Macherey-Nagel) and eluted with 80 μ l of H₂O.

Immunoprecipitated DNA was analyzed in a MX3005P fluorescence temperature cycler (Stratagene) in triplicates by real time PCR starting from 3 μ l of template DNA in a final volume of 20 μ l containing power SYBR Green (Applied Biosystems) and primers at a final concentration of 0.5 μ M. The primers used are summarized in [Supplementary Table S1](#). According to a melting point analysis, only one PCR product was amplified under these conditions. An input control was used to generate a standard curve for each gene. Results were expressed as % input. The experiment was performed twice and a representative experiment is shown.

2.7. Western blotting and antibodies

Proteins were separated by SDS-PAGE and transferred onto nitrocellulose membranes (GE healthcare). After 1 h of blocking in PBSM (PBS with 5% milk), the membranes were incubating overnight at 4 °C with specific primary antibodies in PBSTM (PBSM with 0.1% Tween) and washed three times with PBSN (PBS with 0.1% NP-40). The membranes were next incubated for 1 h at room temperature with secondary antibodies coupled to peroxidase (Amersham) in PBSM, washed three times in PBSN and revealed by chemiluminescence.

The anti-HIC1 antibody has been previously described [17]. Anti-p21, anti-p53 and anti-actin were purchased from Santa-cruz biotechnology and anti γ -H2AX from Abcam.

3. Results

3.1. Effects of modulating HIC1 expression on the levels of p21

To test whether p21 could be a new target gene of HIC1, we first investigated the effects of HIC1 ectopic expression in HIC1-deficient breast cancer cells, MDA-MB-231 or overexpression in BJ-Tert "normal" human fibroblasts on p21 mRNA ([Fig. 1A](#)) and protein levels ([Fig. 1B](#)). After retroviral infection of pBabe-HIC1 in these two cell lines, we observed a significant decrease in p21 mRNA levels ([Fig. 1A](#)) correlated with a lower expression of the protein ([Fig. 1B](#)) in comparison with the cells infected with the empty vector. Conversely, inhibition of endogenous HIC1 expression in BJ-Tert fibroblasts by RNA interference [16,17] leads to a concomitant increase in p21 mRNA and protein levels ([Fig. 2](#)). This increase was observed in basal conditions but more interestingly we demonstrated that knock-down of HIC1 super-induces p21 in response to etoposide-induced DNA damages. In addition, inhibition of HIC1 leads to a modest increase of total p53 protein levels either in non-treated or in the etoposide-treated cells, reflecting the complexity of this regulatory loop. However, our results strongly suggests that the increased p21 expression observed in the cells transfected with the siRNA targeting HIC1 is not merely due to an increased expression of p53 resulting in an increased transcription of p21.

Download English Version:

<https://daneshyari.com/en/article/10760168>

Download Persian Version:

<https://daneshyari.com/article/10760168>

[Daneshyari.com](https://daneshyari.com)