



N-n-Butyl haloperidol iodide inhibits the augmented $\text{Na}^+/\text{Ca}^{2+}$ exchanger currents and L-type Ca^{2+} current induced by hypoxia/reoxygenation or H_2O_2 in cardiomyocytes

Yongpan Huang^a, Fenfei Gao^a, Yanmei Zhang^a, Yicun Chen^a, Bin Wang^a, Yanshan Zheng^a, Ganggang Shi^{a,b,*}

^a Department of Pharmacology, Shantou University Medical College, Shantou, China

^b Department of Cardiovascular Diseases, First Affiliated Hospital, Shantou University Medical College, Shantou, China

ARTICLE INFO

Article history:

Received 13 March 2012

Available online 2 April 2012

Keywords:

N-n-butyl haloperidol iodide

$\text{Na}^+/\text{Ca}^{2+}$ exchanger currents

L-type Ca^{2+} current

Ventricular myocytes

Patch-clamp techniques

ABSTRACT

N-n-butyl haloperidol iodide (F_2), a novel quaternary ammonium salt derivative of haloperidol, was reported to antagonize myocardial ischemia/reperfusion injuries. To investigate its mechanisms, we characterized the effects of F_2 on $\text{Na}^+/\text{Ca}^{2+}$ exchanger currents (I_{NCX}) and the L-type Ca^{2+} channel current ($I_{\text{Ca,L}}$) of cardiomyocytes during either hypoxia/reoxygenation or exposure to H_2O_2 . Using whole-cell patch-clamp techniques, the I_{NCX} and $I_{\text{Ca,L}}$ were recorded from isolated rat ventricular myocytes. Exposure of cardiomyocytes to hypoxia/reoxygenation or H_2O_2 enhanced the amplitude of the inward and outward of I_{NCX} and $I_{\text{Ca,L}}$. F_2 especially inhibited the outward current of $\text{Na}^+/\text{Ca}^{2+}$ exchanger, as well as the $I_{\text{Ca,L}}$, in a concentration-dependent manner. F_2 inhibits cardiomyocyte I_{NCX} and $I_{\text{Ca,L}}$ after exposure to hypoxia/reoxygenation or H_2O_2 to antagonize myocardial ischemia/reperfusion injury by inhibiting Ca^{2+} overload.

© 2012 Elsevier Inc. All rights reserved.

1. Introduction

Ischemic heart disease is the leading cause of morbidity and mortality worldwide. Subsequent reperfusion of acutely ischemic myocardium is essential for myocardial rescue, but also leads to a unique type of injury known as myocardial ischemia/reperfusion (I/R) injury [1]. Such an injury is often related to endothelial and microvascular dysfunction, impaired blood flow, metabolic dysfunction, and cellular necrosis [2], and its mechanism is associated with cytosolic and mitochondrial calcium overload, release of reactive oxygen species (ROS), and an acute inflammatory response [3,4]. As one of the important mechanisms of I/R injury, much research has focused on the precise intracellular signaling pathways and elements responsible for calcium overload in ischemia/reperfusion. Ca^{2+} influx via both activation of L-type calcium channel and reversal of the $\text{Na}^+/\text{Ca}^{2+}$ exchanger (NCX) have been reported to occur in cardiocytes following I/R [1,5,6]. Simultaneously, ROS, including superoxide radicals, hydroxyl radicals, and oxidants such as H_2O_2 are generated in significant amounts

during reperfusion and could contribute to intracellular Ca^{2+} overload in the heart through reversal or inhibition of the NCX [7,8]. Calcium overload may lead to deleterious consequences such as stunning, apoptosis, and necrosis, which contribute to infarct formation [9–12]. Due to the pivotal role of calcium overload in I/R injury, attenuation of cellular calcium overload remains an important therapeutic goal.

N-n-butyl haloperidol iodide, a novel quaternary ammonium salt derivative of haloperidol, was found to maintain the cardiac and vascular effects without adverse extrapyramidal reactions. Our previous studies showed that F_2 could block L-type calcium channels in ventricular myocytes under physiological conditions [13–15]. Subsequently, we demonstrated that F_2 could antagonize myocardial I/R injury in different animal models [13,16]. So, we inferred that the mechanism by which F_2 antagonizes myocardial I/R injury might be related to the inhibition of Ca^{2+} overload via suppression of cardiomyocyte $\text{Na}^+/\text{Ca}^{2+}$ exchanger (I_{NCX}) currents and L-type Ca^{2+} channel ($I_{\text{Ca,L}}$) during I/R. In this study, we established a model of cardiomyocyte hypoxia/reoxygenation (H/R) and exposure to H_2O_2 , to simulate heart I/R conditions, and characterized the changes of I_{NCX} and $I_{\text{Ca,L}}$ during H/R and exposure to H_2O_2 . We further characterized the effects of F_2 on I_{NCX} and $I_{\text{Ca,L}}$ during H/R and exposure to H_2O_2 to elucidate the mechanisms and ability of F_2 to block myocardial I/R injury.

* Corresponding author at: Department of Pharmacology, Shantou University Medical College, 22 Xinling Road, Shantou, Guangdong 515041, China. Fax: +86 754 88557562.

E-mail addresses: yongpanhuangxy@yahoo.cn, ggshi@stu.edu.cn (G. Shi).

2. Materials and methods

2.1. Cell isolation

Adult male Sprague–Dawley rats (180–250 g) were obtained from the Laboratory Animal Breeding and Research Center of Shantou University Medical College. The investigation was in compliance with the *Guide for the Care and Use of Laboratory Animals* published by the US National Institutes of Health (NIH Publication No. 85-23, revised 1996). Cardiomyocytes were isolated by collagenase type 2 and protease (Sigma, Type XIV) perfusion as previously described [15,17]. All experiments were performed at $37 \pm 0.5^\circ\text{C}$. Cardiac ventricular tissue was cut into small pieces. Single myocytes were harvested and stored at 4°C and the myocytes were used for experiments within 6 h.

2.2. Patch clamp recordings

Membrane currents were recorded by whole cell patch-clamp method using pCLAMP 8.2 software (Axon Instruments, Foster City, CA, USA). Single cardiac ventricular cells were placed in a 1 ml recording chamber attached to an inverted microscope (OLYMPUS, Tokyo) and were perfused with Tyrode solution at a rate of 1 ml/min. The temperature of the bath solution was maintained at room temperature ($22\text{--}25^\circ\text{C}$). Patch pipettes were forged from 1.5-mm-diameter glass capillaries with a two-stage microelectrode puller (pp-83; Narishige Scientific Instrument Lab, Tokyo). Pipette resistance was $2\text{--}4\text{ M}\Omega$ when filled with the pipette solution. The electrode was connected to a patch-clamp amplifier (Axopatch-200B, Axon Instruments, Foster City, CA, USA). Recording signals were filtered at 2.5 kHz bandwidth.

2.3. Measurement of I_{NCX}

After establishing the whole-cell configuration in Tyrode solution, the cell was perfused with a special K^+ -free bath solution (140 mM NaCl, 2 mM CaCl_2 , 1 mM MgCl_2 , and 10 mM HEPES, pH 7.2). To block Na^+/K^+ pump currents and currents flowing through K^+ or Ca^{2+} channels, 0.02 mM ouabain, 2 mM CsCl and 0.01 mM nifedipine were added to the solution. After recording the control current, the external solution was switched from the special K^+ -free bath solution to the simulated hypoxic solution including 0.02 mM ouabain, 2 mM CsCl and 0.01 mM nifedipine. The pipette solution contained 120 mM CsOH, 50 mM aspartic acid, 20 mM NaCl, 10 mM CaCl_2 (free Ca^{2+} concentration 226 nM), 20 mM BAPTA, 3 mM MgCl_2 , 5 mM Mg ATP, and 10 mM HEPES, pH 7.2). The ramp pulse was initially depolarized from a holding potential of -60 to $+60$ mV, then hyperpolarized to -150 mV, and depolarized back to the holding potential at a speed of 680 mV/s [18]. The descending limb of the ramp was used to plot current–voltage (I – V) curves without capacitive current compensation. I_{NCX} was identified as a Ni^{2+} -sensitive current because 5 mM Ni^{2+} selectively inhibits I_{NCX} under these ionic conditions and the Ni^{2+} -insensitive current was not affected by H/R.

2.4. Measurement of $I_{\text{Ca,L}}$

$I_{\text{Ca,L}}$ was recorded using a whole-cell patch clamp configuration. The pipette solution contained 150 mM CsCl, 15 mM EGTA, 1 mM MgCl_2 , 5 mM MgATP, and 5 mM HEPES, adjusted to pH 7.2 with CsOH). After establishing a high resistance seal by gentle suction, the cell membrane beneath the tip of the electrode was disrupted by further suction to obtain the whole-cell patch-clamp configuration. $I_{\text{Ca,L}}$ was elicited by 300 ms pulses to potentials ranging from -30 to $+70$ mV in 10 mV increments from a holding potential of

-40 mV (to inactivate I_{Na} and T-type Ca^{2+} currents) at 0.2 Hz [15]. Peak outward K^+ current (I_{to}) was suppressed by 3 mM 4-aminopyridine added to Tyrode solution. Representative current traces and the I – V relationships were obtained from a ventricular myocyte.

2.5. H/R model

H/R conditions were induced by switching the Tyrode solution to the hypoxic solution and then to control extracellular solution. After perfusing with normal Tyrode solution, to mimic hypoxic conditions, isolated ventricular cardiac myocytes were perfused for 15-min with the simulated hypoxic solution (123 mM NaCl, 6 mM NaHCO_3 , 0.9 mM NaH_2PO_4 , 8 mM KCl, 0.5 mM MgSO_4 , 2.5 mM CaCl_2 , and 20 mM sodium lactate, pH 6.8; gassed with 90% N_2 –10% CO_2) [19–21]. The method is convenient and severe enough to produce an H/R single cell model. Perfusion with buffer was controlled by gravity to maintain a flow rate of 6 ml/min.

2.6. Drugs

F_2 (synthesized by our lab and assayed by the Shanghai Organic Chemistry Institute of the Chinese Academy of Sciences; purity greater than 98%) was prepared as a 0.1 M stock solution in DMSO and diluted to the desired drug concentration with extracellular solution before each experiment. A DMSO of less than 0.1% did not affect the I_{NCX} and $I_{\text{Ca,L}}$ at the highest F_2 concentration used. Ouabain, nifedipine, CsCl, HEPES, and BAPTA were purchased from the Sigma Chemical Co., St. Louis, MO. All chemicals used were the highest grade available.

2.7. Analysis of statistics

All values presented are arithmetic means \pm SEM. Statistical significance was determined using a paired Students' t -test. Differences were considered significant when the P value was less than 0.05.

3. Results

3.1. F_2 reduces both outward and inward I_{NCX} under physiological conditions

Bi-directional I_{NCX} was induced by 1 mM Ca^{2+} and 140 mM Na^+ in the external solution and 20 mM Na^+ and 226 nM free Ca^{2+} in the pipette solution. Under these ionic conditions, following establishment of the whole-cell clamp configuration, the external solution was switched from the control external solution to Tyrode solution, while monitoring the increase in current until a steady state was reached. I_{NCX} was recorded under conditions in which we selectively blocked various ion channel currents, such as Na^+/K^+ pump currents, K^+ current, sarcoplasmic reticulum Ca^{2+} release channels, and Ca^{2+} currents. After recording the control current, 1.0 μM F_2 was added to the extracellular solution. Upon stabilization of current, 5 mM Ni^{2+} , a selective NCX inhibitor under these ionic conditions, was added to the extracellular solution to block I_{NCX} (Fig. 1B and C). F_2 inhibited outward I_{NCX} at $+60$ mV by $39.51 \pm 2.62\%$ ($n = 4$) and inward I_{NCX} at -150 mV by $10.68 \pm 0.62\%$ ($n = 4$).

3.2. F_2 inhibits I_{NCX} during H/R

To examine the effect of F_2 on I_{NCX} during H/R, current traces were recorded in the presence and absence of H/R exposure and with and without F_2 . The I – V relationship recorded in the presence

Download English Version:

<https://daneshyari.com/en/article/10761516>

Download Persian Version:

<https://daneshyari.com/article/10761516>

[Daneshyari.com](https://daneshyari.com)