



Removal of sphingomyelin headgroup inhibits the ligand binding function of hippocampal serotonin_{1A} receptors

Pushpendra Singh, Amitabha Chattopadhyay*

Centre for Cellular and Molecular Biology, Council of Scientific and Industrial Research, Uppal Road, Hyderabad 500 007, India

ARTICLE INFO

Article history:

Received 24 January 2012

Available online 10 February 2012

Keywords:

Serotonin_{1A} receptor

Sphingomyelinase

Sphingolipid

Ligand binding

Fluorescence anisotropy

ABSTRACT

Sphingolipids are essential components of eukaryotic cell membranes and are thought to be involved in a variety of cellular functions. Sphingomyelin is the most abundant sphingolipid in the nervous system. In this work, we explored the ligand binding function of the hippocampal serotonin_{1A} receptor upon hydrolyzing sphingomyelin to ceramide and phosphocholine using sphingomyelinase. The serotonin_{1A} receptor is an important neurotransmitter receptor and belongs to the superfamily of G-protein coupled receptors. It is involved in the generation and modulation of various cognitive, behavioral and developmental functions. We show here that specific agonist binding to serotonin_{1A} receptors in native hippocampal membranes is considerably reduced upon sphingomyelinase treatment. Interestingly, the overall membrane order does not exhibit any appreciable change under these conditions. Our results show the importance of sphingomyelin (specifically, the sphingomyelin headgroup) for the function of serotonin_{1A} receptors. These novel results constitute the first report on the effect of enzymatic hydrolysis of sphingomyelin on the ligand binding function of this important neurotransmitter receptor in native hippocampal membranes. Our results assume greater relevance in the broader perspective of the influence of the membrane lipid environment on the function of the serotonin_{1A} receptor in particular, and other G-protein coupled receptors in general.

© 2012 Elsevier Inc. All rights reserved.

1. Introduction

Sphingolipids are essential components of eukaryotic cell membranes and constitute ~10–20% of the total membrane lipids [1]. They represent diverse and dynamic regulators of a large number of cellular processes such as cell signaling, growth, differentiation, and neoplastic transformation. Sphingomyelin is the most abundant sphingolipid in the nervous system. For example, it makes up ~25% of total lipids in the myelin sheath [2]. Sphingolipids such as sphingomyelin are regarded as reservoirs for bioactive second messengers such as ceramide and sphingosine 1-phosphate [3,4]. The distribution of sphingomyelin in the cellular plasma membrane appears heterogeneous and patchy [5]. Importantly, sphingomyelins have been shown to be involved in the development and progression of several neurological diseases such as Alzheimer's disease [4,6] which could be due to impaired neurotransmission.

Abbreviations: 5-HT_{1A} receptor, 5-hydroxytryptamine-1A receptor; 8-OH-DPAT, 8-hydroxy-2-(di-*N*-propylamino)tetralin; BCA, bicinchoninic acid; DMPC, dimyristoyl-*sn*-glycero-3-phosphocholine; DPH, 1,6-diphenyl-1,3,5-hexatriene; GPCR, G-protein coupled receptor; PMSF, phenylmethylsulfonyl fluoride.

* Corresponding author. Fax: +91 40 2716 0311.

E-mail address: amit@ccmb.res.in (A. Chattopadhyay).

The G-protein coupled receptor (GPCR) superfamily is the largest and most diverse protein family in mammals, involved in signal transduction across membranes [7,8]. GPCRs regulate physiological responses to a diverse array of stimuli, and mediate multiple physiological processes. GPCRs have therefore emerged as major targets for the development of novel drug candidates in all clinical areas [9]. It is estimated that ~50% of clinically prescribed drugs act as either agonists or antagonists of GPCRs [10]. The serotonin_{1A} receptor is an important neurotransmitter receptor and belongs to the GPCR superfamily. It is the most extensively studied among the serotonin receptors [11]. The serotonin_{1A} receptor plays a key role in the generation and modulation of various cognitive, behavioral and developmental functions [11–13]. Agonists and antagonists of the receptor have been shown to possess potential therapeutic effects in anxiety or stress-related disorders [11]. The serotonin_{1A} receptor therefore serves as an important target in the development of therapeutic agents for neuropsychiatric disorders such as anxiety and depression. In the context of increasing pharmacological significance of the serotonin_{1A} receptor [14,15], its interaction with the surrounding lipids such as sphingomyelin assumes relevance. In this overall context and keeping in mind the relevance of sphingolipids in the nervous system [4], we explored the importance of the sphingomyelin headgroup on ligand binding function of the serotonin_{1A} receptor in native hippocampal membranes.

2. Materials and methods

2.1. Materials

Sphingomyelinase (EC 3.1.4.12) from *Bacillus cereus*, cholesterol, DMPC, DPH, EDTA, EGTA, MgCl_2 , MnCl_2 , iodoacetamide, PMSF, serotonin, sucrose, polyethylenimine, sodium azide, and Tris were obtained from Sigma Chemical Co. (St. Louis, MO). BCA reagent for protein estimation was from Pierce (Rockford, IL). $[^3\text{H}]8\text{-OH-DPAT}$ (sp. activity 106 Ci/mmol) was purchased from DuPont New England Nuclear (Boston, MA). GF/B glass microfiber filters were from Whatman International (Kent, UK). All solvents used were of analytical grade. Pre-coated silica gel 60 thin layer chromatography (TLC) plates were from Merck (Darmstadt, Germany). All other chemicals used were of the highest purity available. Water was purified through a Millipore (Bedford, MA) Milli-Q system and used throughout. Fresh bovine brains were obtained from a local slaughterhouse within 10 min of death and the hippocampal region was carefully dissected out. The hippocampi were immediately flash frozen in liquid nitrogen and stored at -70°C until further use.

2.2. Preparation of native hippocampal membranes

Native hippocampal membranes were prepared as described previously [16,17]. Native membranes were suspended in a minimum volume of 50 mM Tris, pH 7.4 buffer, homogenized using a hand-held Dounce homogenizer, flash frozen in liquid nitrogen and stored at -70°C . Protein concentration was assayed using the BCA reagent [18].

2.3. Treatment of native membranes with sphingomyelinase

Native membranes were resuspended in 50 mM Tris, pH 7.4 buffer at a protein concentration of 2 mg/ml and treated with sphingomyelinase (aliquoted from a stock solution of 200 U/ml in 10 mM Tris, pH 7.4 buffer) at 25°C with constant shaking for 1 h. Membranes were then spun down at $50,000\times g$ for 10 min at 4°C and resuspended in the same buffer.

2.4. Radioligand binding assays

Receptor binding assays were carried out as described earlier [17]. Tubes in duplicate with 0.5 mg protein in a total volume of 1 ml of buffer (50 mM Tris, 1 mM EDTA, 10 mM MgCl_2 , 5 mM MnCl_2 , pH 7.4) were incubated with the radiolabeled agonist $[^3\text{H}]8\text{-OH-DPAT}$ (final concentration in assay tube being 0.29 nM) for 1 h at 25°C . Nonspecific binding was determined by performing the assay in the presence of 10 μM serotonin. The binding reaction was terminated by rapid filtration under vacuum in a Millipore multiport filtration apparatus through Whatman GF/B 2.5 cm diameter glass microfiber filters (1.0 μm pore size), which were presoaked in 0.15% polyethylenimine for 1 h [19]. Filters were then washed three times with 3 ml of cold water (4°C), dried and the retained radioactivity was measured in a Packard Tri-Carb 1500 liquid scintillation counter using 5 ml of scintillation fluid.

2.5. Estimation of sphingomyelin by thin layer chromatography

Total lipids were extracted from control and sphingomyelinase-treated membranes according to Bligh and Dyer [20]. The lipid extracts were dried under a stream of nitrogen at 45°C . The dried extracts were dissolved in a chloroform/methanol mixture (1:1, v/v). Total lipid extracts were resolved by thin layer chromatography using a chloroform/methanol/water (65:25:4, v/v/v) as the solvent

system [21]. The separated lipids were visualized under ultraviolet light by spraying a fluorescent solution of 0.01% (w/v) primuline prepared in acetone [22]. A sphingomyelin standard was used to identify its position on TLC plates run with total lipid extracts obtained from control and sphingomyelinase-treated membranes. Sphingomyelin bands were scraped from TLC plates, lipids were re-extracted with a chloroform/methanol mixture (1:1, v/v) from samples, and the phosphate content was estimated and normalized to the phosphate content obtained from control samples.

2.6. Estimation of cholesterol content

Cholesterol content in membranes treated with different concentration of sphingomyelinase was estimated using the Amplex Red cholesterol assay kit [23].

2.7. Estimation of phospholipid content

The concentration of lipid phosphate was determined subsequent to total digestion by perchloric acid [24] using Na_2HPO_4 as standard. DMPC was used as an internal standard to assess lipid digestion. Samples without perchloric acid digestion produced negligible readings.

2.8. Fluorescence anisotropy measurements

Fluorescence anisotropy experiments were carried out using the fluorescent probe DPH as described previously [17,25]. Steady state fluorescence was measured in a Hitachi F-4010 spectrofluorometer using 1 cm path length quartz cuvettes at room temperature ($\sim 23^\circ\text{C}$). Excitation and emission wavelengths were set at 358 and 430 nm. Excitation and emission slits with nominal bandpasses of 1.5 and 20 nm were used. The optical density of the samples measured at 358 nm was 0.15 ± 0.01 . The anisotropy values remained identical even after dilution of membrane samples, indicating the absence of any scattering artifact [26]. Fluorescence anisotropy measurements were performed using a Hitachi polarization accessory. Anisotropy (r) values were calculated from the equation [27]:

$$r = \frac{I_{VV} - GI_{VH}}{I_{VV} + 2GI_{VH}}$$

where I_{VV} and I_{VH} are the measured fluorescence intensities (after appropriate background subtraction) with the excitation polarizer vertically oriented and the emission polarizer vertically and horizontally oriented, respectively. G is the grating correction factor and is the ratio of the efficiencies of the detection system for vertically and horizontally polarized light and is equal to I_{HV}/I_{HH} . All experiments were done with multiple sets of samples and average values of fluorescence anisotropy are shown in Fig. 4.

2.9. Statistical analysis

Significance levels were estimated using Student's two-tailed unpaired t -test using Graphpad Prism software version 4.0 (San Diego, CA).

3. Results

Sphingomyelinases are water soluble enzymes that act at the membrane interface and specifically hydrolyze sphingomyelin into ceramide and phosphocholine ([28,29]; see Fig. 1). Phosphocholine, a water soluble moiety, is liberated from sphingomyelin leaving a hydrophobic ceramide backbone of sphingomyelin in the

Download English Version:

<https://daneshyari.com/en/article/10762250>

Download Persian Version:

<https://daneshyari.com/article/10762250>

[Daneshyari.com](https://daneshyari.com)