

Reduced expression of the E-cadherin gene and its aberrant DNA methylation in hamster pancreatic tumors[☆]

Kyoko Shimizu^a, Madoka Hanaoka^a, Ayumi Kato^a, Hiromasa Fujii^b,
Kanya Honoki^b, Toshifumi Tsujiuchi^{a,*}

^a *Laboratory of Cancer Biology and Bioinformatics, Department of Life Science, Faculty of Science and Engineering, Kinki University, 3-4-1, Kowakae, Higashiosaka, Osaka 577-8502, Japan*

^b *Department of Orthopedic Surgery, Nara Medical University, 840 Shijo-cho, Kashihara, Nara 634-8521, Japan*

Received 4 August 2005

Available online 18 August 2005

Abstract

The expression of the E-cadherin gene and its DNA methylation status in the 5' upstream region were investigated in pancreatic duct adenocarcinomas (PDAs) induced by *N*-nitrosobis(2-oxopropyl)amine (BOP) in hamsters. Female Syrian golden hamsters received 70 mg/kg BOP, followed by repeated exposure to an augmentation pressure regimen consisting of a choline-deficient diet combined with DL-ethionine then L-methionine and a further administration of 20 mg/kg BOP. A total of 15 PDAs were obtained, along with total RNA for assessment of expression by real-time quantitative reverse transcription-polymerase chain reaction. The expression of the E-cadherin was significantly reduced in PDAs ($p < 0.05$) compared with normal pancreatic tissue. For the analysis of methylation status, bisulfite sequencing was performed with two normal pancreatic tissues and six tumors. The normal pancreatic tissue was all demethylated in this region of E-cadherin. In contrast, six PDAs were highly methylated, correlating with reduced expression of the E-cadherin. These results suggest that aberrant DNA methylation of the E-cadherin gene may play a role in the development of PDAs induced by BOP in hamsters.

© 2005 Elsevier Inc. All rights reserved.

Keywords: Pancreatic duct adenocarcinoma; DNA methylation; E-cadherin; Hamster; Nitrosamine

E-cadherin is a transmembrane glycoprotein that mediates cell–cell adhesion in epithelial cells [1]. Reduced expression of E-cadherin is found in tumor progression and metastasis, while abnormal expression of E-cadherin is involved in tumor invasion and metastasis [2–6]. Since methylation of cytosine residues at CpG dinucleotides can suppress gene expression in mammalian genomes [7–11], it has been suggested that aberrant

DNA methylation of the promoter regions of E-cadherin may be the major mechanisms of gene silencing in several tumors, including pancreatic cancers [12–14].

Pancreatic duct adenocarcinoma (PDA) has one of the lowest cure rates of human malignancies [15,16]. In order to understand the underlying mechanisms, we have generated a model for the development of hamster PDAs by *N*-nitrosobis(2-oxopropyl)amine (BOP), mirroring the human situation, and this model gives us the ability to gain a better understanding of the underlying mechanisms [17–19]. Indeed, we have reported several genetic changes in this pancreatic carcinogenesis model. For example, Ki-ras mutations were frequently found in preneoplastic lesions as well as PDAs, while p53 mutations were detected as late events [20–22].

[☆] *Abbreviations:* PDA, pancreatic duct adenocarcinoma; BOP, *N*-nitrosobis(2-oxopropyl)amine; PCR, polymerase chain reaction; RT, reverse transcription; GAPDH, glyceraldehyde-3-phosphate dehydrogenase.

* Corresponding author. Fax: +81 6 6721 2721.

E-mail address: tsujiuch@life.kindai.ac.jp (T. Tsujiuchi).

In addition, we have provided evidence that alteration of the *Fhit* gene may also be involved in the development of PDAs [23]. In contrast, the analysis for epigenetic changes, such as DNA methylation status in the promoter region of genes, has been limited, since only a small amount of genetic information is available for hamsters.

Recently, we determined the 5' upstream region of the hamster *p16* gene using the suppression PCR method combined with gene-specific primers and measured its methylation patterns in PDAs [24]. In the present study, we determined the 5' upstream region of the hamster E-cadherin gene with this method, and to better understand epigenetic changes in pancreatic carcinogenesis expression of the E-cadherin gene and its DNA, methylation status in hamster PDAs induced by BOP was examined.

Materials and methods

Production of hamster PDAs. A total of 18 female Syrian golden hamsters, weighing approximately 100 g each, were used (Japan SLC). PDAs were induced in 15 animals according to the rapid production model [17–19]. Briefly, BOP (30 mg/kg body weight) (Nakalai Tesque, Kyoto, Japan) was given subcutaneously as initiation, followed by two cycles of augmentation pressure which consisted of choline-deficient diet administration and ethionine–methionine–BOP injection [17–19]. To obtain normal control tissues including the pancreas, the remaining three animals were maintained free from carcinogen exposure throughout the experimental period. All hamsters were sacrificed under light ether anesthesia at 10 week after the beginning of the experiment and the pancreas was immediately excised. Macroscopically apparent tumors were dissected from the surrounding tissue and frozen in liquid nitrogen.

Real-time quantitative RT-PCR. Total RNA was extracted from frozen tissue using ISOGEN (Nippon Gene, Toyama, Japan) and first-strand cDNA was synthesized from 0.2 µg samples with Ready-To-Go Your-Prime First-Strand Beads (Pharmacia, Tokyo, Japan). To eliminate possible false positives caused by residual genomic DNA, all samples were treated with DNase.

Real-time quantitative RT-PCR analysis using a Smart Cycler II System (TaKaRa Bio, Shiga, Japan) and a SYBR Premix Ex Taq (TaKaRa) was performed according to the manufacturer's protocol. One microliter of the synthesized cDNA was used in the following assay. The primer sets used in this assay were as follows: for E-cadherin, 5'-CTGCAGGTCTCATCATGGC-3' (sense) and 5'-ACCTGTAGACCTCGGCACTG-3' (antisense) and for glyceraldehyde-3-phosphate dehydrogenase (*Gapdh*), 5'-TTGTGAAGGTCGGTGTGAAC-3' (sense) and 5'-AGGGGTCGTTGATGGCAACA-3' (antisense). The PCR conditions were as follows: 95 °C for 20 s, 45 cycles of 95 °C for 15 s and 64 °C (for E-cadherin) or 55 °C (for *Gapdh*) for 30 s. To obtain the standard curve for each gene, a cDNA synthesized from normal lung tissue was used. The amplification plots of the PCR were used to determine the threshold cycle (C_t). The C_t value represented the first cycle in which there was a significant increase in fluorescence above the background fluorescence. The initial copy number of the target mRNA was calculated by a plot of the C_t against the input target quantity. The data of the target genes were normalized to *Gapdh*. Each assay was repeated at least twice for confirmation. Data were statistically analyzed by Student's *t* test.

Determination of 5' upstream region of hamster E-cadherin gene. To determine the nucleotide sequence of the 5' upstream region of the

hamster E-cadherin gene, the suppression PCR combined with gene-specific primers was performed as described recently [24]. Genomic DNA was extracted from the liver of a female Syrian golden hamster (Japan SLC, Shizuoka, Japan) using a DNeasy tissue kit (Qiagen, Hilden, Germany). Next, 5 µg of the genomic DNA was digested with *RsaI* (New England Biolabs, MA, USA). After ethanol precipitation, the digested DNA fragments were ligated with 2 µl of a specific adaptor primer at the *RsaI*-digested end by T4 ligase (TaKaRa) in a total volume of 30 µl at 16 °C. The adaptor primer sequence was as follows: 5'-GTAATACGACTCACTATAGGGCTCGAGCGGCCGCCC GGCCAGGT-3' and 5'-ACCTGCC-3' [24,25]. The primary PCR was performed with the first adaptor primer P1, 5'-GTAATACGACTCACTATAGGGC-3' [24,25], and the first gene-specific primer R1, 5'-CTGCAGCAGGAACAGGATCG-3'. The conditions for the suppression PCR were as previously described [24]. For the second nested PCR amplification, the first PCR product was diluted 20-fold with distilled water and amplified with the second adaptor primer P2, 5'-TGTAGCGTGAAGACGACAGAA-3' [24,25] and the second gene-specific primer R2, 5'-AGCAGGAACAGGATCGCGGA-3'. Both gene-specific primers were designed from rat E-cadherin cDNA sequences (GenBank Accession No. AB017696). The PCR amplification was carried out in a total volume of 20 µl containing 1 µl of each gene primer, 200 µM of each dNTP, 1× PCR buffer (Perkin-Elmer, Applied Biosystems Division, Foster City, CA, USA), and 0.5 U of AmpliTaq Gold (Perkin-Elmer) under the following reaction conditions; primary denaturation for 10 min at 95 °C, 35 cycles of 30 s denaturation at 95 °C, 15 s annealing at 69 °C, and 1 min extension at 72 °C, and a final extension for 10 min at 72 °C. The amplified product was separated in a 1% agarose gel containing 0.05 µg/ml ethidium bromide, extracted from the gel, subcloned using a TOPO TA cloning kit (Invitrogen, CA, USA), and sequenced with a BigDye terminator v3.0 cycle sequencing ready reaction kit (Applied Biosystems Japan, Tokyo, Japan) and an ABI PRISM 310 genetic analyzer (Applied Biosystems Japan).

Bisulfite sequencing. Bisulfite treatment of genomic DNA was performed as previously described [24,26]. Briefly, genomic DNA was extracted with DNeasy tissue kit (Qiagen) from frozen tissues, and 500 ng of each sample was digested with an appropriate restriction enzyme. The digested DNA was denatured in 0.3 N NaOH, and then 2.9 M sodium bisulfite (Sigma, St. Louis, MO, USA) and 0.5 mM hydroquinone (Sigma) were added and the mixture underwent 15 cycles of 30 s denaturation at 95 °C and 15 min incubation at 50 °C. The sample was then desalted with the Wizard DNA cleanup system (Promega, Madison, WI, USA), and desulfonated by treatment with 0.3 N NaOH at room temperature for 5 min. After ethanol precipitation with ammonium acetate, DNA was dissolved in distilled water.

For bisulfite sequencing, PCR was performed with the primer sets as follows; BS-F: 5'-ATTTAGATGAAGAGTAAAGTTTTTTGTA-3', BS-R: 5'-ACACAAAATCCATAACTAACC-3' (annealing temperature: 57 °C). PCR products were subcloned with a TOPO TA cloning kit (Invitrogen) and sequenced with a BigDye terminator v3.0 cycle sequencing ready reaction kit (Applied Biosystems Japan) and ABI PRISM 310 genetic analyzer (Applied Biosystems Japan). For each sample, ten clones were sequenced.

Results and discussion

It has been reported that loss or reduced expression of E-cadherin was detected in several human malignancies [2–6]. In pancreatic tumors, reduced expression of E-cadherin was found in human primary pancreatic carcinomas, correlating with an invasion and metastasis of tumor cells [5,6]. Therefore, it has been suggested that loss or reduced expression of E-cadherin is involved in

Download English Version:

<https://daneshyari.com/en/article/10768250>

Download Persian Version:

<https://daneshyari.com/article/10768250>

[Daneshyari.com](https://daneshyari.com)