

Identification of tick periviscerokinin, the first neurohormone of Ixodidae: Single cell analysis by means of MALDI-TOF/TOF mass spectrometry

Susanne Neupert^{a,b,*}, Reinhard Predel^{a,b}, William K. Russell^c, Ronald Davies^d,
Patricia V. Pietrantonio^e, Ronald J. Nachman^b

^a Saxon Academy of Sciences, Research Group Jena, Erbertstrasse 1, 07743 Jena, Germany

^b Areawide Pest Management Research, Southern Plains Agricultural Research Center, USDA, 2881 F/B Road, College Station, TX 77845, USA

^c The Laboratory for Biological Mass Spectrometry, Department of Chemistry, Texas A&M University, College Station, TX 77843, USA

^d Cattle Fever Tick Research Laboratory, USDA-ARS, 22675 N. Moorefield Road, Edinburg, TX 78541, USA

^e Department of Entomology, Texas A&M University, College Station, TX 77843, USA

Received 20 October 2005

Available online 9 November 2005

Abstract

The first peptidergic neurohormone from the ticks *Ixodes ricinus* and *Boophilus microplus* has been identified by using a combination of immunocytochemistry and mass spectrometric analysis of single cells. The novel peptide (Ixori-PVK, PALIPFPRV-NH₂) shows a high sequence homology with members of the insect periviscerokinin/CAP2_b peptides that in insects are involved in the regulation of water balance. The function of this peptide in ticks is still unknown, but these pests consume large amounts of blood in a single blood meal which is a challenge for the regulation of diuretic processes. Thus, the novel peptide may be involved in one of the key physiological processes in ticks. High energy collision-induced dissociation was successfully used to distinguish between Leu/Ile ambiguities in single cell preparations. This is the first successful de novo sequencing of a peptide from single cell preparations of arthropods.

© 2005 Elsevier Inc. All rights reserved.

Keywords: MALDI-TOF/TOF mass spectrometry; Neuropeptide; Periviscerokinin; *Boophilus*; *Ixodes*; Tick; Single cell analysis

Members of the family Ixodidae, commonly known as ticks, cause great economic losses of livestock; an estimated damage of several million dollars per year is reported for the United States [1]. As obligatory blood feeders, ticks consume large amounts of blood in a single blood meal which is a challenge for the regulation of diuretic processes. The necessity to remove excessive water after a blood meal has also been observed in a number of insects such as mosquitos [2–4], blood-feeding bugs (e.g., *Rhodnius prolixus*) [5,6], fleas [7], and lice [8]. In the insects, diuresis is known to be regulated by several neuropeptides which influence secretion and/or reabsorption processes in the Malpighian tubules and the rectum [9]. Among these neuropeptides, are

the insect kinins and perviscerokinin/CAP2_b-related peptides which may occur with multiple forms in a single species [10–14]. However, nothing is known about the hormonal control of excretion in ticks and no neuropeptides have been unambiguously identified from any tick so far. Recently, a receptor with sequence similarity to an insect kinin receptor (*Drosophila melanogaster*, CG 10626) [15] has been identified from the Southern Cattle Fever tick *Boophilus microplus* (CG10626) [16] and shown to functionally respond to various kinin peptides [17].

In this study, we searched for peptidergic neurohormones that may influence the water balance of ticks by screening the nervous system for peptide homologs to the insect kinins and periviscerokinin/CAP2_b-related peptides. Antisera against these neuropeptides revealed the expression of immunoreactive substances in a few neurons. We

* Corresponding author. Fax: +49 3641 949192.

E-mail address: mail@susanne-neupert.de (S. Neupert).

used the advantages of MALDI-TOF/TOF mass spectrometry to perform an analysis of those neurons that were found to contain immunoreactive material. This strategy resulted in the identification of a tick neuropeptide, which is a member of the periviserokinins, a peptide family that had been found only in insects so far. High energy collision-induced dissociation was used to distinguish between Leu/Ile in single cell preparations. As shown in our study, the novel peptide is expressed in another tick as well.

Materials and methods

Animals. Southern Cattle Fever ticks (*B. microplus*) were maintained at the Cattle Fever Tick Research Laboratory (USDA-ARS, Mission, TX), where all preparations on this pest were performed as well. The *Ixodes ricinus* was caught in Jena (Germany) by sweeping a net in grass fields. Only unfed adult ticks were used throughout the experiments.

Dissection and sample preparation for mass spectrometry. Isolated central nervous system tissue of adult ticks was fixed with microneedles in a chamber containing insect saline. A number of putative neurosecretory neurons could be identified due to their slight bluish color (Tyndall effect). After selection of the neurons of interest, the ganglionic sheath was partially removed and the respective neurons were individualized. Without any enzyme treatment, cells were removed step by step using an uncoated glass capillary and transferred to a stainless steel sample plate for MALDI-TOF mass spectrometry. Subsequent sample preparation was performed as described in Neupert and Predel [18].

MALDI-TOF/TOF tandem mass spectrometry. MALDI analysis was performed on the ABI 4700 proteomics analyzer (Applied Biosystems, Framingham, MA). Due to the nature of the samples all acquisitions were taken in manual mode. Initially the instrument was operated in reflectron mode, in order to determine the parent masses. A laser intensity of 3800 was typically employed for ionizing the neuropeptides. For the tandem MS experiments, the CID acceleration was 1 kV in all cases. The number of laser shots used to obtain a spectrum varied from 500 to 5000, depending on signal quality. In order to change the net amount of activation energy imparted to the primary ions, the collision gas (atmospheric air) pressure was increased. Three gas pressures were employed: none, medium, and high. The fragmentation patterns from these three different settings were used to determine the sequence of the peptide. The fragmentation data obtained in these experiments was handled using the Data ExplorerT software package.

Immunocytochemistry. Dissected nervous system was fixed for 24 h at 4 °C with 4% formaldehyde in phosphate-buffered saline (PBS), pH 7.2. Subsequently, preparations were washed in PBS–4% Triton X-100 and PBS–1% Triton X-100 for 24 h, respectively. Then, the preparations were incubated for 3 days at 4 °C in anti-Pea-PVK-2 serum [19] (1:4000) diluted with PBS–1% Triton X-100 containing 0.25% bovine serum albumin and normal goat serum. Following washing in 0.1 mol/L Tris–HCl, 3% NaCl, and 1% Triton X-100 (pH 7.6) for 24 h, the Fluorochrome-labelled (Cy3; Amersham, Braunschweig, Germany) secondary antibodies were directly used as a mixture in PBS–bovine serum albumin (2.5 mg/ml) at a concentration of 1:3000 for 3 days. The preparations were washed again for 24 h in 0.1 mol/L Tris–HCl, 3% NaCl, 1% Triton X-100 (pH 7.6), and then incubated at 4 °C in anti-leucokinin-1 (DPAFNSWGa) serum (1:4000) diluted with PBS–1% Triton X-100 containing 0.25% bovine serum albumin and normal goat serum. After 3 days, the samples were washed (24 h) in 0.1 mol/L Tris–HCl, 3% NaCl, and 1% Triton X-100 (pH 7.6), the Fluorochrome-labelled secondary antibodies Cy2 were directly used as a mixture in PBS–bovine serum albumin (2.5 mg/ml) at a concentration of 1:3000 for 3 days. Finally, the preparations were washed again 24 h in 0.1 mol/L Tris–HCl, 3% NaCl, and 1% Triton X-100 (pH 7.6) and transferred to glycerol. For visualization, tissues were dehydrated in ethanol, cleared in methyl salicylate, and mounted in entellan (Euromex microscopes Holland).

Documentation. Immunostainings were examined with a confocal laser scanning microscope (ZEISS LSM 510 Meta system, Jena, Germany), equipped with a Helium/Neon1 laser (wavelength 543 nm) and an Argon2 laser (wavelength 458, 477, 488, and 514 nm). Serial optical sections were assembled into combined images. Images were exported and processed with Adobe Photoshop 7.0 software.

Peptide synthesis. Ixori-PVK (PALIPFPRV-NH₂) was synthesized via Fmoc methodology on Rink Amide resin (Novabiochem, San Diego, CA) using Fmoc protected amino acids (Advanced Chemtech, Louisville, KY) on an ABI 433A peptide synthesizer (Applied Biosystems, Foster City, CA) under previously described conditions [20]. The crude product was purified on a Waters C₁₈ Sep Pak cartridge and a Delta-Pak C₁₈ reverse-phase column (8 × 100 mm, 15 μm particle size, and 100 Å pore size) on a Waters 510 HPLC controlled with a Millennium 2010 chromatography manager system (Waters, Milford, MA) with detection at 214 nm at ambient temperature. Solvent A = 0.1% aqueous trifluoroacetic acid (TFA); solvent B = 80% aqueous acetonitrile containing 0.1% TFA. Conditions: initial solvent consisting of 20% B was followed by the Waters linear program to 100% B over 40 min; flow rate, 2 ml/min. Delta-Pak C-18 retention time: Ixori-PVK (PALIPFPRV-NH₂), 9.0 min. The peptide was further purified on a Waters Protein Pak I125 column (7.8 × 300 mm) (Milligen, Milford, MA). Conditions: flow rate: 2.0 ml/min; isocratic with solvent = 80% acetonitrile made to 0.01% TFA. WatPro retention time: Ixori-PVK (PALIPFPRV-NH₂), 6.0 min. Amino acid analysis was carried out under previously reported conditions [20] and used to quantify the peptide and to confirm identity, leading to the following analysis: Ixori-PVK (PALIPFPRV-NH₂): A[1.0], F[1.0], I[1.0], L[1.0], P[2.6], R[1.0], V[1.0]. The identity of the peptide analog was confirmed via MALDI-TOF MS on a Kratos Kompact Probe MALDI-TOF MS machine (Kratos Analytical, Manchester, UK) with the presence of the following molecular ion [M+H⁺]: Ixori-PVK (PALIPFPRV-NH₂) 1008.63 Da [M+H⁺].

Results and discussion

The search for peptidergic hormones in arthropods as tiny as ticks is generally hampered by the small size of the central nervous system that contains low amounts of bioactive substances. In addition, working on ticks provides the researcher with further problems. First of all, and despite their great economic importance, not a single peptide hormone is known from ticks and bioassays to perform conventional neuropeptides purification strategies are not available. This paucity of information was confirmed by a current search for tick neuropeptides in GenBank that only retrieved sequences for two putative salivary secreted neuropeptide-like proteins from the tick *Ixodes pacificus*. Second, the synganglionic tick nervous system does not exhibit well-separated neurohemal release sites as observed in insects, where neurohormones are accumulated and easily accessible for structural elucidation [12]. Having in mind the positive experiences with the analysis of single identified neurons in an insect [18], we decided to perform the following strategy for the identification of putative diuretic hormones. Antisera raised against insect diuretic peptides, namely leucokinin-1 [21] and Pea-periviscerokinin-2 [22], were used to locate neurons in the CNS of *I. ricinus* which may express peptides that are sequence related to the insect kinins and periviscerokinins. Indeed, this procedure yielded several immunoreactive cells, a majority of these neurons were immunostained by both antisera (Fig. 1). Due to their slightly bluish color (Tyndall effect), we succeeded in detecting a number of these neurosecretory neurons in

Download English Version:

<https://daneshyari.com/en/article/10768669>

Download Persian Version:

<https://daneshyari.com/article/10768669>

[Daneshyari.com](https://daneshyari.com)