

Effect of thymosin β 15 on the branching of developing neurons

Jeehyung Choe^a, Woong Sun^a, Seung-Yong Yoon^b, Im Joo Rhyu^a,
Eun Hae Kim^a, Hyun Kim^{a,*}

^a Department of Anatomy and Division of Brain Korea 21 Biomedical Science, Korea University College of Medicine,
126-1, 5-Ka, Anam-Dong, Sungbuk-Gu, Seoul 136-705, Republic of Korea

^b Department of anatomy and Cell Biology, Ulsan University, 388-1 Pungnap-dong, Songpa-gu, Seoul 138-736, Republic of Korea

Received 14 March 2005

Available online 30 March 2005

Abstract

The thymosin β s (T β s) are polypeptide regulators of actin dynamics that are critical for the growth and branching of neurites in developing neurons. We found that mRNAs for T β 4, T β 10, and T β 15 were highly expressed in the developing rat brain during neurogenesis, supporting a role for the T β s in this process. Overexpression of the T β s increased the number of neurite branches per neuron in cultured hippocampal and cerebral cortex neurons, and T β 15 had the greatest effect. Actin binding activity appears to be essential for the branch-promoting activity of T β s because two mutants of T β 15 lacking monomeric actin binding activity failed to stimulate branch formation. We also found that transfection of siRNA against T β 15 reduced branching. Taken together, these data suggest that the three T β s, and especially T β 15, stimulate neurite branching during brain development.

© 2005 Elsevier Inc. All rights reserved.

Keywords: Thymosin β ; Neurogenesis; Branching; Actin dynamics; Actin binding protein

Neurogenesis is essential in development to allow neurons to make several thousand different connections with their partners in different brain regions [1–3]. Neuronal development is initiated upon activation of membrane receptors by extracellular cues [4], and this triggers intracellular cascades some of which alter the organization of the actin cytoskeleton [5]. As a result, the original rounded shape of the cells is broken down giving rise to buds, which become neurites and are later transformed into axons and dendrites [1].

The motility of growth cones, the main determinant of neurite elongation and branching [6–8], is thus largely dependent on the reorganization of the actin cytoskeleton [9], and actin dynamics are believed to be critical for neurite growth, branching, and pathfinding [7,10]. Branching occurs via changes of actin and microtubule structure in primary growth cones or axon shafts [11],

and numerous actin-associated proteins that control actin organization are involved in neurite development [1,12,13].

The thymosin β s (T β s) have G-actin sequestering activity and are key regulators of actin dynamics. They are small (5 kDa), highly conserved proteins [14], and at least three different isoforms (T β 4, T β 10, and T β 15) have been identified in mammals [15,16]. T β 4 and T β 10 are abundant in neural tissue as well as in circulating cells such as platelets, leukocytes, and macrophages [15,17–19]. On the other hand, the expression of T β 15 has not been thoroughly examined. Although the function of T β s in the mammalian brain has yet to be elucidated, it is believed that they play a role in neurite development [15,19]. During zebrafish development, their expression is closely correlated with neuronal growth and differentiation, and inhibition of T β expression impairs axonal tract formation [15].

In this study, we examined the expression of T β s in the developing rat brain, and assessed their function in

* Corresponding author. Fax: +82 2 929 5696.

E-mail address: kimhyun@korea.ac.kr (H. Kim).

neurite development by overexpressing and knocking down a number of T β isoforms *in vitro*. Our data indicate that T β s, especially T β 15, are involved in the regulation of axonal branches during brain development.

Materials and methods

Construction of expression vectors. PCR-generated DNA fragments containing the full coding regions of various T β s were ligated in-frame into the *Bam*HI and *Hind*III site of the N-terminally tagged expression vector pYFP-C1 (Clontech). T β 15-T12P and -L18A were generated with a QuickChange site-directed mutagenesis kit (Stratagene, La Jolla, CA). Western blotting confirmed that the YFP-tagged T β s were properly expressed (data not shown). To construct the siT β 15 vector, pSilencer.U6.1.0 (Ambion) was digested with *Apa*I and *Eco*RI, and blunt-end ligated with insert fragments. The target sequence of siT β 15, derived from the NCBI database (Accession No.: U25684), was 5'-AGAATACTCTTCCTTCGAA-3'. A BLAST search revealed no significant sequence homology of this sequence to other genes. Sequences of 19–23 nt separated by a 9 nt hairpin loop from the reverse complementary repeat, with five thymidines as termination signal, were synthesized by Genotech (Korea) and inserted into pSilencer.U6.1.0.

Cell culture and transfection. Cultures were prepared from the cerebral cortex and hippocampus of gestation day 17 Sprague–Dawley rat embryos (E17) [20]. Briefly, cortical and hippocampal cells were isolated by trypsinization and cultured in serum-free conditions (Neurobasal media + B27 supplements) at a density of 5×10^4 cells/well in 24-well plates. Cultures were maintained in a 37 °C humidified atmosphere of 5% CO₂ for up to 2 days and then transfected with 1 μ g of the various T β constructs using Lipofectamine 2000 (Invitrogen).

To examine the effect of T β 15 on F-actin structure, COS-7 cells were cultured in DMEM supplemented with 10% fetal bovine serum (FBS) and antibiotics. One day before transfection, the cells were seeded in 24-well plates, and transfection was performed with a CalPhos Mammalian Transfection Kit (BD Bioscience).

Immunocytochemistry. Cells were fixed for 15 min with 4% paraformaldehyde in PBS. After blocking in 3% BSA and 0.2% Triton X-100 in PBS, primary antibodies (rabbit-anti-T β 4, BIODESIGN International; rabbit-anti-T β 10, BIODESIGN international; or chicken-anti-T β 15 kindly provided by Dr. B.R. Zetter, Harvard University, 1:500 with mouse-anti-GAP43 (Sigma) 1:500, or mouse-anti- β -tubulin isotype III (Sigma) 1:1000) were applied to the slides. The slides were then rinsed with PBS and incubated in secondary antibody (goat-anti-rabbit (Vector Laboratories, Burlingame, CA) 1:500, or goat-anti-chicken (Amersham) 1:500 with goat anti-mouse (Vector Laboratories, Burlingame, CA) or TRITC labeled phalloidin (Sigma) 1:500) in 3% BSA and 0.2% Triton X-100 in PBS for 30 min at RT. Slides were rinsed with PBS and mounted.

In situ hybridization. In situ hybridization was performed as described previously [21,22]. Frozen sections (12 μ m thick) were thaw-mounted, fixed in 4% paraformaldehyde, and incubated with 0.25% acetic anhydride in 0.1 M triethanolamine/0.9% NaCl (pH 8.0) for 10 min. They were then dehydrated/delipidated in ethanol and chloroform, and finally air-dried. Hybridization probes were prepared from pGEM-T Easy plasmids containing nt 1–499 of T β 4 (AA819609), nt 32–445 of rat T β 10 (M58405.1), and nt 1–437 of T β 15 (AA997865) by transcription with appropriate RNA polymerases using a Riboprobe System (Promega, Madison, WI) and ³⁵S-UTP (Amersham Pharmacia Biotech). Sections were hybridized with ³⁵S-labeled probes (1.2 \times 10⁶ cpm/slide) at 52 °C overnight, followed by four washes in 4 \times SSC at RT. The slides were then incubated with RNase buffer [RNase A (10 mg/ml in DW), 0.5 M NaCl, 10 mM Tris–HCl, and 1 mM EDTA, pH 8.0] at 37 °C for 30 min, washed with 2 \times SSC/10 mM DTT (twice for 5 min), 1 \times SSC/10 mM DTT for 10 min, 0.5 \times SSC/10 mM DTT for 10 min, and 0.1 \times SSC/10 mM DTT for 30 min at 62 °C. Fi-

nally, the sections were dehydrated, air-dried, and exposed to X-ray film (BioMAX-MR; Kodak) for 5 days. All experimental procedures using animals were in accordance with the NIH *Guide for the Care and Use of Laboratory Animals* and were approved by the Committee at Korea University College of Medicine.

Quantitative analysis of neurites. Stained and transfected cells were examined with a Zeiss Axioskop2 microscope equipped with a digital camera (CoolSNAP, Photometrics) using metaview software, or a Zeiss Axiovert-equipped digital camera (CoolSNAP, Photometrics) with IPlab3.0 software. The total number of neurites was determined for each cell. Neurons that were confounded by unresolvable processes were not included. All data were analyzed by one-way ANOVA and Student's *t* test, with *P* < 0.05 as the accepted level of statistical significance.

Results

Thymosin β s are strongly expressed in developing neuronal processes

We examined the expression of three T β s (T β 4, 10, and 15) in the developing rat brain by *in situ* hybridization. At E18, expression of T β 15 was predominantly observed in the central nervous system (CNS), whereas T β 4 and T β 10 were found throughout the embryo (Figs. 1A–C). There were differences in mRNA signal intensities (T β 4 > T β 10 > T β 15), consistent with previous reports [23,24]. These differences in expression level were maintained on P3 (Figs. 1D–F). In the adult, T β 4 expression was substantially reduced but a moderate level remained in most brain regions (Fig. 1G). Expression of T β 10 was restricted to particular areas including olfactory bulb, deep layer of cerebral cortex, and hippocampal formation (Fig. 1H). At the same time a low level of T β 15 expression was observed in the olfactory bulb (Fig. 1I).

Immunocytochemical analyses of cultured primary neurons demonstrated that the T β s were enriched in developing neurite processes and growth cones (Figs. 1J–U). For instance, T β 15 immunoreactivity (IR) colocalized with GAP43 (a marker for growth cones), phalloidin (a marker for F-actin), and Tuj1 (a marker for neurites). Similar cellular distributions of T β 4 and T β 10 were observed (data not shown).

Overexpression of the T β s increases neurite branching in primary cultured neurons

We tested whether overexpression of T β s modified neurite development. Overexpression of YFP-tagged T β 15 in DIV2 cerebral cortex neurons increased the number of neurite branches (Fig. 2A). Although overexpression of YFP-tagged T β 4 and T β 10 appeared to have similar branch-promoting effects, these effects did not reach statistical significance. To rule out a cell type-specific effect of T β 15, we tested whether overexpression of T β 15 had the same effect on cultured hippocampal

Download English Version:

<https://daneshyari.com/en/article/10771319>

Download Persian Version:

<https://daneshyari.com/article/10771319>

[Daneshyari.com](https://daneshyari.com)