

Molecular identification of a myosuppressin receptor from the malaria mosquito *Anopheles gambiae*[☆]

Susanne Schöller, Martin Belmont, Giuseppe Cazzamali, Frank Hauser, Michael Williamson, Cornelis J.P. Grimmelikhuijzen*

Department of Cell Biology and Comparative Zoology, Institute of Biology, University of Copenhagen, Universitetsparken 15, DK-2100 Copenhagen, Denmark

Received 23 November 2004
Available online 8 December 2004

Abstract

The insect myosuppressins (X₁DVX₂HX₃FLRFamide) are neuropeptides that generally block insect muscle activities. We have used the genomic sequence information from the malaria mosquito *Anopheles gambiae* Genome Project to clone a G protein-coupled receptor that was closely related to the two previously cloned and characterized myosuppressin receptors from *Drosophila* [Proc. Natl. Acad. Sci. USA 100 (2003) 9808]. The mosquito receptor cDNA was expressed in Chinese hamster ovary cells and was found to be activated by low concentrations of *Anopheles* myosuppressin (TDVDHVFLRFamide; EC₅₀, 1.6 × 10⁻⁸ M). The receptor was not activated by a library of 35 other insect neuropeptides and monoamines, including neuropeptides that resembled myosuppressin in their C-terminal moiety, such as PDRNFLRFamide (*Anopheles* FMRFamide-3), other *Anopheles* FMRFamide peptides, or neuropeptide F-like peptides, showing that the receptor was quite selective for myosuppressin. These results also showed that the myosuppressin receptor needs a much larger portion than the C-terminal FLRFamide sequence for its activation. The insect myosuppressins are often grouped together with the insect FMRFamides under the name FaRPs (FMRFamide-related peptides). However, this is not justified anymore, because the insect myosuppressin receptor/ligand couple is both functionally and evolutionarily fully unrelated to the insect FMRFamide receptor/ligand couple. To our knowledge, this is the first report on the molecular identification of a mosquito neuropeptide receptor.

© 2004 Elsevier Inc. All rights reserved.

Keywords: G protein-coupled receptor; Neuropeptide; FMRFamide; Myosuppressin; Leucomyosuppressin; *Anopheles*; Mosquito; Malaria

The insect myosuppressins are neuropeptides with the C-terminal sequence X₁DVX₂HX₃FLRFamide (where X₁ is pQ, P, T; X₂ is D, G, V; and X₃ is V, S) [1]. The insect myosuppressins obtained their name because they inhibit most insect visceral muscles, including those that are involved in the passage of food along the insect alimentary canal [1–5]. The myosuppressins have been iso-

lated from a variety of insects and it can be assumed that they occur in all insect species [1–4,6–10].

The C-terminal four amino acid moiety of the insect myosuppressins (FLRFamide) resembles that of the FMRFamide neuropeptides. However, the actions of myosuppressins are often different from those of the insect FMRFamides [1,11] and also their preprohormones have a quite different organization (one immature myosuppressin sequence is contained in the myosuppressin preprohormones, while typically a large number of FMRFamide sequences—up to 24—are contained in the insect FMRFamide preprohormones) [12,13], suggesting that the two neuropeptide genes are not evolutionarily related.

[☆] The nucleotide sequence reported in this paper has been submitted to the GenBank Data Bank with Accession No. AY 345586.

* Corresponding author. Fax: +45 353231200.

E-mail address: cgrimmelikhuijzen@bi.ku.dk (C.J.P. Grimmelikhuijzen).

URL: <http://www.bi.ku.dk/staff/staff-vip-details.asp?ID=90>.

The FMRFamide preprohormone from the malaria mosquito *Anopheles gambiae* contains a neuropeptide sequence, PDRNFLRFamide (*Anopheles* FMRFamide-3), that resembles a myosuppressin (Cazzamali et al., in preparation). *Anopheles* FMRFamide-3 acts as a genuine FMRFamide on the *Anopheles* FMRFamide receptor (Cazzamali et al., in preparation). It would be important to see whether *Anopheles* FMRFamide-3 also has myosuppressin properties, which would intermingle the insect FMRFamide and myosuppressin systems. This, however, would require the cloning of the *Anopheles* myosuppressin receptor. We have previously cloned and identified (“deorphanized”) the first two insect myosuppressin receptors from the fruitfly *Drosophila melanogaster* [14]. Here, we describe the cloning and functional characterization of a myosuppressin receptor from *Anopheles*. This is, to our knowledge, the first report on the molecular identification of a mosquito neuropeptide receptor.

Materials and methods

Total RNA was isolated from adult *A. gambiae* (strain KWA, kindly supplied by Drs. N. Hill and P. Aiyenuro, London School of Hygiene and Tropical Medicine, UK) using TRIzol Reagent (Life Technologies) and treated with DNase using the DNA-free kit (Ambion). cDNA was synthesized using the SMART RACE cDNA Amplification Kit (Clontech). For 3' RACE, the sense primer 5'-CGTGATGGACGTGCTGGCGTTGGTCAA-3' and the nested sense primer 5'-GGACCGGTGGATGGCGGTGCCG-3' (corresponding to positions 1080–1108 and 1190–1212 of Fig. 1) were used. For 5' RACE, the antisense primer 5'-GCTGCCAAAGATGCACACCAGCAGGCAG-3' followed by the nested antisense primer 5'-GGCCCCATCCTCATCGCCTGCCAG (corresponding to positions 174–201 and 57–81 of Fig. 1) were used. The PCR program was 94 °C for 3 min, then 20 cycles touchdown, 94 °C for 30 s, 68 °C for 45 s, decreasing 0.5 °C per cycle, 72 °C for 3 min, followed by 25 cycles at 94 °C for 30 s, 58 °C for 45 s, and 72 °C for 3 min.

The coding region was amplified using the sense primer 5'-CACCATGAGCTCGACGGAATTGG-3' and the antisense primer 5'-GGACGTCCCGCGTGGCGCTA-3' (the underlined nucleotides correspond to positions 1–19 and 1281–1301 of Fig. 1), cloned into pCR4-TOPO (Invitrogen), sequenced, and cloned into the pIRES-hrGFP-1a/Neo-module (Stratagene), using *EcoRI*. Cell culture, cell transfection, and the bioluminescence assay were performed as in [14–17].

DNA sequence comparisons were carried out using the Lasergene software package (DNASTAR). ClustalW was used for protein sequence alignments (www.npsa-pbil.ibcp.fr).

Results

We have previously identified two myosuppressin receptors from *Drosophila* [14]. Blasting of their sequences against the genomic database sequences from the malaria mosquito (*A. gambiae*) Genome Project (<http://www.ncbi.nlm.nih.gov/BLAST/Genome/Fly-Blast.html>) revealed the sequence of a putative mosquito

myosuppressin receptor. To obtain the cDNA, encoding this receptor, we carried out PCR, and 5' and 3' RACE PCR, using primers directed against the exons of the putative myosuppressin receptor gene, and cDNA from adult *A. gambiae* as a template.

The receptor cDNA contains a polyadenylation signal and a poly(A)⁺ tail. The start codon is preceded by several in-frame stop codons (Fig. 1). The cDNA codes for a protein of 427 amino acid residues, containing seven transmembrane helices, which is characteristic for G protein-coupled receptors (Fig. 1).

A comparison of the cDNA of Fig. 1 with the genomic sequence of *A. gambiae* revealed a small number of nucleotide differences, which, however, did not result in amino acid residue differences (Table 1). This comparison also showed the presence of two introns in the receptor gene (Table 2).

An alignment of the mosquito receptor with the two *Drosophila* myosuppressin receptors showed 47–48% amino acid residue identity (52–53% in the transmembrane region) and 57–58% similar residues (63–64% in the transmembrane region) (Fig. 2). Furthermore, all three receptors had two introns with the same intron phasings in common, showing that the three receptor genes are evolutionarily closely related (Fig. 2; Table 2).

We stably expressed the coding region of the mosquito receptor gene (Fig. 1) in Chinese hamster ovary (CHO) cells and established clonal cell lines that expressed the receptor efficiently. These cells also stably expressed the promiscuous G protein, G-16 [15]. Two days before the bioassay, we transiently transfected the cells with DNA, coding for apoaquorin and 3 h before the assay, we added coelenterazine to the cell culture medium. An activation of G-16-coupled receptors in these pretreated cells, would lead to an IP₃/Ca²⁺-mediated bioluminescence response, which could easily be measured and quantified [14–17].

We tested a library of 35 insect neuropeptides and biogenic amines on the transfected cells. Low concentrations of *Drosophila* myosuppressin (TDVDHVFLRFamide) activated the receptor (EC₅₀, 1.6 × 10^{−8} M) (Fig. 3), whereas the other neuropeptides and neurohormones, including those that resembled *Drosophila* myosuppressin in their C-terminal moieties, such as the *Anopheles* FMRFamides (we tested SALDKNFMRamide, PDRNFLRFamide, STGSGYMRamide, and AGNLMRamide), short neuropeptide F-1 (AQ RSPSLRLRFamide), and perisulfakinin [EQFDDY(SO₃H)GHMRamide] [1,11,18], did not give a response (all tested up to 10^{−5} M). This showed that the receptor is specific for myosuppressin.

Searching of the genomic database from *A. gambiae* (<http://www.ncbi.nlm.nih.gov/BLAST/Genome/Fly-Blast.html>) revealed only one gene coding for a myosuppressin preprohormone, which is in agreement with

Download English Version:

<https://daneshyari.com/en/article/10771570>

Download Persian Version:

<https://daneshyari.com/article/10771570>

[Daneshyari.com](https://daneshyari.com)