



Review

Resonance Raman applications in investigations of cytochrome *c* oxidase[☆]

Takashi Ogura

Graduate School of Life Science, University of Hyogo, Koto 3-2-1, Kamigori-cho, Ako-gun, Hyogo 678-1297, Japan

ARTICLE INFO

Article history:

Received 22 September 2011

Received in revised form 24 November 2011

Accepted 24 November 2011

Available online 8 December 2011

Keywords:

Cytochrome *c* oxidase

Proton pump

Oxygen activation

Resonance Raman

Mitochondria

ABSTRACT

Recent applications of resonance Raman (RR) spectroscopy in investigations of cytochrome *c* oxidase (CcO) are reviewed. Red-excited RR spectra for the fully oxidized “as-isolated” CcO tuned to the ligand-to-metal charge transfer absorption band at 655 nm exhibit a Raman band at 755 cm^{-1} assignable to the ν_{OO} stretching mode of a peroxide. Binding of CN^- diminishes the RR band concomitant with the loss of the charge transfer absorption band. This suggests that a peroxide forms a bridge between heme α_3 and Cu_B . Time-resolved RR spectroscopy of whole mitochondria identified a band at 571 cm^{-1} arising from the oxygenated intermediate at $\Delta t = 0.4, 0.6$ and 1.4 ms . Bands at 804 and 780 cm^{-1} of the P and F intermediates were observed at $\Delta t = 0.6$ and 1.4 ms , respectively. The coordination geometries of the three intermediates are essentially the same as the respective species observed for solubilized CcO. However, the lifetime of the oxygenated intermediate in mitochondria was significantly longer than the lifetime of this intermediate determined for solubilized CcO. This phenomenon is due either to the pH effect of mitochondrial matrix, the effect of ΔpH and/or $\Delta\psi$ across the membrane, or the effect of interactions with other membrane components and/or phospholipids. This article is part of a Special Issue entitled: Respiratory Oxidases.

© 2011 Elsevier B.V. All rights reserved.

1. Introduction

Three-dimensional structures provide the basis for clarifying the mechanisms of reactions catalyzed by respiratory enzymes. In such structures, the coordinates of individual atoms can be determined at resolutions as high as 0.1 \AA by X-ray crystallography. Vibrational spectroscopy, on the other hand, determines the energy of vibrational levels and it is indirectly correlated with the lengths of bonds between atoms. A correlation between a bond length in a typical molecule and the vibrational frequency of the bond can be determined from data obtained by vibrational spectroscopy and X-ray crystallography. A distance change of 0.01 \AA (1 pm) or smaller may be detected by vibrational spectroscopy based on the established correlations (see below). Such a subtle change in bond length reflects a change in the electronic state and thus, the reactivity of a functional group. The ability to provide information with respect to the reactivity of a functional group is unique to vibrational spectroscopy and is essential in order to elucidate the role of the functional group in the reaction mechanism. For example, a linear relationship between the O—O stretching frequency ($\nu_{\text{OO}}/\text{cm}^{-1}$) and the corresponding O—O bond length ($r_{\text{OO}}/\text{\AA}$) ($\nu_{\text{OO}} = 5098.4 - 2963.3 \times r_{\text{OO}}$) is observed for a series of different species of O_2 [1,2] (O_2^+ : 1858 cm^{-1} and 1.12 \AA , O_2 : 1555 cm^{-1} and 1.21 \AA , O_2^- : 1108 cm^{-1} and 1.28 \AA , O_2^{2-} : 760 cm^{-1} and 1.49 \AA) when

bound to metal complexes. The O—O and M—O (metal–oxygen) bond orders are inversely correlated. This is rationalized by variable charge transfer from the metal to O_2 through a covalent interaction between the metal d_{xy} orbital and the in-plane π^* orbital of O_2 . The degree of charge transfer serves as the basis of the properties and indicates the reactivity of the metal–oxygen adduct. In addition, since vibrational spectroscopy has a time resolution in the picosecond range, it is powerful for determining the structure of transient species in chemical reactions. The advantage of such time resolution is that if there are multiple conformers in a given molecule, the molecule gives multiple signals originating from the respective conformers. In this case, the structures of individual conformers could be determined. However, the three dimensional structure of proteins cannot be determined by vibrational spectroscopy. For many enzymes, both vibrational spectroscopic data and X-ray crystallographic data are now available and this combination of vibrational spectroscopy and X-ray crystallography is becoming more and more important for elucidating reaction mechanisms as a result of the advantages of these two techniques.

Respiratory enzymes such as cytochrome *c* oxidase (CcO) pump protons across the membrane [3]. The proton pump is driven by the free energy change accompanied by dioxygen reduction. Respiratory control is a phenomenon in which the electron transfer rate is affected by the proton motive force $\Delta\mu_{\text{H}^+}^+$, consisting of the proton concentration gradient (ΔpH) and the membrane potential ($\Delta\psi$). These parameters have been regarded as providing physiologically important control in order to attain the highest efficiency of energy conversion as a whole. Three quarters of the weight of the inner mitochondrial membrane is occupied by membrane proteins and thus the

[☆] This article is part of a Special Issue entitled: Respiratory Oxidases.
E-mail address: ogura@sci.u-hyogo.ac.jp.

respiratory protein complexes are interacting with other proteins and phospholipids. The reactions of the membrane proteins might be affected by these interactions. For solubilized membrane proteins, the interactions with other proteins and the respiratory control are lost. Thus, it is desirable to determine the structure and pursue the reaction of the membrane proteins *in situ* to clarify the reaction mechanisms under physiological conditions. However, the development of such a technique has not yet been attained. The improved RR method has been applied to whole mitochondria and the oxygen activation reaction by CcO has been successfully detected.

Recent applications of RR spectroscopy to investigations of CcO are described in this review. The first application is with respect to the fully oxidized “as-isolated” state. A resonance Raman band assignable to a peroxide molecule was detected, suggesting a Fe—O—O—Cu_B structure with an O—O distance of 1.5 Å. The second application is with respect to the oxygen activation reaction in whole mitochondria. The results show that essentially the same reaction intermediates are involved and that the lifetime of the oxygenated intermediate is significantly longer than that of solubilized CcO. This indicates the presence of a controlling mechanism in mitochondria which could be related to respiratory control.

2. Cytochrome c oxidase

Cytochrome c oxidase (CcO) (ferrocytochrome c: oxygen oxidoreductase, EC 1.9.3.1) is the terminal component of the mitochondrial respiratory chain and catalyzes the four electron reduction of dioxygen. The reduction of dioxygen is coupled with the process of proton pumping across the mitochondrial inner membrane with a stoichiometry of H⁺/e⁻. Since a single chemical proton is consumed in the mitochondrial matrix per electron, a proton concentration difference of 3 is generated by transfer of 1 electron. A functional unit of bovine CcO consists of 13 different subunits and its molecular weight is 210 kD [4]. CcO has four redox-active metal centers: the Cu_A site (a binuclear Cu site), heme *a*, heme *a*₃ and the Cu_B. Cu_A accepts electrons from cytochrome *c* and donates them to heme *a*. Heme *a* delivers electrons to the O₂ reduction site and is proposed to act as the driver of the proton pump. The O₂ reduction site is a binuclear metal center consisting of heme *a*₃ and Cu_B. The mechanism of dioxygen reduction and the coupling mechanism between electron transfer and proton pumping have been extensively studied. The oxygenated species, the P intermediate, the F intermediate and the O intermediate have been respectively determined to have Fe—O₂, Fe O, Fe O and Fe—OH units at the heme *a*₃ site based primarily on the iron–oxygen stretching ($\nu_{\text{Fe—O}}$) frequencies detected by RR spectroscopy [5–8].

The fully oxidized form of CcO generated under turnover conditions pumps protons upon receiving electrons, while the resting oxidized form does not [9]. Thus, the functions of the two oxidized forms are distinct from each other and the key to understanding the different functions should be found in their different structures. The coordination geometry of the heme *a*₃ site in the fully oxidized form under turnover conditions has been determined by RR spectroscopy to be Fe³⁺—OH⁻ [10,11]. However, the coordination geometry of the resting oxidized form (the fully oxidized “as isolated” form) was not well established. The presence of a bridging ligand between heme *a*₃ and Cu_B was postulated based on data obtained by EPR spectroscopy [12]. Four electron equivalents should be sufficient for full reduction of the protein, since there are four redox-active metal centers in a functional unit of CcO. However, the “as-isolated” form was shown to require six electron equivalents for full reduction, while the oxidized form under turnover conditions requires four electron equivalents [13]. The results suggested that the bridging ligand is peroxide (O₂²⁻). The X-ray structure of the resting oxidized CcO showed an area of electron density comparable to that of two oxygen atoms [14]. Although the possibility of attributing the electron density to Cl⁻ was suggested as an alternative interpretation [15], it was

recently confirmed by an X-ray anomalous dispersion analysis that the electron density is due to the presence of O₂²⁻ [16]. The O—O distance of the Fe—O—O—Cu_B bridged structure was determined to be 1.70 Å. However, the O—O bond distance is too long to represent an ordinary peroxide with a typical O—O distance of 1.5 Å. To our knowledge, no peroxide species with an O—O bond distance of 1.7 Å has been reported. Thus, identifying the factors responsible for the discrepancy should help to clarify the chemical nature of the bridging ligand between heme *a*₃ and Cu_B.

Fig. 1 depicts RR spectra of the fully oxidized “as-isolated” (A), fully oxidized “as-isolated” and cyanide-bound CcO (B) and their difference spectrum (C=A-B) [17]. It is evident in Spectrum 1C that there is a band at 755 cm⁻¹. This means that the band at 755 cm⁻¹ for the fully oxidized “as-isolated” form loses its intensity upon binding of CN⁻. The inset shows that the absorption band at 655 nm for fully oxidized “as-isolated” CcO (A') loses its intensity upon cyanide binding (B'). The RR scattering was excited at 647.1 nm in resonance with the ligand-to-metal charge transfer transition at 655 nm [18,19]. The prolonged laser illumination with higher power (140 mW) diminished the Raman band at 755 cm⁻¹ due to photoreduction [20]. The enhancement of only one Raman band at 755 cm⁻¹ in Spectrum 1C indicates that the band arises from a diatomic molecule, since we expect only one stretching vibration for a diatomic molecule. These results led to the conclusion that the band at 755 cm⁻¹ originates from the ν_{OO} mode of the bridging peroxide. If the peroxide resides at a position different from the binuclear site, then enhancement of the RR band is not expected. There is no RR band assignable to the Fe—O or Cu—O stretching mode in Spectrum 1C, while both the O—O and Fe—O stretching modes are detectable for a side-on peroxy and an end-on hydroperoxy Fe-porphyrin model complexes [21]. Since the RR spectral characteristics of the Fe—O—O—Cu type bridging peroxide has not yet been well established, the reason of the undetectability of neither the Fe—O nor Cu—O stretching mode is not clear at present. As mentioned previously, there is a linear relationship between the O—O stretching frequency and the O—O bond length [1,2]. Based on the relationship ($\nu_{\text{OO}} = 5098.4 - 2963.3 \times r_{\text{OO}}$), the r_{OO} of the fully oxidized “as-isolated” CcO was calculated to be 1.47 Å, which is close to the bond distance of an ordinary peroxide, but the value (1.7 Å) obtained by X-ray crystallography was significantly larger. The discrepancy between the X-ray and Raman results might be due to inaccuracies in the X-ray structural analysis even at 1.8 Å resolution [14]. However, the ¹⁸O₂-isotopic substitution effects on the RR spectra have not been confirmed because the method to prepare the fully oxidized “as-isolated” form from the fully reduced form under an ¹⁸O₂ atmosphere has not been




Fig. 1. Resonance Raman spectra of fully oxidized “as-isolated” cytochrome c oxidase (A) and its cyanide adduct (B) excited at 647.1 nm. Spectrum C is the Raman difference spectrum (A–B). The laser power was 11 mW. The accumulation period for spectra A and B was 90 min. Spectral features marked with an asterisk are due to emissions of the Kr⁺ laser used. The inset shows the visible absorption spectra (A') and (B') for the species corresponding to Raman spectra (A) and (B), respectively. (Reproduced with permission from Sakaguchi et al., Ref. [17]).

Download English Version:

<https://daneshyari.com/en/article/10795897>

Download Persian Version:

<https://daneshyari.com/article/10795897>

[Daneshyari.com](https://daneshyari.com)