

Transforming growth factor type beta (TGF- β) requires reactive oxygen species to induce skeletal muscle atrophy

Johanna Abrigo, Juan Carlos Rivera, Felipe Simon, Daniel Cabrera, Claudio Cabello-Verrugio

PII: S0898-6568(16)30010-9
DOI: doi: 10.1016/j.cellsig.2016.01.010
Reference: CLS 8620

To appear in: *Cellular Signalling*

Received date: 19 September 2015
Revised date: 28 December 2015
Accepted date: 24 January 2016

Please cite this article as: Johanna Abrigo, Juan Carlos Rivera, Felipe Simon, Daniel Cabrera, Claudio Cabello-Verrugio, Transforming growth factor type beta (TGF- β) requires reactive oxygen species to induce skeletal muscle atrophy, *Cellular Signalling* (2016), doi: 10.1016/j.cellsig.2016.01.010

Transforming growth factor type beta (TGF-β) requires reactive oxygen species to induce skeletal muscle atrophy

Johanna Abrigo[a,b], Juan Carlos Rivera [a,b], Felipe Simon[b,c], Daniel Cabrera[d,e] and Claudio Cabello-Verrugio[a,b,*]

[a] Laboratory of Biology and Molecular Physiopathology, Department of Biological Sciences, Faculty of Biological Sciences & Faculty of Medicine, Universidad Andrés Bello. Santiago, Chile.

[b] Millennium Institute on Immunology and Immunotherapy, Santiago, Chile.

[c] Laboratory of Integrative Physiopathology. Department of Biological Sciences, Faculty of Biological Sciences & Faculty of Medicine, Universidad Andrés Bello. Santiago, Chile.

[d] Departamento de Gastroenterología. Facultad de Medicina. Pontificia Universidad Católica de Chile. Santiago. Chile.

[e] Departamento de Ciencias Químicas y Biológicas. Facultad de Salud, Universidad Bernardo O Higgins. Santiago, Chile.

*** Corresponding author:** Departamento de Ciencias Biológicas, Facultad de Ciencias Biológicas & Facultad de Medicina, Universidad Andrés Bello, Santiago, Chile. Avenida República 239, Santiago, 8370146, Chile. *E-mail address:* claudio.cabello@unab.cl (C. Cabello-Verrugio).