

# Kinetics of the plastoquinone pool oxidation following illumination Oxygen incorporation into photosynthetic electron transport chain

Boris Ivanov\*, Maria Mubarakshina, Sergey Khorobrykh

*Institute of Basic Biological Problems, Russian Academy of Sciences, Photosynthetic Electron Transport Lab,  
Pushchino 142290, Moscow Region, Russian Federation*

Received 29 December 2006; revised 18 February 2007; accepted 22 February 2007

Available online 1 March 2007

Edited by Richard Cogdell

**Abstract** The oxidation of the PQ-pool after illumination with 50 or 500  $\mu\text{mol quanta m}^{-2} \text{s}^{-1}$  was measured in isolated thylakoids as the increase in  $\Delta A_{263}$ , i.e., as the appearance of PQ. While it was not observed under anaerobic conditions, under aerobic conditions it was biphasic. The first faster phase constituted 26% or 44% of total reappearance of PQ, after weak or strong light respectively. The dependence on oxygen presence as well as the correlation with the rate of oxygen consumption led to conclusion that this phase represents the appearance of PQ from  $\text{PQ}^{\cdot-}$  produced in the course of  $\text{PQH}_2$  oxidation by superoxide accumulated in the light within the membrane.

© 2007 Federation of European Biochemical Societies. Published by Elsevier B.V. All rights reserved.

**Keywords:** Thylakoids; Electron transport chain; Plastoquinone; Oxygen reduction

## 1. Introduction

Plastoquinone pool (PQ-pool) oxidation estimated as an increase in the area over the chlorophyll fluorescence induction curve after illumination of thylakoids demonstrated biphasic kinetics in a seconds range with a first rapid phase followed by slow tail [1,2]. With the same approach, biphasic kinetics of PQ-pool oxidation were found also in leaf disks of *Arabidopsis* and it was stressed that its rate was insensitive to inhibitors of oxidases [3]. The reasons of such kinetics were not discussed. The components of PQ-pool are the only membrane-embedded electron carriers which are not included into the molecular assemblies where electron pathways are strictly determined. This circumstance provides them with more possibilities for reacting not only with adjoining carriers of photosynthetic electron transport chain (PETC). It was shown that PQ-pool oxidation did not occur in the absence of oxygen [4]. Oxygen concentration inside the thylakoid membrane can be estimated as 2–10  $\text{O}_2$  per 1000 Chl, with higher values

for more hydrophobic areas where its solubility is higher. It is comparable with concentrations of the electron carriers of PETC. Oxygen was found to be reduced by PQ-pool when  $\text{PQH}_2$  oxidation by *cytb<sub>6</sub>f*-complex was inhibited [5]. According the data in [5] this reduction, occurs at first as the univalent reduction of dioxygen by  $\text{PQ}^{\cdot-}$  followed by reduction of superoxide radical,  $\text{O}_2^{\cdot-}$ , by  $\text{PQH}_2$ . It is moderate and is saturated at low-light intensity [5], while the PQ-pool participation in oxygen reduction in the entire PETC, being low in weak light attains 70% in strong light [6]. To explain such dependence, it was hypothesized [6] that  $\text{PQH}_2$  can reduce superoxide radicals produced in both PQ-pool and PS I.

In this work we have found that PQ-pool oxidation measured as PQ reappearance after switching off the light consisted of two phases; after stronger light the extent of the initial rapid phase was higher, that coincided with a higher rate of oxygen consumption in the light. The data are discussed as indicating the participation of superoxide generated within the membrane in the light, in the first phase of the post-illumination PQ-pool oxidation.

## 2. Materials and methods

Thylakoids from leaves of pea plants grown in a greenhouse were isolated as described [5]. The basic reaction medium contained 0.4 M sucrose, 20 mM NaCl, 5 mM  $\text{MgCl}_2$ , 10  $\mu\text{M}$  Gr D, 50 mM Hepes-KOH (pH 7.8); thylakoids were with 100, 15 and 30  $\mu\text{g}$  Chl per ml in the measurements of absorbance changes, oxygen uptake, and Chl *a* fluorescence, respectively. Absorbance changes were measured, using a dual-wavelength spectrophotometer (Hitachi 553, Japan), in a quartz cuvette with optical length 5 mm and the suspension thickness 4 mm perpendicular to actinic light. The absorption changes at 263 nm, – the minimum in the reduced-minus-oxidized spectrum of PQ [7], – were corrected for changes in transmittance and/or redox-state of other components as follows. For every variant the kinetics of both  $\Delta A_{243-283}$  and  $\Delta A_{263-283}$  were recorded; the line connecting zero on the Y-axis at 283 nm ( $\Delta A_{283-283}$ ) and the value of  $\Delta A_{243-283}$  at 243 nm at any moment after switching off the light was taken as the base line; the difference between the point at 263 nm at this line and  $\Delta A_{263-283}$  at a corresponding moment was taken as  $\Delta A$  due to PQ appearance. The measurements started 2 s after cessation of illumination. The differential extinction coefficient at 263 nm was taken as  $13 \text{ mM}^{-1} \text{ cm}^{-1}$  [8]. The anaerobic conditions were created by adding 10 mM glucose, 25  $\mu\text{g/ml}$  glucose oxidase and 500  $\mu\text{g/ml}$  catalase. Light-induced *cyt f* redox-changes were determined as  $\Delta A_{554-545}$ ; the blue-green filter SZS-22 (Russia) screened the photomultiplier from scattered light. Chlorophyll *a* fluorescence was measured under stirring in square quartz cuvette using a PAM fluorometer (WALZ, Germany). Oxygen concentration changes in a stirred thylakoid suspension (3.2 ml) were measured at 21 °C in a glass vessel with a Clark-type oxygen electrode; in these experiments, sucrose was 0.1 M, and Gr D was

\*Corresponding author. Fax: +7 4967 330532.

E-mail address: ivabor@issp.serpukhov.su (B. Ivanov).

**Abbreviations:** Chl, chlorophyll; DCMU, 3-(3,4-dichlorophenyl)-1,1-dimethylurea; DNP-INT, dinitrophenylether of 2-iodo-4-nitrothymol; Gr D, gramicidin D; MV, methyl viologen; PETC, photosynthetic electron transport chain; PQ, plastoquinone;  $\text{PQH}^+$ ,  $\text{PQ}^{\cdot-}$  plastoquinone;  $\text{PQH}_2$ , plastoquinone; PS I, photosystem I; PS II, photosystem II; TMPD, *N,N,N',N'*-tetramethyl-*p*-phenylene diamine

1  $\mu\text{M}$ . The separate PS I operation was achieved by additions of 10  $\mu\text{M}$  DCMU, 5 mM ascorbate and 0.1 mM TMPD to provide electron donation to plastocyanin/P700. A saturating amount of superoxide dismutase was added to reaction mixtures in these experiments to prevent the reaction of superoxide with ascorbate. It was also added in the experiments without donor pair to provide the same conditions in the both analyses. Light was filtered through a red cut-off ( $\lambda > 600$  nm) and heat absorbing filters. Light intensity measured with a Li-Cor quantum meter was varied using neutral filters. Chlorophyll was determined in 95% ethanolic extracts [9]. Stock solutions of DCMU and Gr D were in dimethylsulfoxide.

### 3. Results

Under aerobic conditions, a gradual increase in absorption at 263 nm toward the initial dark level occurred after switching off the actinic light, and the fraction of PQ appearing was estimated as the ratio of the difference between ‘aerobic’ and ‘anaerobic’  $\Delta A_{263}$  to ‘anaerobic’  $\Delta A_{263}$  (Fig. 1A and A’). Under anaerobic conditions only a small and rapid (within 3–4 s) increase in absorption at 263 nm occurred, and the concentration of the reduced PQ molecules preserved after illumination with weak or strong light was similar, 22–24 per 1000 Chl (after correction for flattening effect at 263 nm), that is close to total PQ concentration in thylakoids [2,7]. The kinetics of PQ reappearance was biphasic, and the semi-logarithmic plots of the kinetics are shown as broken lines (Fig. 1A, A’ and inserts). The slopes of the fast phases after illumination with weak or strong lights were  $0.08\text{ s}^{-1}$  and  $0.14\text{ s}^{-1}$ , respectively, and the contributions of fast phase were 26% and 44%, when the possible slow phase contributions, which could be found by extension of a line approximating this phase to the intersection with Y-axis, were subtracted. The slopes of first quick phases could be considered as apparent ‘constants’ of the pseudo-first-order reactions characterizing PQ-pool oxidation in weak and strong light, respectively.

The extents of PQ-pool reduction under aerobic conditions under both light intensities were similar as it is seen from the chlorophyll *a* fluorescence yields (Fig. 1B and B’). Therefore a difference in the level of PQ-pool reduction was not responsible for the difference in the contributions of fast phase of PQ reappearance after illumination. The high-potential components of PETC might be the first acceptors of electrons from the PQ-pool, if they were oxidized in the light. Under our experimental conditions, we did not observe any light-induced redox changes of *cyt f* in both strong and weak light (Fig. 1C and C’) (in the presence of Mv and in the absence of Gr D it demonstrated the characteristic redox-changes, dashed line in Fig. 1C’). Thus, the initial rapid phase of PQH<sub>2</sub> oxidation could not be explained by electron flow toward FeS-Rieske centre, *cyt f* and plastocyanin.


Since we subtracted the small changes in absorption at 263 nm under anaerobic conditions, the oxygen has to be implicated into process of PQ-pool oxidation. The rate of oxygen consumption in the light did increase with the increase in light intensity under operation of whole PETC as well as under conditions where electron transfer occurred only through PS I. The oxygen reduction rates (Fig. 2) were calculated from oxygen consumption rates, using the stoichiometries:  $4\text{ e}^-$  per 1  $\text{O}_2$  consumed, for the entire chain, and  $2\text{ e}^-$  per 1  $\text{O}_2$  consumed, for ‘isolated’ PS I [10]. In both cases the rates were limited by oxygen reduction [6], and therefore it was possible to com-

pare them. It is seen that electron transport increased more steeply in the entire PETC.

### 4. Discussion

The implication of oxygen in the process of PQ-oxidation allowed the comparison between the higher fast phase of PQ appearance in stronger light and the higher rate of oxygen reduction. A putative plastoquinone oxidase was not involved in the process determining the biphasic kinetics of PQ-pool oxidation in the above experiments. The addition of catalase after illumination of thylakoids increased the oxygen concentration in the thylakoid suspension to the level before illumination (not shown, but see [11]), and this indicated that the final product of oxygen reduction was  $\text{H}_2\text{O}_2$ , but not  $\text{H}_2\text{O}$ . The role of enzymes in the kinetics of PQH<sub>2</sub> oxidation was not detected in [3] (see Section 1).

During illumination there is some accumulation of superoxide within the thylakoid membrane, since  $\text{O}_2^-$  is stable there over the seconds range [12,13]. Since superoxide production depended on the light intensity (Fig. 2, curve 1) superoxide accumulation apparently determined the extent of the fast phase of PQ appearance after switching off the light. The process could be the rapid production of  $\text{PQ}^{\cdot-}$  in the reaction of PQH<sub>2</sub> with the accumulated  $\text{O}_2^-$ , followed by PQ appearance from both  $\text{PQ}^{\cdot-}$  dismutation and  $\text{PQ}^{\cdot-}$  oxidation. The difference in  $E'_0$  values of pairs  $\text{PQ}^{\cdot-}/\text{PQH}_2$  (0.37 V) [14,15] and  $\text{O}_2^-/\text{H}_2\text{O}_2$  (0.94 V) [16] provides high-equilibrium constant,  $>1.5 \times 10^4$ , of the reaction



and the electrostatic repulsion does not hamper it, since  $pK_1$  of PQH<sub>2</sub> is close to 11. An increase of this reaction in stronger light was confirmed by an increase of the intramembrane  $\text{H}_2\text{O}_2$  production with increasing light intensity [11]. The produced  $\text{PQ}^{\cdot-}$  could either dismutate,



or be oxidized by dioxygen



The difference of redox potentials of pairs  $\text{PQ}/\text{PQ}^{\cdot-}$ ,  $-170\text{ mV}$ , and  $\text{O}_2/\text{O}_2^{\cdot-}$ ,  $-160\text{ mV}$ , is not high. The dismutation is possibly more significant due to high-equilibrium constant,  $10^{10}$  [17], however, it can be partially hampered by an electrostatic repulsion ( $pK$  of  $\text{PQ}^{\cdot-}$  is 7.0) as well as the distance between  $\text{PQ}^{\cdot-}$  molecules formed.

After exhaustion of the superoxide that was accumulated in the light, the production of additional (to that produced in the reverse reaction (2))  $\text{PQ}^{\cdot-}$  molecules sharply decreased. The oxidation of residual PQH<sub>2</sub> molecules could proceed autocatalytically as follows: reverse reaction (2)  $\rightarrow$  reaction (3)  $\rightarrow$  reaction (1) producing additional  $\text{PQ}^{\cdot-}$ , and so on.  $\text{PQ}^{\cdot-}$  was produced in a limited amount, and this determined the slow second phase of PQ appearance in Fig. 1A and A’. Obviously, the same process determined the slow phase of the dark PQ-pool oxidation, which was measured using fluorescence induction [1–3]. The oxidation of reduced plastoquinone placed into egg yolk liposomes is very slow: the second order rate constant is close to  $10^{-7}\text{ }\mu\text{M}^{-1}\text{ s}^{-1}$  [18], and an apparent first-order rate constant we estimated from the data in this

Download English Version:

<https://daneshyari.com/en/article/10872613>

Download Persian Version:

<https://daneshyari.com/article/10872613>

[Daneshyari.com](https://daneshyari.com)