



Short Communication

High-intensity Fitness Training Among a National Sample of Male Career Firefighters



Sara A. Jahnke^{1,*}, Melissa L. Hyder¹, Christopher K. Haddock¹, Nattinee Jitnarin¹, R. Sue Day², Walker S. Carlos Poston¹

¹ Center for Fire, Rescue, and EMS Health Research, Institute for Biobehavioral Health Research, National Development and Research Institutes, Leawood, KS, USA

² University of Texas Health Science Center at Houston, Houston, TX, USA

ARTICLE INFO

Article history:

Received 12 September 2014

Received in revised form

18 November 2014

Accepted 6 December 2014

Available online 12 January 2015

Keywords:

firefighter

fitness

high intensity training

ABSTRACT

Obesity and fitness have been identified as key health concerns among USA firefighters yet little is known about the current habits related to exercise and diet. In particular, high-intensity training (HIT) has gained increasing popularity among this population but limited quantitative data are available about how often it is used and the relationship between HIT and other outcomes. Using survey methodology, the current study evaluated self-reported HIT and diet practice among 625 male firefighters. Almost one-third (32.3%) of participants reported engaging in HIT. Body composition, as measured by waist circumference and percentage body fat, was significantly related to HIT training, with HIT participants being approximately half as likely to be classified as obese using body fat [odds ratio (OR) = 0.52, 95% confidence interval (CI) = 0.34–0.78] or waist circumference (OR = 0.61, 95% CI = 0.37–0.98). Those who engaged in HIT were more than twice as likely as those who did not (OR = 2.24, 95% CI = 1.42–3.55) to meet fitness recommendations. Findings highlight directions for future prevention and intervention efforts.

© 2015, Occupational Safety and Health Research Institute. Published by Elsevier. All rights reserved.

1. Introduction

Firefighting is an inherently dangerous, physically demanding occupation requiring adequate levels of fitness to fulfill mandatory job duties. Physically taxing activities performed by firefighters may include hauling hose, using heavy extrication equipment, or lifting and transporting patients [1–3]. Despite the need for high levels of readiness and fitness, recent research suggests high rates of obesity among firefighters. Rates of overweight and obesity in the fire service (estimated at about 80% combined) [4,5] are higher than those found among the general United States (US) public (68% are overweight or obese) [6]. Obesity is an even greater risk for firefighters given the duties they are required to perform at a moment's notice, often in suboptimal conditions such as heat, smoke, and chemical exposure [7].

Previous studies documented that overweight and obese firefighters have elevated rates of hypertension, low High-density

lipoprotein (HDL) cholesterol, high Low-density lipoprotein (LDL) cholesterol, high triglycerides, significant yearly weight gain, lower cardiorespiratory fitness, reduced muscle strength, and more frequent fatal cardiac events [7,8]. In turn, cardiovascular disease (CVD) is the leading cause of line of duty deaths (LODDs) among firefighters, implicated in nearly half of LODDs, with the majority occurring during fire suppression activities [9]. In addition to serving as a risk factor for CVD, a study by Jahnke et al [10] found that obese firefighters were 5.2 times more likely to incur a musculoskeletal injury than their healthy weight peers. In addition, obesity has been found to be related to low rates of fitness in firefighters [4]. Evidence suggests that firefighting duties require approximately 12 metabolic equivalent units of effort (METs) [11], which is recommended by national standards as a return to work standard for firefighters following a cardiac event [12]. In a population-based sample of firefighters, nearly all of whom were obese (80% career, 95% volunteer) by body mass index (BMI)

* Corresponding author. Center for Fire, Rescue, and EMS Health Research, National Development and Research Institutes, 1920 West 143rd Street, Leawood, KS 66224, USA.
E-mail address: jahnke@ndri.org (S.A. Jahnke).

This is an Open Access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (<http://creativecommons.org/licenses/by-nc/3.0>) which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

standards would not be able to meet this fitness recommendation [4]. Given the risks firefighters face and the need for high levels of fitness, high rates of overweight and obesity are of particular concern.

Given the importance of fitness for firefighters, innovative techniques for improving their overall fitness responsive to the unique training needs, culture, and environment of the fire service [1] are needed. One promising approach to promoting fitness among firefighters is high-intensity training (HIT), which is an approach to exercise including functional movements at a high level of effort, but short in duration [13]. These exercise programs may be a better way for firefighters to prepare for what their job requires because they focus on full body movements, closely mimicking actual fire ground activities [1,13]. Roberts et al [14] evaluated a 16-week HIT program for new firefighter recruits and found 3 days per week of training for 1 hour per session resulted in significant improvements in maximal oxygen consumption (VO_{2max}). Training with HIT not only provides improvement in cardiorespiratory fitness but does so in less time than traditional training, and will be likely to prove beneficial for firefighters' constrained and unpredictable schedules [14].

While the fire service is working to improve health and wellness among its members, limited data are available about current fitness in general and HIT in particular that firefighters engage in. Thus, the purpose of the current study is to assess the self-reported health practices of male firefighters, specifically related to fitness and HIT. In addition, we will explore the relationship between self-reported engagement in HIT, obesity, and fitness among this important population. Findings will provide a foundation of knowledge from which to build future prevention and intervention efforts.

2. Materials and methods

2.1. Procedures

The data reported are from the follow-up evaluation of a longitudinal cohort study of firefighters examining general health and quality of life including measured height and weight, estimated percentage body fat (BF%), blood pressure, and self-reported measures of diet, physical activity, substance use, mood, sleep, injury, and job satisfaction. All data were collected between 2011 and 2013. Departments ($n = 20$) ranged in size from small (≤ 3 stations; $n = 7$), medium (> 3 stations, but < 350 personnel; $n = 8$), and large (≥ 350 personnel; $n = 5$) based on the fire services' classification [15]. They also encompassed different settings (4 rural, 10 suburban, 2 urban, and 4 mixed urban) and regions based on US Census regions (8 Pacific/West, 8 Mountain/Central, and 4 East/Southeast). All departments were recruited by a request for participation on The Secret List (www.firefighterclosecalls.com), which is a widely distributed listserv within the US Fire Service. Data were collected for all participants at baseline and 6-month follow-up. More detailed recruitment procedures can be found in Poston et al [16]. The protocol for the protection of participants for this study was approved by the Institutional Review Board of the National Development and Research Institutes, Inc., Leawood, KS, USA and the University of Texas Houston Health Sciences Center, Houston, TX, USA.

2.2. Measures

2.2.1. Demographics

Standard individual (e.g., age, marital status, educational level, and race) and occupational (rank and years in the fire service)

demographic data were collected using previously published questions [4].

2.2.2. Body composition: weight, BMI, BF%, and waist circumference

Height was assessed by using a portable stadiometer. Body weight and BF% were assessed using foot-to-foot bioelectrical impedance (Tanita 300; Tanita Corporation of America, Inc., Arlington Heights, IL, USA). The Tanita 300 has demonstrated strong evidence of concurrent validity ($r = 0.94$; $p < 0.001$) [17] when compared with the "criterion standard" of dual energy X-ray absorptiometry for BF assessment. It is a commonly used bioelectrical impedance field measure because of its portability and accuracy in determining BF% [17]. Waist circumference (WC) was assessed using a spring-loaded nonstretchable tape measure as recommended by the US Government obesity guidelines [18]. Standard cut-points were used for obesity ($BMI \geq 30$, $BF\% \geq 25$, $WC \geq 40$) [18].

2.2.3. HIT

Participation in HIT was assessed by asking the following question: "Do you currently base your workouts on high-intensity fitness training programs such as CrossFit, P90X, or Insanity?" Response options were: If the participant answered yes, they were asked to indicate "How long (in months) have you participated in a high-intensity training program?" and "On average, how many days a week do you do high-intensity workouts?" For analyses of HIT, participants were divided into two categories (HIT and No HIT).

2.2.4. Physical activity

The self-report of physical activity (SRPA) was used to assess a global self-rating of physical activity patterns [19]. Participants selected a value (0–7) on the SRPA most closely indicating their physical activity over the past 30 days. Indicators of the questionnaire's validity in adult populations have been established (i.e., significant correlations between SRPA ratings and measured VO_{2max}) [20]. Scores on the SRPA, BMI, and age were used to estimate VO_{2max} with a method developed and tested in comparison with measured VO_{2max} . The method has been found to be equal or better in accuracy compared to submaximal exercise testing [20]. METs were calculated based on VO_{2max} and used to classify firefighters into those meeting the suggested minimal fitness standard of 12 METs and those who did not. The cut off is based on findings of Donovan et al [2] indicating firefighting activities require 12 METs of energy as well as National Fire Protection Association standards showing a firefighter must be able to achieve 12 METs to return to work postcardiac event [12].

2.3. Statistical analysis

Data are presented as mean (M) and standard deviation (SD) for continuous variables and percentages for categorical variables. Logistic models were used to determine the relationships between engaging in HIT and body composition (BMI, WC, BF%) as well as the relationship between HIT and meeting minimum fitness recommendations. Preliminary analyses revealed that those who used HIT were significantly younger than those who did not ($M = 37.5$ years, $SD = 8.3$ years and $M = 40.4$ years, $SD = 8.9$ years respectively; $t = 3.8$, $p < 0.001$). In addition, departments engaging in wellness or fitness programs have previously been found to be significantly related to body composition [16]; therefore, all HIT logistic models were controlled for department type and age. We also examined the dose-response relationship between HIT frequency and body composition among HIT users with a logistic model. Finally, to determine whether there was a difference between firefighters who practiced HIT and those who did not, we

Download English Version:

<https://daneshyari.com/en/article/1092009>

Download Persian Version:

<https://daneshyari.com/article/1092009>

[Daneshyari.com](https://daneshyari.com)