



Contents lists available at ScienceDirect

Cryobiology

journal homepage: www.elsevier.com/locate/ycryo

Comparing ethylene glycol with glycerol and with or without dithiothreitol and sucrose for cryopreservation of bull semen in egg-yolk containing extenders [☆]

Serhat Büyükleblebici ^a, Pürhan Barbaros Tuncer ^{b,*}, Mustafa N. Bucak ^c, Umut Taşdemir ^b, Ayşe Eken ^d, Olga Büyükleblebici ^e, Emre Durmaz ^f, Serpil Sarıözkan ^g, Burcu Ü. Endirlik ^d

^a Aksaray University, Faculty of Veterinary Medicine, Department of Reproduction and Artificial Insemination, TR-68100 Aksaray, Turkey

^b Aksaray University, Aksaray Vocational School, TR-68100 Aksaray, Turkey

^c Selçuk University, Faculty of Veterinary Medicine, Department of Reproduction and Artificial Insemination, TR-42003 Konya, Turkey

^d Erciyes University, Faculty of Pharmacy, Department of Pharmaceutical Toxicology, TR-38039 Kayseri, Turkey

^e Aksaray University, Faculty of Veterinary Medicine, Department of Biochemistry, TR-68100 Aksaray, Turkey

^f Department of Toxicology, Faculty of Pharmacy, Gazi University, TR-06330 Ankara, Turkey

^g Erciyes University, Faculty of Veterinary Medicine, Department of Reproduction and Artificial Insemination, TR-38039 Kayseri, Turkey

ARTICLE INFO

Article history:

Received 11 March 2014

Accepted 12 May 2014

Available online xxx

Keywords:

Bovine

Antioxidants

Cryoprotectants

Sperm parameters

Freeze–thawing

ABSTRACT

There are few studies performed for investigating the roles of different ratio and cryoprotectants with dithiothreitol or sucrose on sperm motility characteristics and antioxidant capacities of post-thawed bull spermatozoa. The objectives of this study were to compare glycerol (G) and ethylene glycol (EG) at different concentrations as cryoprotectants and dithiothreitol (D) or sucrose (S) (with/without) as antioxidants in Tris extender for cryopreservation of bull semen. Twenty-four ejaculates obtained from three bulls were included in the study. Each ejaculate was split into four equal aliquots and diluted using both of the Tris extenders with glycerol (5% or 7%) or ethylene glycol (3% or 5%). After that, each extenders were split into three equal aliquots and diluted using both of the dithiothreitol 5 mM or sucrose 25 mM, and control (without additives) was cooled to 4 °C and frozen in 0.25-ml French straws. When compared to control, different doses cryoprotectants and antioxidants addition no significantly increased the percentages of post-thaw sperm progressive and motilities, acrosome abnormality and plasma membrane integrity ($P > 0.05$). However, EG3 + S yielded the greatest percentages of the total abnormality ($P < 0.05$). As regard to antioxidant activities G7 and EG5 led to lowest MDA activity with or without D or S but, these results were not supported to the GPx activity ($P < 0.01$). The sperm motion characteristics such as VAP, VCL, ALH and BCF gave significantly different results ($P < 0.05$). When compared the DNA integrity, different doses cryoprotectants without antioxidants addition significantly increased the percentages of the tail intensity and tail moment ($P < 0.05$). There were no significant differences observed in non-return rates among all treatment groups ($P > 0.05$).

© 2014 Elsevier Inc. All rights reserved.

Introduction

The success of cryopreservation depends on the specific susceptibility of sperm cells at low temperatures. Changes in sperm cell and its function associated to cold shock have often been reported [5,29]. Cryopreservation processes lead to the generation of reactive oxygen species (ROS) that impair sperm motility, membrane

integrity, and fertilizing ability [7,18,20]. Although bovine semen has natural defense system against the ROS attack, it is insufficient under cryopreservation due to stress [28]. Sperm plasma and acrosomal membranes are especially important with regard to survival following thawing and it is considered that there are changes on the primary organelles caused by the cryopreservation process. These changes are due to oxidative and osmotic stresses [19,43]. The level and type of cryoprotectants influence the sperm cells during freezing in semen extender [17]. In recent years, many cryoprotective agents have been used as a cryoprotectant in bull [37] and goat [40]. Glycerol (G) can cause chemical and osmotic deleterious effects on sperm cells and changes in the structure of the plasma membrane, these effects and changes can result in a lower fertility

[☆] This study was supported by the Republic of Turkey, Ministry of Agriculture and Rural Affairs, General Directorate of Agricultural Research (GDAR) Project number: 09/01/01/01.

* Corresponding author.

E-mail address: barbarostuncerp@hotmail.com (P.B. Tuncer).

rate when AI is used, which demonstrates the need for alternative cryoprotectants [6]. Besides, improvement of semen extenders with suitable antioxidant is suggested to reduce oxidative damage during freeze–thawing of bull spermatozoa [1]. Dithiothreitol (D) is known as an antioxidant and it decreases protamine disulfide bond [42]. It prevents the oxidation of sulfhydryl groups, also it has a mucolytic effect on mucoprotein disulfide bonds, which may probably damage the frozen membranes [29]. It is stated that D provides a protective effect against apoptosis and oxidative damage [15]. The addition of D seems to improve bull and human sperm motility during liquid storage or in the frozen state [14]. Sugars are known to maintain the osmotic pressure of extenders by inducing cell dehydration and less ice crystal formation into the sperm cells [30]. They also interact with phospholipid membranes at low hydration and thus cause stabilization of the membranes. Furthermore, sugars are utilized by spermatozoa as an energy source during glycolysis and mitochondrial oxidative phosphorylation to support sperm motility [27]. Addition of sugars in extenders, such as sucrose, tends to protect the sperm cells. This sugar, as non-permeating cryoprotectant, is not able to diffuse across the plasma membrane, but creates an osmotic pressure that induces cell dehydration before freezing. Thus it decreases the extent of cell injury by intracellular ice formation [21].

There are few studies performed for investigating the roles of different ratio and cryoprotectants with D or S on sperm motility characteristics and antioxidant capacities of post-thawed bull spermatozoa. Thus, the aim of this study is to determine the effects of G or EG (with or without D or S) on post-thaw sperm parameters (motility, acrosome and total abnormalities, plasma membrane integrity), antioxidant activities (MDA and GPx) and DNA integrity as well as conception rate.

Material and methods

Animals, semen collection and semen processing

Three Holstein bulls (3–4 years of age) were housed at the Lalahan Livestock Central Research Institute (Ankara, Turkey), and maintained under uniform feeding and housing conditions. A total number of 24 ejaculates (8 ejaculates for each Holstein bull) were collected from the bulls with the aid of an artificial vagina twice a week, according to AI standard procedures. The volume of ejaculates was measured in a conical tube graduated at 0.1 ml intervals and sperm concentration was determined by means of an Accucell photometer (IMV, L'Aigle, France). Sperm motility was estimated using phase-contrast microscopy (200 \times). The ejaculates containing spermatozoa with >80% forward progressive motility and concentrations higher than 1.0×10^9 spermatozoa/ml were used in the study. Immediately after collection, the ejaculates were immersed in a warm water bath at 35 °C until their assessment in the laboratory. A Tris-based extender (T) (189.5 mM of Tris, 63.2 mM of citric acid, 55.5 mM of fructose, 20% v/v egg yolk, and 1000 ml of distilled water at a pH of 6.8) was used as the base for the experimental extenders. Each ejaculate was split into four equal aliquots and diluted using both of the T extenders with G (5% or 7%) or EG (3% or 5%). After that, each extenders were split into three equal aliquots and diluted using both of 25 mM sucrose (S) or 5 mM dithiothreitol (D), and control (without additives) to a final concentration of approximately 60×10^6 spermatozoa per ml (15×10^6 sperm in each 0.25 ml straw). Diluted semen samples cooled down to 4 °C in 4 h. After that, frozen at a programmed rate of -3 °C/min from +4 to -10 °C; -40 °C/min from -10 to -100 °C; and -20 °C/min from -100 to -140 °C in a digital freezing machine (Digitcool 5300 ZB 250, IMV, France). Thereafter, the straws were plunged into liquid

nitrogen. After at least 24 h, the frozen straws were thawed in a 37 °C water bath for 20 s immediately before use.

Post-thaw semen evaluation

Analysis of microscopic sperm parameters

A computer-assisted sperm motility analysis (CASA; IVOS version 12; Hamilton-Thorne Biosciences, Beverly MA, USA) was also used to analyze sperm motion characteristics. CASA was pre-adjusted for bovine sperm analysis. Thawed semen was diluted (5 μ l semen + 95 μ l extender) in a Tris based extender (without egg yolk and cryoprotectant) and evaluated immediately after dilution. A 4- μ l sample of diluted semen was put onto a prewarmed chamber slide (20 mm; Leja 4; Leja Products Luzernestraat B.V., Holland), and sperm motility characteristics were determined with a $\times 10$ objective at 37 °C. The following spermatological characteristics were recorded; CASA motility (%), progressive motility (%), VAP (average path velocity, μ m/s), VSL (straight linear velocity, μ m/s), VCL (curvilinear velocity, μ m/s), ALH (amplitude of lateral head displacement, μ m), LIN (linearity index; $LIN = [VSL/VCL] \times 100$), BCF (beat cross frequency, Hz), STR (Straightness, VSL/VAP). For each evaluation, 10 microscopic fields were analyzed to include at least 300 cells.

For the assessment of sperm abnormalities, at least three drops of each sample were added to Eppendorf tubes containing 1 ml Hancock solution [62.5 ml formalin (37%), 150 ml sodium saline solution, 150 ml buffer solution, and 500 ml double-distilled water] [35]. One drop of this mixture was put on a slide and covered with a coverslip. The percentages of total sperm abnormalities (acrosome and other abnormalities) were determined by counting a total of 200 spermatozoa under phase-contrast microscopy ($\times 1000$, oil immersion, Olympus BX43, Tokyo, Japan).

The hypo-osmotic swelling test (HOS test) was used to assess the functional integrity of the spermatozoal membranes. The HOS test was performed by incubating 30 μ l of semen with 300 μ l of a 100 mOsm hypo-osmotic solution at 37 °C for 60 min. After incubation, 0.2 ml of the mixture was spreading on a warm slide with a cover slip and then was examined using a phase-contrast microscope ($\times 400$, Olympus BX43, Tokyo, Japan) [31]. The number of swollen spermatozoa out of 200 was counted; swelling is characterized by a coiled tail, indicating that the plasma membrane is intact.

For the assessment of sperm DNA damage was investigated using the single cell gel electrophoresis (comet) assay by the method of previously described by Tuncer et al. [40]. Analysis of comet was performed at high alkaline conditions. After analysis, the images of 100 randomly chosen nuclei were analyzed using a fluorescent microscope ($\times 400$, Zeiss, Germany). Nucleotide DNA extends under electrophoresis to form “comet tails,” and the relative intensity of DNA in the tail images DNA break frequency. The percentage of the total DNA in the comet tail was taken as a measure of DNA break frequency. Tail DNA (%) was assessed in 100 cells by using Comet Assay III image analysis system (Perceptive Instruments, UK). Analysis was performed blindly by one slide reader.

Biochemical assays of post-thaw semen

Malondialdehyde (MDA) concentrations. Lipid peroxidation was estimated by the measuring of MDA in sperm samples by the method of previously described by Eken et al. [16]. After the reaction of MDA with thiobarbituric acid reactive substances (TBARS), the reaction product was followed spectrophotometrically at 532 nm, using tetramethoxypropane as a standard. The results were expressed as nmol/ml.

GPx activity. GPx activity was measured as previously described by Tüzün et al. [41]. Briefly, a reaction mixture containing 1 mmol/l

Download English Version:

<https://daneshyari.com/en/article/10928182>

Download Persian Version:

<https://daneshyari.com/article/10928182>

[Daneshyari.com](https://daneshyari.com)