

Two HAP2-GCS1 homologs responsible for gamete interactions in the cellular slime mold with multiple mating types: Implication for common mechanisms of sexual reproduction shared by plants and protozoa and for male-female differentiation

Marina Okamoto, Lixy Yamada, Yukie Fujisaki, Gareth Bloomfield, Kentaro Yoshida, Hidekazu Kuwayama, Hitoshi Sawada, Toshiyuki Mori, Hideko Urushihara



PII: S0012-1606(16)30094-X
DOI: <http://dx.doi.org/10.1016/j.ydbio.2016.05.018>
Reference: YDBIO7125

To appear in: *Developmental Biology*

Received date: 19 February 2016
Revised date: 23 April 2016
Accepted date: 13 May 2016

Cite this article as: Marina Okamoto, Lixy Yamada, Yukie Fujisaki, Gareth Bloomfield, Kentaro Yoshida, Hidekazu Kuwayama, Hitoshi Sawada, Toshiyuki Mori and Hideko Urushihara, Two HAP2-GCS1 homologs responsible for gamete interactions in the cellular slime mold with multiple mating types: Implication for common mechanisms of sexual reproduction shared by plants and protozoa and for male-female differentiation, *Developmental Biology*, <http://dx.doi.org/10.1016/j.ydbio.2016.05.018>

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting galley proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

Two HAP2-GCS1 homologs responsible for gamete interactions in the cellular slime mold with multiple mating types: Implication for common mechanisms of sexual reproduction shared by plants and protozoa and for male-female differentiation

Marina Okamoto ^a, Lixy Yamada ^b, Yukie Fujisaki ^a, Gareth Bloomfield ^c, Kentaro Yoshida ^a, Hidekazu Kuwayama ^a, Hitoshi Sawada ^b, Toshiyuki Mori ^d, and Hideko Urushihara ^{a#}

^a Faculty of Life and Environmental Sciences, University of Tsukuba, 1-1-1 Tennodai, Tsukuba, Ibaraki 305-8572, Japan

^b Sugashima Marine Biological Laboratory, Graduate School of Science, Nagoya University, 429-63 Sugashima, Toba, Mie 517-0004, Japan

^c Medical Research Council Laboratory of Molecular Biology, Francis Crick Avenue, Cambridge Biomedical Campus, Cambridge, CB2 0QH, UK

^d Department of Biological Sciences, Graduate School of Science, The University of Tokyo, 7-3-1 Hongo, Bunkyo-ku, Tokyo 113-0033, Japan

Running title: HAP2-GCS1 homologs in Dictyostelium

[#] To whom correspondence should be addressed.

Hideko Urushihara

Faculty of Life and Environmental Sciences, University of Tsukuba

1-1-1 Tennodai, Tsukuba, Ibaraki 305-8572, Japan

Tel.: +81-29-853-4664

Fax: +81-29-853-6614,

E-mail: hideko@biol.tsukuba.ac.jp

Download English Version:

<https://daneshyari.com/en/article/10931143>

Download Persian Version:

<https://daneshyari.com/article/10931143>

[Daneshyari.com](https://daneshyari.com)