



Contents lists available at [ScienceDirect](#)

Vaccine

journal homepage: www.elsevier.com/locate/vaccine



Maternal immunization in Argentina: A storyline from the prospective of a middle income country

C. Vizzotti^a, S. Neyro^{a,*}, N. Katz^a, M.V. Juárez^a, M.E. Perez Carrega^a, A. Aquino^a, F. Kaski Fullone^b

^a Dirección Nacional de Control de Enfermedades, Ministerio de Salud de la Nación, Argentina

^b Secretaría de Promoción y Programas Sanitarios, Ministerio de Salud de la Nación, Argentina

ARTICLE INFO

Article history:

Received 27 April 2015
Received in revised form 10 July 2015
Accepted 28 July 2015
Available online xxx

Keywords:

Vaccination
Pregnancy
Flu
Tdap
Whooping cough
Vaccination coverage
Health policies

ABSTRACT

The importance of vaccination during pregnancy lies not only in directly protecting vaccinated women, but also by indirectly protecting small infants during the first few months of life. Vaccination against the flu and whooping cough is a priority within the comprehensive care strategy for pregnant women and small infants in Argentina, in the context of transitioning from child vaccination to family vaccination.

In 2011, the flu vaccine was included in the National Immunization Schedule (NIS) as mandatory and free of charge, with the aim of decreasing complications and death due to influenza in the at-risk population in Argentina. The national vaccination coverage attained in pregnant women in the past 4 years (2011–2014) has been satisfactory; 88% coverage was attained in the year this program was introduced to the schedule. In the following years, coverage was maintained at greater than 95%. In February 2012, Argentina became the first country in Latin America to have universal vaccination strategy for pregnant women against whooping cough. This recommendation was implemented throughout the country by vaccination with the diphtheria toxoid, tetanus toxoid, and acellular pertussis (Tdap) vaccine starting at 20 weeks of pregnancy, with the aim of decreasing morbimortality due to whooping cough in infants under 6 months of age. The vaccine was incorporated into the NIS in 2014. More than 1,200,000 doses were applied in this period. Both vaccines showed a suitable safety profile and no serious events were reported.

Argentina is an example of a middle-income country that has been able to implement a successful strategy for primary prevention through vaccines, making it a health policy.

© 2015 The Authors. Published by Elsevier Ltd. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

1. Introduction

The Republic of Argentina is part of the Region of the Americas. Located at the southeastern part of the American continent, its continental surface is around 2,780,000 km² and it extends 3800 km from north to south and 1425 km from east to west. The territory is quite diverse, including mountainous regions, plateaus, and plains, with extensive climatic and cultural diversity. Organized as a republican, representative, and federal system, it is composed of 24 jurisdictions: 23 provinces and one autonomous city. There are more than 40 million inhabitants [1] and more than 750,000 births are reported each year [2].

Argentina is a middle-income country in which the primary prevention strategy, by way of vaccines, was established as a priority in public health, comprising a policy of state in the last decade.

The health system is composed of three subsectors: public, private, and social security. Nevertheless, all inhabitants in the country are guaranteed universal and free access to health by means of the public system and through the national government. The system guarantees all vaccines included in the NIS, as overseen by the Argentinean State at the more than 8000 public vaccination centers located throughout the country.

The evolution and growth of the NIS in this period has made it one of the most modern and comprehensive in the region [3], with 19 free and mandatory vaccines available universally for all stages of life and in all sectors of the population. It is therefore a robust indicator of social equity and inclusion.

This vision of public health prioritizes the strategy for primary prevention through vaccines, which is reflected by the incorporation of 13 of the 19 vaccines as part of the schedule, and an over

* Corresponding author. Tel.: +54 1143427388.

E-mail addresses: sneyro@dinacei.msal.gov.ar, silvinaneyro@gmail.com (S. Neyro).

<http://dx.doi.org/10.1016/j.vaccine.2015.07.109>

0264-410X/© 2015 The Authors. Published by Elsevier Ltd. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

17-fold increase in the national budget earmarked for vaccination in the last decade.

One of the most important challenges in this period has been the transition from child vaccination to family vaccination, in which the vaccination of pregnant women is a cornerstone. Part of the challenge was the inclusion of the obstetricians and health personnel who take care of pregnant women and the newborns as a part of the strategy. The epidemiological circumstances faced before the start of the strategy of maternal immunization, reflected in the mortality of pregnant women with influenza during 2009–2010 and young infants during the outbreak of pertussis in 2011, aware health personnel and general population, attaching high strategy adherence. This allowed turning a potential threat into an opportunity for reaching satisfactory vaccine coverages.

As pregnancy vaccination takes place at primary care facilities which belong to the Expanded Program on Immunization, and not at the obstetrician office, it is offered to all women even if they did not achieve the obstetrician recommendation.

The importance of vaccination during pregnancy lies not only in directly protecting vaccinated women, but also by indirectly protecting small infants during the first few months of life. This is based on passive immunization generated by the transplacental transfer of maternal antibodies to infants who cannot be vaccinated due to their chronological age. Therefore, flu and whooping cough vaccination is a priority within the comprehensive care strategy of pregnant women and small infants in Argentina.

Safety monitoring in vaccines is also a paramount objective in Argentina. The notification of adverse events following immunization (AEFI) is performed passively. This implies that cases must be spontaneously reported by health effectors. The final evaluation and classification of serious AEFI is performed by the Comisión Nacional de Seguridad en Vacunas (CoNaSeVa) [National Commission for Safety in Vaccines]. It is composed of external consultants who belong to national scientific societies such as the Argentine Society of Pediatrics, the Argentine Society of Infectious Diseases, the Argentine Society for Pediatric Infectious Diseases, the national regulatory agency – Administración Nacional de Medicamentos, Alimentos y Tecnología Médica (ANMAT) – [National Administration of Drugs, Food and Medical Technology], the Pan American Health Organization (PAHO), and representatives of the Dirección Nacional de Control de Enfermedades Inmunoprevenibles (DiNaCEI) [National Immunization Program].

2. Development

2.1. Influenza

Pregnant women infected by the influenza virus present a greater risk of serious progression, hospitalization, and death compared to the rest of the healthy adult population. The product of conception can also be affected, generating intrauterine growth restriction, premature birth, fetal and/or neonatal death [4–7].

Flu vaccination in pregnant women has been shown to be safe and effective in providing protection not only to the woman, but also to the fetus and infant [8,9], through the passive transfer of antibodies through the placenta [10,11].

The available scientific evidence sustains the safety of flu vaccination [12,13] as a preventative measure during pregnancy and shows the cost-effectiveness of the strategy [14] in terms of flu prevention and its complications, both in pregnancy and in the product of conception. The PAHO establishes pregnant women as a priority group among the population in whom flu vaccination is aimed.

2.1.1. Experience in Argentina

The analysis of 332 total deaths occurring in Argentina during the influenza A (H1N1) pandemic in 2009 evidenced that 6%

(20/332) of the deaths occurred in pregnant women and women in the immediate postpartum. Of these, only 47% presented a comorbidity that would determine a greater risk of complications due to the flu, unlike 70% of the non-pregnant women who died [15]; this highlights the vulnerability generated by the condition of being pregnant itself in infection by the flu virus.

The high number of pregnant and post-partum women who died is consistent with international reports that identify pregnancy as an independent risk factor for serious disease and death due to influenza [7,15,16]. The increased vulnerability of this population, demonstrated nationally and internationally [6,15], led to the subsequent definition of a sustained preventative strategy, through the implementation of a free and mandatory flu vaccine for at-risk groups, including pregnant women.

In 2010, the largest national flu vaccination campaign in the history of our country was carried out, with the objective of decreasing mortality due to the influenza A (H1N1) virus; it included 12 million adjuvant monovalent vaccine doses. This strategy was aimed at pregnant women in any trimester of pregnancy, postpartum women, health personnel, children 6–59 months of age and people aged 5–64 years with risk factors (heart, kidney and chronic respiratory diseases, immunocompromised, diabetes, morbid obesity, etc.), and people over 65 years of age [17]. Vaccination coverage in pregnant women that year reached 98.1% (total number of doses applied: 444,397), with an average national coverage for all risk groups of 93.4%. Eighty-five percent (85%) of the target population was vaccinated at least 14 days prior to the start of the winter season. In 2010, there were no cases of death due to confirmed flu based on real time or conventional chain polymerase reaction (CPR), reported in patients belonging to the aforementioned risk groups.

In 2011, under the framework of a comprehensive prevention strategy in respiratory diseases, Argentina incorporated the flu vaccine into the NIS, as free and mandatory for all vulnerable groups (children aged 6–24 months, postpartum women, health personnel, people aged 2–64 years of age with risk factors, and people over the age of 65), including pregnant women in any trimester of pregnancy, with the objective of decreasing complications, hospitalizations, and deaths due to the influenza virus [3,18].

Since its incorporation as part of the NIS in 2011 and through 2014, 1,422,059 flu vaccine doses have been given to pregnant women. The flu vaccines, overseen by the national Ministry of Health in this time period and used in this population, were Agrippal® S1 (in 2011 and 2012) and Virafllu® (in 2013 and 2014); both are inactivated, trivalent vaccines without adjuvants.

The national vaccination coverage attained in this group in the past 4 years (2011–2014) has been satisfactory. Eighty-eight percent (88%) coverage was achieved in the year this program was introduced to the schedule and in the following years, coverage was maintained at greater than 95% [19] (Graph 1). National Immunization Program estimates flu vaccination coverages in pregnant women based on a 50% cohort of live newborns as denominator.

During the 2011–2014 period, 10 AEFI were reported (rate: 0.7/100,000 doses applied [d.a.]), of which 4 were mild events associated with the vaccine (rate: 0.28/100,000 d.a.) and 5 were program errors (rate: 0.35/100,000 d.a.). One case of ascending acute flaccid paralysis was reported and classified as non-conclusive due to the lack of complementary studies. No deaths were recorded due to influenza in promptly vaccinated pregnant women, nor were there any serious adverse events reported in relation to vaccination (Table 1).

2.2. Whooping Cough

Whooping cough is an acute contagious and infectious disease that is vaccine-preventable; it essentially compromises the

Download English Version:

<https://daneshyari.com/en/article/10963441>

Download Persian Version:

<https://daneshyari.com/article/10963441>

[Daneshyari.com](https://daneshyari.com)