



## Short communication: Viable *Mycobacterium avium* subspecies *paratuberculosis* in retail artisanal Coalho cheese from Northeastern Brazil

A. C. S. Faria,\* D. G. G. Schwarz,† I. A. Carvalho,† B. B. Rocha,† K. N. De Carvalho Castro,‡ M. R. Silva,§ and M. A. S. Moreira†<sup>1</sup>

\*Faculdade de Ciências Biológicas e da Saúde (FACISA/UNIVIÇOSA), Av. Maria de Paula Santana, 3815, Silvestre, CEP 36570-000, Viçosa, Minas Gerais, Brazil

†Departamento de Medicina Veterinária, Universidade Federal de Viçosa (UFV), Av. P. H. Rolfs, s/n, Campus Universitário, CEP 36570-900, Viçosa, Minas Gerais, Brazil

‡Empresa Brasileira de Pesquisa Agropecuária do Meio Norte/Uep Parnaíba (EMBRAPA), Br 343, km 35, Cep 64200-970, Parnaíba, Piauí, Brazil

§Empresa Brasileira de Pesquisa Agropecuária Gado de Leite (EMBRAPA), Rua Eugênio do Nascimento, 610, CEP 36038-330, Juiz de Fora, MG, Brazil

### ABSTRACT

*Mycobacterium avium* ssp. *paratuberculosis* (MAP) is the etiologic agent of paratuberculosis and it potentially plays a role in Crohn's disease. In humans, the main route of transmission of MAP might be the intake of contaminated milk and dairy products. Considering that MAP has already been detected in many types of cheese in different counties, and that Coalho cheese is an important dairy product in northeastern Brazil, the aim of this study was to report the first detection of MAP in retail Coalho cheese in Brazil by PCR and culture. Of 30 retail Coalho cheese samples, 3 (10%) amplified fragments of a similar size to that expected (626 bp) were obtained and viable MAP was recovered by culture from 1 (3.3%) sample. The DNA from the positive culture sample was sequenced and showed 99% identity with the insertion sequence IS900 deposited in GenBank. It was possible to identify the presence of MAP-specific DNA in the analyzed samples for the first time in Brazil, and to recover viable cells from retail Coalho cheese.

**Key words:** paratuberculosis, cheese, viable, culture

### Short Communication

*Mycobacterium avium* ssp. *paratuberculosis* (MAP) is the etiological agent of paratuberculosis that affects all ruminant species worldwide (Chiodini et al., 1984), and the spread of MAP occurs through feces, semen, and milk (Ayele et al., 2004; Slana et al., 2008). In Brazil, the presence of MAP DNA and the viability of MAP in raw and retail pasteurized milk (Carvalho et al., 2009, 2012) has been proven, suggesting that live

organisms might be transmitted to humans by ingestion of contaminated products. Artisanal Coalho cheese is a popular dairy product widely consumed in the northeastern region of Brazil, being produced with raw or pasteurized milk, added to rennet, and subjected to cooking. According to Brazilian standards, Coalho cheese is categorized as a semi-hard cheese, presenting medium to high moisture, and a solids content ranging from 35 to 60% (Brasil, 2001). Under Brazilian legislation, Coalho cheese is produced following milk coagulation using rennet or another suitable coagulating enzyme and is sometimes supplemented with specific lactic acid bacteria (Brasil, 2001). In terms of sensory characteristics, the cheese has a slightly salty and sour taste, and it is widely used as a "roast cheese" because of its heat resistance and resistance to melting. The deficient sanitary management of dairy herds, as well as inappropriate care in cheese processing, implies an increasing risk of contamination by many microorganisms that depreciates the product and endangers human health.

Considering the implications to public health and the quality of Brazilian artisanal cheese, this study aimed to evaluate the presence of MAP in retail Coalho cheese in the region of Parnaíba, Piauí State, in northeastern Brazil. From November 2011 to January 2012, 30 samples of artisanal Coalho cheese were randomly collected from formal (bakery, small market, and supermarket) and informal (street stall) traders in Parnaíba city, Piauí State. The samples were stored at  $-25^{\circ}\text{C}$  until analysis at the Bacterial Disease Laboratory of the Federal University of Viçosa (Minas Gerais, Brazil). The outside of each sample bag was decontaminated with 70% ethanol and exposed to room decontamination with UV radiation for 30 min. Each sample package was then aseptically opened and 30 g was weighed and transferred to a new sterile Whirl-Pak bag (Nasco, Fort Atkinson, WI) for culture and DNA extraction. A 1%

Received December 15, 2013.

Accepted March 20, 2014.

<sup>1</sup>Corresponding author: [masm@ufv.br](mailto:masm@ufv.br)

NaCl solution was preheated at 37°C for 30 min, and 125 mL was added to each bag, which were homogenized in a stomacher blender (Kleinfeld Labortechnik, Gehrden, Germany) for 2 min at 260 rpm. The homogenate was then incubated in a water bath at 37°C for 30 min, and 30 mL of the resultant suspension was transferred to a sterile falcon tube for DNA extraction. Another 30 mL of the same suspension was transferred to another sterile falcon tube for culture with 0.75% hexadecyl pyridinium chloride (**HPC**; Sigma Chemical Co., St. Louis, MO).

After transfer of 30 mL of the homogenate to a 50-mL falcon tube for DNA extraction, the solution was centrifuged at  $4,200 \times g$  for 15 min for the PCR assay, and the pellet was resuspended in 30 mL of 1% NaCl solution and centrifuged at  $4,200 \times g$  for 15 min. The pellet was washed in 1 mL of PBS and the supernatant was transferred to a 1.5-mL microtube and centrifuged at  $14,100 \times g$  for 10 min. Extraction of bacterial genomic DNA from cheese samples was performed using a Wizard Genomic DNA Purification Kit (Promega, Madison, WI), according to the manufacturer's instructions. The DNA was eluted in a final volume of 30  $\mu$ L of MilliQ water (Millipore, Billerica, MA), and its concentration was determined using a NanoDrop Lite spectrophotometer (Thermo Scientific, Waltham, MA). For PCR reactions, the PCR kit GoTaq Green Master Mix (Promega) was used according to the manufacturer's instructions. Reactions were performed using the BN1 (5'-GTT ATT AAC GAC GCC CAG C-3') and BN2 (5'-ACG ATG CTG TGT TGG GCG TTA G-3') primers that amplify a 626-bp fragment, based on the insertion sequence *IS900*, according to Sivakumar et al. (2005). Briefly, 12.5  $\mu$ L of mix, 1  $\mu$ L of each oligonucleotide, 6.5  $\mu$ L of MilliQ water, and 4  $\mu$ L of extracted DNA at a concentration of 200 ng/ $\mu$ L were used in a total reaction volume of 25  $\mu$ L. Amplifications were carried out in a thermocycler (MaxyGene II Thermal Cycler, Corning Inc., Corning, NY) using an initial denaturation at 94°C for 4 min, followed by 30 cycles of denaturation at 94°C for 1 min, annealing at 60°C for 1 min, and extension at 72°C for 1 min, with a final extension at 72°C for 4 min. Amplicons were analyzed by gel electrophoresis in 1% (wt/vol) agarose gels (Invitrogen, Washington, DC) in 0.5 tris-borate-EDTA buffer stained with 1  $\mu$ L of GelRed (diluted 1:500; Biotium Inc., Hayward, CA) using UV radiation in an Eagle Eye II transilluminator (Stratagene, La Jolla, CA). The K-10 strain of MAP was used as the positive control, and ultrapure water was used as a negative control. A 100-bp DNA ladder (Promega) was used as a molecular weight standard.

A further 30 mL of the homogenate was transferred to a 50-mL sterile falcon tube for culture and centri-

fuged at  $2,500 \times g$  for 15 min. The supernatant was discarded and the pellet resuspended in 10 mL of 0.75% HPC in a dark room at room temperature for 5 h, as previously described (Ikonomopoulos et al., 2005). Subsequently, samples were centrifuged at  $2,500 \times g$  for 15 min and the supernatant was discarded. Finally, the remaining pellet was resuspended and washed with 1 mL of PBS, pH 7.2, and a 250- $\mu$ L aliquot of the suspension was inoculated onto 2 Herrold's egg yolk medium (**HEYM**) slopes with or without 2  $\mu$ L/mL of mycobactin *J*. After growth, the colonies were confirmed by *IS900* PCR and the amplified fragments were extracted and purified from agarose gels using the Wizard SV Gel kit and PCR Clean-Up System (Promega), according to the manufacturer's instructions. Both strands were sequenced in triplicate and the sequences obtained were aligned, edited, and compared with other sequences deposited in GenBank using BLAST (Basic Local Alignment Search Tool) software (<http://blast.ncbi.nlm.nih.gov/Blast.cgi>), available from the National Center for Biotechnology Information. Isolates grown in HEYM were collected, heat-fixed onto slides, and stained with the Ziehl-Neelsen (ZN) kit (Laborclin, Pinhais, Brazil) according to the manufacturer's instructions. Samples were then checked by optical microscopy for the presence of acid-fast bacilli.

The detection of MAP-specific DNA was observed in 10% (3/30) of Coalho cheese samples. Although PCR techniques sensitively detect the presence of the microorganism at certain points in cheese production, they do not distinguish living cells from dead cells. The detection of MAP has been shown in cheese produced by raw milk (Spahr and Schaefroth, 2001; Stephan et al., 2007) as well as pasteurized milk (Sung and Collins, 2000; Donaghy et al., 2004; Clark et al., 2006) and it was shown that cheese ripening significantly decreases the viability of MAP. However, the efficiency of maturation is related to the degree of contamination of the raw milk used and to the efficiency of pasteurization. In general, using pasteurized milk and a 60-d curing period in cheese manufacture decreases the amount of initial MAP (Sung and Collins, 2000). In relation specifically to the Brazilian artisanal Coalho cheese, information is available regarding the presence and resistance of microorganisms in products produced from raw or pasteurized milk. However, the antimicrobial activity of the water-soluble peptide extracts, due to the proteolytic activity of the cheese process, recently showed that artisanal Coalho cheese can be considered a "functional food." Antimicrobial activity was found against *Enterococcus faecalis*, *Bacillus subtilis*, *Escherichia coli*, and *Pseudomonas aeruginosa* (Silva et al., 2012), but was not evaluated against *Mycobacterium* spp. The MAP viability in this type of cheese is likely

Download English Version:

<https://daneshyari.com/en/article/10976276>

Download Persian Version:

<https://daneshyari.com/article/10976276>

[Daneshyari.com](https://daneshyari.com)